

# IOWA PRENATAL CARE BARRIERS PROJECT DATA FROM 2015 SURVEYS

2015 Respondents from All Hospitals in Iowa: 23,212

2015 Respondents with Babies Born in **Hardin**: 111

<b>Demographic Indicators</b>	<b>State</b>	<b>County</b>
<b>Mother's Age</b>		
Less than 18 years	01%	<01%
18-19 years	04%	<01%
20-25 years	27%	30%
26-30 years	35%	35%
31-35 years	25%	25%
More than 35 years	08%	10%
<b>Mother's Ethnicity</b>		
Hispanic	07%	05%
Non-Hispanic	93%	95%
White	91%	01%
Black	05%	97%
Asian or Pacific Islander	03%	01%
American Indian or Native Alaskan	01%	01%
<b>Socioeconomic Indicators</b>		
<b>Mother's Education</b>		
Less than high school	07%	04%
High school	20%	12%
Some college	35%	49%
College graduate (Bachelor's degree)	27%	27%
Graduate school	11%	08%
<b>Household Income</b>		
Less than \$10,000	14%	07%
\$10,000 - \$19,999	10%	07%
\$20,000 - \$29,999	09%	13%
\$30,000 - \$39,999	09%	11%
\$40,000 - \$49,999	07%	07%
\$50,000 or more	52%	54%

<b>Information Received in 12 Months Prior to Becoming Pregnant</b>	<b>State</b>	<b>County</b>
---------------------------------------------------------------------	--------------	---------------

<b>Health Care Provider Talked About:</b>		
-------------------------------------------	--	--

Birth control	50%	47%
Timing of pregnancy	20%	18%
Spacing of children	11%	08%

<b>Doctor, Nurse, or Other Health Care Provider Talked About:</b>		
-------------------------------------------------------------------	--	--

Taking folic acid or a multivitamin	45%	41%
Smoking	32%	26%
Drinking alcohol	27%	19%
Your weight	25%	22%
Regular exercise	28%	22%
How romantic relationships affect your health	09%	10%
Did not see a doctor, nurse, or other health care professional for a check-up in the 12 months prior to becoming pregnant	24%	24%

<b>Behavioral Indicators During Pregnancy</b>	<b>State</b>	<b>County</b>
-----------------------------------------------	--------------	---------------

<b>No Birth Control was Used at Time of Conception</b>	86%	89%
--------------------------------------------------------	-----	-----

<b>Unintended Pregnancy and Did NOT Use Birth Control</b>	67%	73%
-----------------------------------------------------------	-----	-----

<b>Cigarettes Smoked During Last 3 Months of Pregnancy</b>		
------------------------------------------------------------	--	--

Not smoked in past 2 years	76%	74%
Zero cigarettes per day	54%	59%
1-10 per day	40%	34%
11-20 per day	06%	07%
More than 20 per day	<01%	<01%

<b>Alcoholic Drinks During Last 3 Months of Pregnancy</b>		
-----------------------------------------------------------	--	--

Not had any alcoholic drinks in past 2 years	30%	24%
Zero drinks per week	98%	96%
1 or more drinks per week	02%	04%

<b>Health Care Services During Pregnancy</b>	<b>State</b>	<b>County</b>
<b>Visited a Dentist, Dental Clinic, or Received Dental Care</b>	55%	52%
<b>Health Care Professional or Other Agency Staff Person Came to Home to Help Care for You or for Your New Baby</b>	03%	02%
<b>Diagnosed with Depression During Pregnancy</b>		
Yes	02%	02%
No	90%	92%
No, Diagnosed with depression before becoming pregnant	08%	06%
<b>Took Prescription Medication for Depression</b>	43%	67%
<b>Received Counseling for Depression During Pregnancy</b>	26%	33%
<b>HIV/AIDS Testing During Pregnancy</b>		
Yes	45%	52%
No	30%	31%
No, Testing was offered, but did not want to be tested	02%	01%
Don't know	23%	16%
<b>Received Flu Shot During Pregnancy</b>		
Yes	63%	56%
No, Received flu shot before became pregnant	07%	09%
No, Did not want flu shot during pregnancy	12%	16%
No, Doctor said to wait until after pregnancy	01%	01%
No, Do not ever get flu shots	15%	16%
No, Flu shot was not available	02%	02%
<b>Hospital Services</b>	<b>State</b>	<b>County</b>
<b>Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby</b>		
Yes	84%	80%
No	14%	18%
Don't remember	03%	02%
<b>During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other Healthcare Professional Provided Information or Scheduled a Time to Discuss:</b>		
Breastfeeding or breastfeeding techniques	81%	77%
Action steps to decrease frustration when infant cries	41%	36%
How to lay the baby down to sleep	52%	36%
Dangers of shaking an infant	40%	35%
Did not receive any information and have not scheduled a time to discuss	07%	11%

*Technical Note:* Because percentages were rounded, the sum of values may not equal 100%.

*For more information:* Call the University of Northern Iowa Center for Social & Behavioral Research: (319) 273-2105.

*Data source:* Iowa Barriers to Prenatal Care Project, Iowa Department of Public Health, University of Northern Iowa Center for Behavioral Research: 2015 Data.