

IOWA PRENATAL CARE BARRIERS PROJECT DATA FROM 2014 SURVEYS

2014 Respondents from All Hospitals in Iowa: 24,696

2014 Respondents with Babies Born in Henry: 174

Demographic Indicators	State	County
Mother's Age		
Less than 18 years	01%	02%
18-19 years	04%	08%
20-25 years	28%	40%
26-30 years	35%	31%
31-35 years	25%	15%
More than 35 years	07%	03%
Mother's Ethnicity		
Hispanic	07%	06%
Non-Hispanic	93%	94%
White	91%	94%
Black	05%	04%
Asian or Pacific Islander	03%	01%
American Indian or Native Alaskan	01%	01%
Socioeconomic Indicators		
Mother's Education		
Less than high school	08%	13%
High school	19%	23%
Some college	36%	43%
College graduate (Bachelor's degree)	27%	16%
Graduate school	10%	04%
Household Income		
Less than \$10,000	14%	22%
\$10,000 - \$19,999	10%	13%
\$20,000 - \$29,999	10%	13%
\$30,000 - \$39,999	09%	12%
\$40,000 - \$49,999	08%	09%
\$50,000 or more	49%	32%

Information Received in 12 Months Prior to Becoming Pregnant	State	County
Doctor, Nurse, or Other Health Care Provider Talked About:		
Taking folic acid or a multivitamin	47%	38%
Smoking	37%	33%
Drinking alcohol	32%	29%
Your weight	30%	27%
Regular exercise	33%	27%
How romantic relationships affect your health	13%	12%
Did not see a doctor, nurse, or other health care professional for a check-up in the 12 months prior to becoming pregnant	34%	41%
Behavioral Indicators During Pregnancy		
No Birth Control was Used at Time of Conception	86%	86%
Unintended Pregnancy and Did NOT Use Birth Control	68%	70%
Plan to return to work outside of home following birth of this baby		
Yes	72%	71%
No	28%	29%
Does worksite have designated room/location (not counting bathroom stalls) for mothers to breastfeed or pump/express their breast milk		
Yes	48%	44%
No	32%	32%
Don't Know	19%	24%
Does worksite have written policy about breastfeeding?		
Yes	18%	14%
No	30%	35%
Don't Know	52%	50%
Cigarettes Smoked During Last 3 Months of Pregnancy		
Not smoked in past 2 years	77%	64%
Zero cigarettes per day	03%	18%
1-10 per day	13%	14%
11-20 per day	06%	03%
More than 20 per day	01%	01%
Alcoholic Drinks During Last 3 Months of Pregnancy		
Not had any alcoholic drinks in past 2 years	23%	37%
Zero drinks per week	26%	62%
1 drink per week	32%	01%
2 or more drinks per week	41%	00%

Prenatal Care	State	County
Mother received prenatal care as early in the pregnancy as wanted		
Yes	93%	90%
No	07%	10%
Where mother went for <u>most</u> of prenatal care		
Private doctor's office or clinic	81%	76%
Midwife's office or home office from a midwife	05%	04%
Public clinic	13%	20%
Other	01%	00%
HIV/AIDS Testing During Pregnancy		
Yes	50%	56%
No	25%	20%
No, Testing was offered, but did not want to be tested	04%	05%
Don't know	22%	19%
Received Flu Shot During Pregnancy		
Yes	62%	68%
No, Received flu shot before became pregnant	07%	05%
No, Did not want flu shot during pregnancy	13%	12%
No, Doctor said to wait until after pregnancy	01%	00%
No, Do not ever get flu shots	14%	13%
No, Flu shot was not available	02%	02%
Hospital Services		
Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby		
Yes	83%	80%
No	15%	18%
Don't remember	03%	01%
During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other Healthcare Professional Provided Information or Scheduled a Time to Discuss:		
Breastfeeding or breastfeeding techniques	78%	81%
Action steps to decrease frustration when infant cries	49%	55%
How to lay the baby down to sleep	56%	64%
Dangers of shaking an infant	48%	57%
Did not receive any information and have not scheduled a time to discuss	09%	06%

Technical Note: Because percentages were rounded, the sum of values may not equal 100%.

For more information: Call the University of Northern Iowa Center for Social & Behavioral Research: (319) 273-2105.

Data source: Iowa Barriers to Prenatal Care Project, Iowa Department of Public Health, University of Northern Iowa Center for Behavioral Research: 2014 Data.