## IOWA PRENATAL CARE BARRIERS PROJECT DATA FROM 2014 SURVEYS

2014 Respondents from All Hospitals in Iowa: 24,696 2014 Respondents with Babies Born in **Polk: 5,050** 

Demographic Indicators	State	County
Mother's Age		
Less than 18 years	01%	01%
18-19 years	04%	03%
20-25 years	28%	24%
26-30 years	35%	34%
31-35 years	25%	28%
More than 35 years	07%	10%
Mother's Ethnicity		
Hispanic	07%	08%
Non-Hispanic	93%	92%
White	91%	86%
Black	05%	08%
Asian or Pacific Islander	03%	05%
American Indian or Native Alaskan	01%	01%
Socioeconomic Indicators	State	County
Mother's Education		,
Less than high school	08%	07%
High school	19%	18%
		10/0
Some college	36%	31%
Some college College graduate (Bachelor's degree)	36% 27%	31% 32%
Some college College graduate (Bachelor's degree) Graduate school	36% 27% 10%	31% 32% 12%
College graduate (Bachelor's degree)	27%	32%
College graduate (Bachelor's degree) Graduate school	27%	32%
College graduate (Bachelor's degree) Graduate school  Household Income	27% 10%	32% 12%
College graduate (Bachelor's degree) Graduate school  Household Income Less than \$10,000	27% 10% 14%	32% 12% 13%
College graduate (Bachelor's degree) Graduate school  Household Income Less than \$10,000 \$10,000 - \$19,999	27% 10% 14% 10%	32% 12% 13% 09%
College graduate (Bachelor's degree) Graduate school  Household Income Less than \$10,000 \$10,000 - \$19,999 \$20,000 - \$29,999	27% 10% 14% 10% 10%	32% 12% 13% 09% 09%

Information Received in 12 Months Prior to Becoming Pregnant	State	County
Doctor, Nurse, or Other Health Care Provider Talked About:		
Taking folic acid or a multivitamin	47%	49%
Smoking	37%	35%
Drinking alcohol	32%	31%
Your weight	30%	30%
Regular exercise	33%	33%
How romantic relationships affect your health	13%	12%
Did not see a doctor, nurse, or other health care professional for a check-up	34%	28%
in the 12 months prior to becoming pregnant		
	<b>.</b> .	
Behavioral Indicators During Pregnancy	State	County
No Birth Control was Used at Time of Conception	86%	87%
Unintended Pregnancy and Did NOT Use Birth Control	68%	69%
Onintended (Tegriancy and Did NOT Ose Shan Control	0070	0370
Plan to return to work outside of home following birth of this baby		
Yes	72%	72%
No	28%	28%
Does worksite have designated room/location (not counting bathroom		
stalls) for mothers to breastfeed or pump/express their breast milk		
Yes	48%	54%
No Double Market	32%	29%
Don't Know	19%	17%
Does worksite have written policy about breastfeeding?		
Yes	18%	21%
No	30%	26%
Don't Know	52%	52%
Cigarettes Smoked During Last 3 Months of Pregnancy		
Not smoked in past 2 years	77%	77%
Zero cigarettes per day	03%	13%
1-10 per day	13%	09%
11-20 per day	06%	01%
More than 20 per day	01%	00%
Alcoholic Drinks During Last 2 Months of Prognancy		
Alcoholic Drinks During Last 3 Months of Pregnancy Not had any alcoholic drinks in past 2 years	23%	29%
Zero drinks per week	25% 26%	70%
1 drink per week	32%	01%
2 or more drinks per week	41%	00%
2 of more drinks per week	71/0	00/0

Prenatal Care	State	County
Mother received prenatal care as early in the pregnancy as wanted		
Yes	93%	95%
No	07%	08%
Where mother went for most of prenatal care		
Private doctor's office or clinic	81%	84%
Midwife's office or home office from a midwife	05%	08%
Public clinic	13%	07%
Other	01%	00%
HIV/AIDS Testing During Pregnancy		
Yes	50%	46%
No	25%	28%
No, Testing was offered, but did not want to be tested	04%	02%
Don't know	22%	23%
Received Flu Shot During Pregnancy		
Yes	62%	61%
No, Received flu shot before became pregnant	07%	08%
No, Did not want flu shot during pregnancy	13%	13%
No, Doctor said to wait until after pregnancy	01%	01%
No, Do not ever get flu shots	14%	15%
No, Flu shot was not available	02%	03%
Hospital Services	State	County
Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby		
Yes	83%	84%
No	15%	13%
Don't remember	03%	03%
During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other		
Healthcare Professional Provided Information or Scheduled a Time to		
Discuss: Breastfeeding or breastfeeding techniques	700/	79%
· ·	78% 40%	
Action steps to decrease frustration when infant cries	49%	41%
How to lay the baby down to sleep	56%	50%
Dangers of shaking an infant  Did not receive any information and have not scheduled a time to discuss	48%	40%
Did not receive any information and have not scheduled a time to discuss	09%	10%

Technical Note: Because percentages were rounded, the sum of values may not equal 100%. For more information: Call the University of Northern Iowa Center for Social & Behavioral Research: (319) 273-2105. Data source: Iowa Barriers to Prenatal Care Project, Iowa Department of Public Health, University of Northern Iowa

Center for Behavioral Research: 2014 Data.