## IOWA PRENATAL CARE BARRIERS PROJECT DATA FROM 2015 SURVEYS

2015 Respondents from All Hospitals in Iowa: 23,212 2015 Respondents with Babies Born in **Webster**: 380

| Demographic Indicators            | State | County |
|-----------------------------------|-------|--------|
| Mother's Age                      |       |        |
| Less than 18 years                | 01%   | 02%    |
| 18-19 years                       | 04%   | 07%    |
| 20-25 years                       | 27%   | 34%    |
| 26-30 years                       | 35%   | 38%    |
| 31-35 years                       | 25%   | 17%    |
| More than 35 years                | 08%   | 04%    |
| Mother's Ethnicity                |       |        |
| Hispanic                          | 07%   | 04%    |
| Non-Hispanic                      | 93%   | 96%    |
| White                             | 91%   | 89%    |
| Black                             | 05%   | 09%    |
| Asian or Pacific Islander         | 03%   | 01%    |
| American Indian or Native Alaskan | 01%   | 01%    |

| Socioeconomic Indicators             | State | County |
|--------------------------------------|-------|--------|
| Mother's Education                   |       |        |
| Less than high school                | 07%   | 09%    |
| High school                          | 20%   | 25%    |
| Some college                         | 35%   | 44%    |
| College graduate (Bachelor's degree) | 27%   | 16%    |
| Graduate school                      | 11%   | 06%    |
| Household Income                     |       |        |
| Less than \$10,000                   | 14%   | 27%    |
| \$10,000 - \$19,999                  | 10%   | 14%    |
| \$20,000 - \$29,999                  | 09%   | 09%    |
| \$30,000 - \$39,999                  | 09%   | 07%    |
| \$40,000 - \$49,999                  | 07%   | 06%    |
| \$50,000 or more                     | 52%   | 36%    |

| Information Received in 12 Months Prior to Becoming Pregnant  | State | County |
|---|-------|--------|
| Health Care Provider Talked About:  |       |        |
| Birth control   | 50%   | 56%    |
| Timing of pregnancy   | 20%   | 16%    |
| Spacing of children   | 11%   | 10%    |
| Doctor, Nurse, or Other Health Care Provider Talked About:  |       |        |
| Taking folic acid or a multivitamin   | 45%   | 34%    |
| Smoking   | 32%   | 37%    |
| Drinking alcohol  | 27%   | 27%    |
| Your weight   | 25%   | 25%    |
| Regular exercise  | 28%   | 25%    |
| How romantic relationships affect your health   | 09%   | 08%    |
| Did not see a doctor, nurse, or other health care professional for a check-up in the 12 months prior to becoming pregnant | 24%   | 27%    |

| Behavioral Indicators During Pregnancy              | State | County |
|---|-------|--------|
| No Birth Control was Used at Time of Conception     | 86%   | 86%    |
| Unintended Pregnancy and Did NOT Use Birth Control  | 67%   | 75%    |
| Cigarettes Smoked During Last 3 Months of Pregnancy |       |        |
| Not smoked in past 2 years                          | 76%   | 63%    |
| Zero cigarettes per day                             | 54%   | 54%    |
| 1-10 per day  | 40%   | 41%    |
| 11-20 per day                                       | 06%   | 04%    |
| More than 20 per day                                | <01%  | 01%    |
| Alcoholic Drinks During Last 3 Months of Pregnancy  |       |        |
| Not had any alcoholic drinks in past 2 years        | 30%   | 27%    |
| Zero drinks per week                                | 98%   | 99%    |
| 1 or more drinks per week                           | 02%   | 01%    |

|  | State      | County     |
|--|------------|------------|
| Visited a Dentist, Dental Clinic, or Received Dental Care  | 55%        | 47%        |
| Health Care Professional or Other Agency Staff Person Came to Home to<br>Help Care for You or for Your New Baby                            | 03%        | 06%        |
| Diagnosed with Depression During Pregnancy   |            |            |
| Yes  | 02%        | 04%        |
| No   | 90%        | 86%        |
| No, Diagnosed with depression before becoming pregnant   | 08%        | 10%        |
| Took Prescription Medication for Depression  | 43%        | 30%        |
| Received Counseling for Depression During Pregnancy  | 26%        | 32%        |
| HIV/AIDS Testing During Pregnancy  |            |            |
| Yes  | 45%        | 58%        |
| No   | 30%        | 19%        |
| No, Testing was offered, but did not want to be tested   | 02%        | 01%        |
| Don't know   | 23%        | 23%        |
| Received Flu Shot During Pregnancy   |            |            |
| Yes  | 63%        | 66%        |
| No, Received flu shot before became pregnant   | 07%        | 06%        |
| No, Did not want flu shot during pregnancy   | 12%        | 12%        |
| No, Doctor said to wait until after pregnancy  | 01%        | 01%        |
| No, Do not ever get flu shots  | 15%        | 11%        |
| No, Flu shot was not available   | 02%        | 04%        |
| Hospital Services  | State      | County     |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth  |            |            |
| of the Baby  | 0/10/      | 700/       |
| Yes<br>No  | 84%<br>14% | 78%<br>20% |
| Don't remember   | 03%        | 02%        |
| Don tremember  | 0378       | 0270       |
| During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other<br>Healthcare Professional Provided Information or Scheduled a Time to |            |            |
| Discuss:   |            |            |
| Breastfeeding or breastfeeding techniques  | 81%        | 79%        |
| Action steps to decrease frustration when infant cries   | 41%        | 42%        |
| •  | 52%        | 51%        |
| How to lay the baby down to sleep  |            |            |
| How to lay the baby down to sleep<br>Dangers of shaking an infant  | 40%        | 42%        |

Technical Note: Because percentages were rounded, the sum of values may not equal 100%. For more information: Call the University of Northern Iowa Center for Social & Behavioral Research: (319) 273-2105. Data source: Iowa Barriers to Prenatal Care Project, Iowa Department of Public Health, University of Northern Iowa Center for Behavioral Research: 2015 Data.