County: Washington											
	Percent who smoked (2-yr avg)			Number who smoked (2-yr avg)			(2-yr avg)		who	Percent and number who quit (3-yr avg)	
Years	3 months prepreg.	3rd trimester	Any trimester	3 months prepreg.	3rd trimester	Any trimester	Total live births	Years	% Quit before 3rd trimester	# Quit before 3rd trimester	
2007-08	24.2	15.0	19.0	67	42	53	277	2007-09	39	24	
2013-14	17.6	10.7	13.2	56	34	42	319	2010-12	43	25	
2014-15	16.6	10.2	12.1	53	33	39	320	2013-15	41	22	

Percent and Number of New Moms Who Smoked, Who Quit Cigarettes, by County, 2007-15

Iowa											
	Perc	ent who smo	oked	Number who smoked					Percent and a	number who	
		(2-yr avg)		(2-yr avg)			(2-yr avg)		quit (3-yr avg)		
									% Quit	# Quit	
	3 months	3rd	Any	3 months	3rd	Any	Total		before third	before third	
Years	prepreg.	trimester	trimester	prepreg.	trimester	trimester	live births	Years	trimester	trimester	
2007-08	24.0	14.6	18.3	9,722	5,921	7,409	40,528	2007-09	40	3,803	
2013-14	20.1	11.7	15.0	7,912	4,610	5,891	39,349	2010-12	41	3,309	
2014-15	19.1	11.1	14.2	7,543	4,398	5,609	39,576	2013-15	42	3,276	

-Pre-pregnancy smoking: Smoking in three months before becoming pregnant

-Third trimester smoking: Smoking during the last three months of pregnancy

-Any trimester smoking: Smoking at any time while pregnant

-Quit rate: Smoked in the three months before pregnancy or in first or second trimester, quit by start of third trimester


Due to small numbers in some counties, two-year averages are shown for smoking prevalence and three-year averages for quit rates. Due to updates in the birth certificate databases, the counts here may vary slightly from those in the vital statistics reports published annually by the Iowa Department of Public Health.

Source: Iowa resident certificates of live birth, multi-year annual average rate, 2007-15


Iowa Department of Public Health, Division of Tobacco Use Prevention and Control, July 2017 https://idph.iowa.gov/tupac/statistics



Percent of new moms who smoked during pregnancy, 2014-15



Percent of new moms who smoked but quit by third trimester, 2013-15



- -Two-year average annual percent of new moms giving live birth who smoked cigarettes at any time during pregnancy, 2014-15.
- -Three-year average annual percent of new moms who smoked in the three months before pregnancy or during the first or second trimester but who quit by the start of the third trimester, 2013-15.



Source: Iowa certificates of live birth, 2013-15, Iowa Department of Public Health, July 2017.