WIC PNSS & PedNSS

Run Date: 2/7/2020

Table 3F

Page : 1

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Local Agency

Comparison of Racial and Ethnic Distribution (2)								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Agency 31	2,230	42.2	20.8	23.5	0.2	8.9	4.4	0.0
Agency 33	151	77.5	3.3	17.9	0.0	0.7	0.7	0.0
Agency 34	818	57.6	29.7	6.7	0.1	1.1	4.8	0.0
Agency 35	618	74.1	13.3	7.0	0.2	3.2	2.3	0.0
Agency 37	347	86.2	1.2	10.7	0.0	0.3	1.7	0.0
Agency 38	755	59.1	5.7	27.5	0.9	5.6	1.2	0.0
Agency 39	258	57.0	1.6	38.4	0.8	1.2	1.2	0.0
Agency 41	524	76.9	5.2	13.4	0.2	2.1	2.3	0.0
Agency 42	192	79.7	2.1	17.7	0.0	0.5	0.0	0.0
Agency 43	898	58.2	23.8	9.7	0.2	5.2	2.8	0.0
Agency 45	585	70.1	9.1	17.9	0.2	2.2	0.5	0.0
Agency 46	532	75.2	7.5	12.2	0.0	3.9	0.9	0.2
Agency 47	302	53.3	3.6	34.4	0.3	7.6	0.7	0.0
Agency 48	297	69.0	1.0	26.3	0.7	3.0	0.0	0.0
Agency 49	683	41.9	11.6	36.7	3.2	3.8	2.8	0.0
Agency 50	396	63.6	7.1	25.8	0.5	2.0	1.0	0.0
Agency 51	437	41.6	37.3	16.2	0.5	1.6	2.7	0.0
Agency 52	427	77.5	3.7	16.9	0.2	0.2	1.4	0.0
Agency 53	214	91.6	2.8	2.8	0.5	1.4	0.9	0.0
Agency 54	650	55.4	24.6	11.2	0.8	2.8	5.2	0.0
Iowa	**11,313**	**59.6**	**14.6**	**18.7**	**0.5**	**4.1**	**2.6**	**0.0**
National (Prior Year)	**133,781**	**53.3**	**21.7**	**17.1**	**1.1**	**2.8**	**3.5**	**0.4**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Local Agency

Comparison of Age Distribution (2)								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Agency 31	2,230	0.1	2.0	5.5	57.9	32.2	2.3	0.0
Agency 33	151	0.0	2.0	8.6	62.3	25.8	1.3	0.0
Agency 34	818	0.0	2.2	6.5	57.7	31.5	2.1	0.0
Agency 35	618	0.0	2.8	7.1	61.0	27.2	1.8	0.2
Agency 37	347	0.3	1.4	7.2	65.1	24.5	1.4	0.0
Agency 38	755	0.0	2.8	6.5	59.7	28.9	2.0	0.1
Agency 39	258	0.0	1.9	9.7	62.4	23.6	2.3	0.0
Agency 41	524	0.0	1.3	5.3	64.5	27.9	1.0	0.0
Agency 42	192	0.0	1.6	4.2	61.5	32.3	0.5	0.0
Agency 43	898	0.1	1.9	5.6	59.8	30.3	2.2	0.1
Agency 45	585	0.0	2.1	5.8	63.4	27.0	1.7	0.0
Agency 46	532	0.0	2.1	5.1	63.2	27.4	2.3	0.0
Agency 47	302	0.0	1.3	4.3	61.6	30.5	2.3	0.0
Agency 48	297	0.0	1.7	5.4	61.3	29.3	2.4	0.0
Agency 49	683	0.0	1.9	6.3	59.4	30.3	2.0	0.0
Agency 50	396	0.0	2.0	6.1	63.4	27.0	1.5	0.0
Agency 51	437	0.0	0.9	4.3	49.9	41.4	3.4	0.0
Agency 52	427	0.0	1.4	8.0	67.0	21.8	1.9	0.0
Agency 53	214	0.0	2.3	8.4	62.1	25.2	1.9	0.0
Agency 54	650	0.2	3.2	7.4	61.1	25.4	2.8	0.0
Iowa	**11,314**	**0.0**	**2.0**	**6.1**	**60.4**	**29.3**	**2.1**	**0.0**
National (Prior Year)	**133,790**	**0.1**	**2.7**	**8.0**	**62.0**	**25.2**	**1.9**	**0.1**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators (2,3)

Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agency 31	1,753	4.4 (14)	57.6 (1)	1,685	24.8 (16)	41.1 (4)	518	2.3 (5)	185	5.4 (7)	121	100.0 (1)	67	100.0 (1)
Agency 33	130	3.8 (10)	65.4 (18)	127	19.7 (3)	55.1 (20)	40	27.5 (13)	38	21.1 (12)	17	100.0 (1)	9	100.0 (1)
Agency 34	597	2.8 (5)	61.0 (10)	580	21.7 (7)	49.8 (16)	195	41.5 (19)	92	28.3 (17)	73	100.0 (1)	25	100.0 (1)
Agency 35	486	3.5 (8)	63.2 (14)	469	22.6 (11)	47.5 (14)	159	43.4 (20)	57	38.6 (20)	33	100.0 (1)	20	100.0 (1)
Agency 37	261	5.4 (19)	61.3 (11)	256	25.8 (17)	40.6 (3)	80	35.0 (18)	58	32.8 (18)	32	100.0 (1)	7	100.0 (1)
Agency 38	567	4.2 (12)	60.1 (8)	543	27.1 (19)	41.6 (5)	148	14.9 (12)	83	22.9 (14)	55	100.0 (1)	9	100.0 (1)
Agency 39	184	4.3 (13)	59.2 (6)	180	23.3 (13)	38.9 (2)	54	1.9 (2)	81	2.5 (3)	16	100.0 (1)	2	100.0 (1)
Agency 41	403	3.2 (7)	70.5 (20)	391	19.2 (2)	46.8 (13)	94	5.3 (9)	85	5.9 (8)	48	100.0 (1)	20	100.0 (1)
Agency 42	147	4.8 (17)	61.9 (12)	141	17.7 (1)	51.8 (19)	55	29.1 (15)	53	37.7 (19)	16	100.0 (1)	9	100.0 (1)
Agency 43	695	3.6 (9)	60.0 (7)	671	21.3 (6)	49.9 (17)	173	3.5 (6)	99	1.0 (2)	43	100.0 (1)	40	100.0 (1)
Agency 45	440	4.5 (15)	59.1 (5)	421	21.1 (5)	49.2 (15)	150	4.0 (7)	65	4.6 (5)	56	100.0 (1)	36	100.0 (1)
Agency 46	438	5.5 (20)	60.3 (9)	413	20.6 (4)	43.6 (7)	166	27.7 (14)	74	27.0 (16)	6	100.0 (1)	16	100.0 (1)
Agency 47	195	2.1 (4)	63.1 (13)	180	26.1 (18)	45.6 (9)	67	0.0 (1)	43	4.7 (6)	32	100.0 (1)	15	100.0 (1)
Agency 48	248	2.0 (3)	63.3 (15)	243	28.8 (20)	42.8 (6)	92	2.2 (4)	73	2.7 (4)	2	100.0 (1)	13	100.0 (1)
Agency 49	525	1.3 (1)	63.8 (16)	510	22.4 (10)	45.7 (10)	190	4.7 (8)	164	6.1 (9)	37	100.0 (1)	15	100.0 (1)
Agency 50	327	3.1 (6)	64.8 (17)	319	23.5 (14)	44.2 (8)	104	1.9 (2)	18	0.0 (1)	18	100.0 (1)	10	100.0 (1)
Agency 51	309	4.5 (15)	58.6 (3)	289	23.5 (14)	38.4 (1)	113	33.6 (16)	130	22.3 (13)	21	100.0 (1)	16	100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Local Agency

Comparison of Maternal Health Indicators (2,3)

Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agency 52	313	1.9 (2)	65.5 (19)	300	23.0 (12)	50.7 (18)	112	8.0 (11)	39	7.7 (10)	17	100.0 (1)	17	100.0 (1)
Agency 53	157	5.1 (18)	58.0 (2)	155	21.9 (8)	46.5 (12)	36	5.6 (10)	21	9.5 (11)	4	100.0 (1)	6	100.0 (1)
Agency 54	502	4.0 (11)	59.0 (4)	476	22.3 (9)	45.8 (11)	155	34.2 (17)	99	26.3 (15)	8	100.0 (1)	24	100.0 (1)
Iowa	**8,677**	**3.8**	**61.0**	**8,349**	**23.1**	**45.1**	**2,701**	**15.5**	**1,557**	**14.7**	**655**	**100.0**	**376**	**100.0**
National (Prior Year)	**112,231**	**4.0**	**60.3**	**89,229**	**22.3**	**47.7**	**23,550**	**28.2**	**52,310**	**18.4**	**96,728**	**10.0**	**95,873**	**9.4**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Local Agency

Comparison of Maternal Behavioral Indicators (2,3)									
Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31	1,451	27.8 (13)	13.3 (13)	1,960	0.5 (15)	652	16.1 (18)	858	88.5 (16)
Agency 33	113	32.7 (5)	8.0 (8)	146	0.0 (18)	42	19.0 (15)	43	93.0 (8)
Agency 34	570	25.4 (18)	8.1 (9)	699	0.6 (13)	361	15.2 (19)	491	90.8 (12)
Agency 35	467	36.6 (2)	10.7 (11)	503	1.4 (6)	358	20.4 (12)	423	92.0 (9)
Agency 37	239	32.2 (7)	7.1 (6)	296	0.0 (18)	128	25.8 (5)	127	90.6 (13)
Agency 38	447	32.0 (8)	8.7 (10)	600	1.5 (4)	332	19.3 (14)	346	87.9 (17)
Agency 39	143	27.3 (15)	7.0 (5)	234	1.3 (7)	68	23.5 (7)	75	96.0 (2)
Agency 41	392	38.3 (1)	15.1 (15)	450	0.4 (16)	345	20.9 (11)	403	87.3 (18)
Agency 42	134	26.9 (16)	14.2 (14)	161	1.2 (9)	139	23.0 (8)	142	91.5 (11)
Agency 43	448	31.7 (9)	17.4 (17)	748	0.8 (11)	308	25.3 (6)	433	90.1 (14)
Agency 45	437	29.7 (11)	4.3 (1)	477	1.0 (10)	403	20.3 (13)	447	94.4 (4)
Agency 46	319	34.2 (4)	25.1 (19)	463	1.7 (3)	40	22.5 (9)	41	82.9 (20)
Agency 47	182	27.5 (14)	7.1 (6)	237	1.3 (7)	180	47.8 (1)	190	94.7 (3)
Agency 48	189	24.9 (19)	6.3 (2)	274	0.4 (16)	31	32.3 (2)	48	93.8 (6)
Agency 49	460	30.2 (10)	6.5 (4)	614	0.8 (11)	79	31.6 (3)	107	94.4 (4)
Agency 50	295	26.1 (17)	25.4 (20)	344	1.5 (4)	233	17.6 (16)	286	84.3 (19)
Agency 51	259	18.1 (20)	18.5 (18)	367	1.9 (1)	94	21.3 (10)	107	91.6 (10)
Agency 52	288	35.1 (3)	6.3 (2)	373	0.0 (18)	118	16.9 (17)	120	89.2 (15)
Agency 53	126	32.5 (6)	15.9 (16)	158	0.6 (13)	88	27.3 (4)	93	93.5 (7)
Agency 54	420	29.5 (12)	13.1 (12)	569	1.9 (1)	24	8.3 (20)	25	100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Local Agency

Iowa	**7,379**	**29.9**	**12.1**	**9,673**	**0.9**	**4,023**	**21.3**	**4,805**	**90.3**
National (Prior Year)	**99,175**	**68.5**	**7.4**	**127,401**	**22.9**	**94,432**	**19.4**	**93,223**	**87.9**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2,3)

Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31	1,216	20.6 (2)	1,194	14.0 (6)	270	26.3 (10)	1,395	5.2 (6)	1,488	1.5 (10)
Agency 33	117	40.2 (19)	147	20.4 (15)	49	40.8 (2)	116	8.6 (15)	146	4.1 (17)
Agency 34	574	26.7 (8)	623	14.6 (7)	173	34.1 (4)	609	5.6 (7)	708	1.6 (11)
Agency 35	492	26.0 (7)	532	12.4 (5)	118	26.3 (10)	488	2.3 (1)	528	0.2 (2)
Agency 37	249	34.5 (16)	247	17.0 (12)	76	21.1 (15)	249	7.2 (11)	227	2.2 (13)
Agency 38	513	21.2 (5)	539	15.0 (8)	104	22.1 (14)	530	6.4 (9)	559	1.1 (6)
Agency 39	157	21.0 (3)	190	11.6 (3)	32	28.1 (8)	156	5.1 (5)	187	3.2 (16)
Agency 41	417	27.1 (9)	499	15.8 (9)	112	32.1 (6)	409	3.7 (3)	466	1.1 (6)
Agency 42	153	30.1 (12)	162	16.7 (11)	47	31.9 (7)	153	17.6 (20)	161	10.6 (20)
Agency 43	535	31.2 (13)	468	19.0 (14)	154	19.5 (16)	544	9.4 (17)	473	1.3 (8)
Agency 45	455	36.9 (17)	549	20.4 (15)	165	33.9 (5)	454	10.8 (19)	541	1.3 (8)
Agency 46	365	32.1 (14)	252	21.4 (18)	105	9.5 (20)	355	7.3 (12)	221	2.3 (14)
Agency 47	194	21.1 (4)	277	8.7 (2)	38	52.6 (1)	195	7.7 (13)	264	9.8 (19)
Agency 48	199	27.6 (10)	172	20.9 (17)	54	18.5 (17)	198	3.0 (2)	168	0.0 (1)
Agency 49	477	21.6 (6)	377	11.7 (4)	87	23.0 (13)	486	6.0 (8)	413	1.0 (5)
Agency 50	323	33.1 (15)	372	18.3 (13)	112	38.4 (3)	320	7.8 (14)	368	1.6 (11)
Agency 51	271	14.4 (1)	237	8.4 (1)	30	26.7 (9)	269	3.7 (3)	227	0.4 (3)
Agency 52	300	39.3 (18)	282	22.3 (19)	114	26.3 (10)	299	6.4 (9)	272	0.4 (3)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2,3)

Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 53	109	44.0 (20)	103	26.2 (20)	45	17.8 (18)	121	9.1 (16)	103	2.9 (15)
Agency 54	444	28.4 (11)	316	16.5 (10)	102	17.6 (19)	442	9.5 (18)	309	4.2 (18)
Iowa	**7,560**	**27.2**	**7,538**	**15.8**	**1,987**	**26.8**	**7,788**	**6.6**	**7,829**	**1.9**
National (Prior Year)	**108,764**	**27.1**	**119,789**	**14.3**	**24,778**	**36.6**	**97,290**	**10.0**	**120,070**	**3.5**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators (2,3)

Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31	2,068	9.0 (14)	7.6 (7)	1,895	9.3 (7)	1,697	4.1 (14)	2,098	77.6 (3)
Agency 33	143	7.0 (6)	6.3 (2)	142	7.0 (1)	132	3.0 (7)	143	70.6 (10)
Agency 34	743	7.1 (7)	9.4 (14)	676	8.1 (3)	599	3.7 (12)	773	71.5 (8)
Agency 35	586	8.9 (12)	6.7 (4)	485	9.7 (8)	438	4.8 (17)	590	56.8 (20)
Agency 37	330	5.8 (2)	9.4 (14)	293	9.9 (9)	264	1.9 (5)	329	69.9 (14)
Agency 38	708	7.8 (10)	10.3 (18)	582	8.9 (4)	522	3.1 (8)	716	74.9 (4)
Agency 39	245	7.3 (8)	6.5 (3)	230	10.9 (12)	203	3.4 (10)	247	85.0 (1)
Agency 41	492	7.7 (9)	10.4 (19)	439	11.2 (13)	388	3.1 (8)	494	70.2 (13)
Agency 42	184	9.8 (16)	10.9 (20)	158	8.9 (4)	144	4.9 (19)	184	73.9 (5)
Agency 43	847	8.9 (12)	9.2 (12)	730	11.5 (15)	642	3.7 (12)	853	70.5 (11)
Agency 45	560	8.4 (11)	7.1 (5)	464	10.3 (10)	415	4.1 (14)	561	64.5 (19)
Agency 46	492	5.3 (1)	8.1 (9)	438	7.3 (2)	406	1.2 (1)	492	70.3 (12)
Agency 47	282	6.7 (5)	9.2 (12)	234	12.4 (18)	204	1.5 (2)	283	73.5 (6)
Agency 48	288	6.6 (4)	5.6 (1)	274	13.1 (20)	238	1.7 (3)	288	64.6 (18)
Agency 49	635	6.5 (3)	7.2 (6)	601	12.1 (17)	512	1.8 (4)	653	68.3 (16)
Agency 50	374	9.6 (15)	8.3 (11)	338	12.4 (18)	296	2.0 (6)	374	70.9 (9)
Agency 51	397	10.8 (20)	8.1 (9)	349	8.9 (4)	314	4.8 (17)	401	84.8 (2)
Agency 52	394	10.2 (18)	7.6 (7)	362	11.3 (14)	319	5.0 (20)	399	72.7 (7)
Agency 53	209	10.0 (17)	9.6 (16)	158	10.8 (11)	141	3.5 (11)	208	64.9 (17)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8F

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators (2,3)

Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 54	613	10.4 (19)	9.6 (16)	553	11.9 (16)	486	4.1 (14)	614	68.6 (15)
Iowa	**10,590**	**8.3**	**8.4**	**9,401**	**10.2**	**8,360**	**3.4**	**10,700**	**71.7**
National (Prior Year)	**117,623**	**9.3**	**6.6**	**117,628**	**11.8**	**100,072**	**3.9**	**116,293**	**57.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Local Agency

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Local Agency	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Agency 31	6,995	43.4	19.5	24.2	0.3	8.1	4.5	0.0
Agency 33	567	80.6	3.0	14.6	0.0	1.1	0.7	0.0
Agency 34	2,630	59.0	26.7	7.4	0.1	1.9	4.8	0.0
Agency 35	1,903	76.2	11.5	7.1	0.3	2.8	2.1	0.0
Agency 37	865	85.7	1.4	11.8	0.0	0.5	0.7	0.0
Agency 38	2,429	60.5	4.0	26.2	1.3	6.4	1.5	0.0
Agency 39	792	59.1	1.3	37.0	0.6	1.1	0.9	0.0
Agency 41	1,488	77.3	4.9	13.6	0.3	2.1	1.7	0.0
Agency 42	598	79.3	2.8	16.6	0.0	0.5	0.8	0.0
Agency 43	2,740	61.5	21.2	9.0	0.5	4.8	3.1	0.0
Agency 45	1,883	70.3	8.9	18.0	0.1	2.0	0.8	0.0
Agency 46	1,520	75.9	6.6	13.0	0.1	3.4	1.1	0.1
Agency 47	1,042	55.0	3.1	33.3	0.2	7.8	0.7	0.0
Agency 48	982	70.7	1.5	24.9	0.4	2.2	0.2	0.0
Agency 49	1,940	43.1	10.8	36.5	3.5	3.5	2.5	0.0
Agency 50	1,096	66.8	5.3	25.2	0.3	1.1	1.4	0.0
Agency 51	1,383	41.9	36.5	17.0	0.2	2.1	2.2	0.0
Agency 52	1,479	78.2	3.2	15.8	0.7	0.5	1.6	0.0
Agency 53	664	91.1	2.7	3.9	0.3	0.9	1.1	0.0
Agency 54	1,954	58.7	23.2	10.5	0.5	1.9	5.1	0.0
Iowa	**34,950**	**60.9**	**13.4**	**18.6**	**0.5**	**3.9**	**2.6**	**0.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Local Agency

Combined 3 Years

Comparison of Age Distribution (2)								
Local Agency	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Agency 31	6,995	0.1	2.4	6.2	57.9	30.5	2.9	0.0
Agency 33	567	0.0	1.4	6.2	64.0	26.5	1.9	0.0
Agency 34	2,630	0.0	1.7	6.2	61.0	29.1	2.0	0.0
Agency 35	1,903	0.0	2.5	6.6	63.5	25.9	1.3	0.2
Agency 37	865	0.1	1.7	8.0	64.0	24.6	1.4	0.1
Agency 38	2,429	0.0	2.2	6.1	60.2	29.4	2.0	0.0
Agency 39	792	0.1	2.8	9.1	58.8	26.6	2.4	0.1
Agency 41	1,488	0.1	1.1	6.2	63.7	27.5	1.4	0.0
Agency 42	598	0.0	2.0	4.7	61.0	31.4	0.8	0.0
Agency 43	2,740	0.1	1.8	4.9	62.8	28.9	1.5	0.0
Agency 45	1,883	0.1	3.2	7.5	61.9	25.7	1.6	0.0
Agency 46	1,520	0.0	2.2	6.7	64.1	25.2	1.8	0.1
Agency 47	1,042	0.0	2.1	5.6	63.1	27.1	2.1	0.0
Agency 48	982	0.0	1.7	5.4	62.1	29.0	1.6	0.1
Agency 49	1,940	0.1	2.0	6.2	60.0	29.7	2.1	0.0
Agency 50	1,096	0.2	3.1	6.7	60.9	27.6	1.5	0.0
Agency 51	1,383	0.0	0.8	3.8	53.6	38.8	3.0	0.0
Agency 52	1,479	0.0	2.6	7.2	64.8	23.8	1.6	0.0
Agency 53	664	0.0	2.0	6.2	63.9	26.2	1.8	0.0
Agency 54	1,954	0.2	3.1	7.8	62.6	24.8	1.6	0.0
Iowa	**34,950**	**0.1**	**2.2**	**6.3**	**61.0**	**28.4**	**2.0**	**0.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Local Agency

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agency 31	5,520	4.1 (15)	56.8 (1)	5,289	24.5 (16)	41.5 (2)	1,567	2.6 (4)	511	5.1 (8)	393	100.0 (1)	181	100.0 (1)
Agency 33	464	3.4 (9)	64.2 (19)	444	18.7 (1)	49.8 (19)	152	31.6 (15)	132	21.2 (13)	45	100.0 (1)	34	100.0 (1)
Agency 34	1,935	3.2 (6)	61.0 (8)	1,857	21.8 (7)	49.1 (18)	544	36.4 (19)	361	34.9 (19)	212	100.0 (1)	78	100.0 (1)
Agency 35	1,478	4.0 (14)	61.8 (13)	1,405	21.5 (4)	50.2 (20)	400	43.5 (20)	162	35.2 (20)	99	100.0 (1)	43	100.0 (1)
Agency 37	612	4.7 (19)	60.1 (7)	585	25.0 (19)	44.1 (6)	155	35.5 (18)	116	28.4 (17)	46	100.0 (1)	18	100.0 (1)
Agency 38	1,915	4.1 (15)	59.9 (5)	1,842	24.9 (18)	42.5 (3)	442	17.2 (12)	399	19.5 (12)	188	100.0 (1)	28	100.0 (1)
Agency 39	601	3.3 (8)	61.2 (9)	582	27.0 (20)	42.6 (4)	175	0.6 (1)	224	1.8 (1)	29	100.0 (1)	8	100.0 (1)
Agency 41	1,123	3.2 (6)	66.7 (20)	1,083	19.3 (2)	48.3 (15)	238	5.0 (8)	208	7.2 (10)	123	100.0 (1)	44	100.0 (1)
Agency 42	454	3.1 (5)	61.5 (11)	435	22.1 (9)	47.1 (13)	147	29.3 (13)	158	30.4 (18)	51	100.0 (1)	24	100.0 (1)
Agency 43	2,170	3.5 (11)	61.2 (9)	2,057	21.6 (5)	47.1 (13)	472	5.7 (10)	256	4.7 (4)	130	100.0 (1)	89	100.0 (1)
Agency 45	1,401	3.9 (12)	59.9 (5)	1,344	20.4 (3)	48.3 (15)	439	3.0 (5)	196	4.6 (3)	152	100.0 (1)	76	100.0 (1)
Agency 46	1,278	4.1 (15)	61.7 (12)	1,203	22.6 (11)	45.3 (9)	388	29.6 (14)	287	25.1 (15)	11	100.0 (1)	36	100.0 (1)
Agency 47	772	2.7 (2)	62.2 (15)	727	23.5 (12)	45.0 (8)	245	0.8 (2)	146	4.8 (7)	75	100.0 (1)	34	100.0 (1)
Agency 48	802	3.0 (4)	62.6 (17)	781	24.7 (17)	46.1 (10)	277	1.8 (3)	221	3.2 (2)	4	100.0 (1)	26	100.0 (1)
Agency 49	1,552	2.6 (1)	62.6 (17)	1,503	21.6 (5)	43.7 (5)	527	4.0 (6)	594	4.7 (4)	130	100.0 (1)	52	100.0 (1)
Agency 50	903	2.7 (2)	62.5 (16)	864	23.7 (13)	44.1 (6)	249	4.8 (7)	94	10.6 (11)	60	100.0 (1)	26	100.0 (1)
Agency 51	980	3.9 (12)	57.4 (2)	915	24.2 (15)	40.4 (1)	328	35.1 (17)	378	25.4 (16)	61	100.0 (1)	37	100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Local Agency

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agency 52	1,131	3.4 (9)	62.0 (14)	1,087	22.3 (10)	48.9 (17)	357	7.0 (11)	171	4.7 (4)	50	100.0 (1)	51	100.0 (1)
Agency 53	462	5.4 (20)	59.7 (4)	451	23.7 (13)	46.1 (10)	134	5.2 (9)	102	5.9 (9)	28	100.0 (1)	13	100.0 (1)
Agency 54	1,418	4.1 (15)	59.2 (3)	1,337	21.9 (8)	46.2 (12)	403	34.0 (16)	234	22.2 (14)	15	100.0 (1)	75	100.0 (1)
Iowa	**26,971**	**3.7**	**60.4**	**25,791**	**22.9**	**45.2**	**7,639**	**14.7**	**4,950**	**14.6**	**1,902**	**100.0**	**973**	**100.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Local Agency

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31	4,463	29.5 (13)	12.3 (12)	6,152	1.1 (11)	1,891	14.1 (20)	2,549	87.4 (14)
Agency 33	405	30.9 (10)	7.7 (4)	550	1.1 (11)	182	20.3 (12)	192	84.4 (18)
Agency 34	1,775	26.1 (18)	10.2 (9)	2,240	0.9 (14)	1,262	16.2 (19)	1,628	90.3 (8)
Agency 35	1,404	35.8 (4)	11.9 (11)	1,525	2.0 (5)	1,141	17.8 (15)	1,311	89.7 (10)
Agency 37	529	37.6 (2)	8.5 (6)	673	1.6 (7)	271	27.7 (3)	265	90.2 (9)
Agency 38	1,600	33.9 (6)	10.3 (10)	2,033	1.6 (7)	1,334	17.5 (16)	1,414	87.3 (15)
Agency 39	507	28.8 (15)	8.5 (6)	710	1.5 (9)	156	23.7 (7)	164	92.7 (3)
Agency 41	1,082	37.7 (1)	17.7 (16)	1,234	1.0 (13)	1,026	21.8 (9)	1,104	86.8 (17)
Agency 42	430	28.6 (16)	12.3 (12)	498	0.4 (20)	441	25.6 (4)	446	91.0 (6)
Agency 43	1,622	35.4 (5)	19.7 (18)	2,340	2.1 (4)	1,054	22.5 (8)	1,235	88.7 (11)
Agency 45	1,392	31.4 (8)	6.5 (2)	1,533	1.2 (10)	1,319	18.7 (14)	1,407	92.8 (2)
Agency 46	986	31.1 (9)	23.5 (20)	1,342	2.5 (1)	210	21.4 (10)	215	79.5 (20)
Agency 47	738	24.9 (19)	8.3 (5)	864	0.8 (15)	635	50.2 (1)	762	94.0 (1)
Agency 48	645	30.7 (11)	5.9 (1)	935	0.6 (18)	139	34.5 (2)	199	87.9 (13)
Agency 49	1,319	29.8 (12)	8.5 (6)	1,762	0.8 (15)	339	25.1 (5)	434	91.5 (5)
Agency 50	847	28.9 (14)	22.1 (19)	965	2.4 (2)	771	17.0 (18)	843	80.2 (19)
Agency 51	798	18.0 (20)	18.5 (17)	1,209	1.7 (6)	292	20.9 (11)	322	88.5 (12)
Agency 52	953	36.1 (3)	7.3 (3)	1,347	0.7 (17)	469	19.4 (13)	478	87.0 (16)
Agency 53	410	27.8 (17)	16.3 (15)	525	0.6 (18)	361	24.4 (6)	376	90.7 (7)
Agency 54	1,188	32.1 (7)	12.5 (14)	1,609	2.3 (3)	109	17.4 (17)	112	92.0 (4)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Local Agency

Combined 3 Years

Iowa	**23,093**	**30.9**	**12.6**	**30,046**	**1.4**	**13,402**	**20.6**	**15,456**	**88.7**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Local Agency

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31	3,345	23.0 (3)	3,299	14.8 (6)	790	22.5 (14)	3,751	5.7 (6)	3,874	1.9 (11)
Agency 33	430	39.8 (20)	527	21.3 (17)	173	35.3 (3)	429	8.6 (12)	525	2.5 (15)
Agency 34	1,840	30.1 (10)	2,162	16.8 (8)	630	34.3 (4)	1,949	7.0 (9)	2,349	1.4 (6)
Agency 35	1,479	23.6 (5)	1,643	13.8 (4)	333	24.9 (11)	1,472	1.8 (1)	1,612	0.7 (2)
Agency 37	576	37.5 (19)	550	22.9 (18)	199	18.6 (19)	575	10.6 (16)	517	2.3 (13)
Agency 38	1,802	23.6 (5)	1,992	14.9 (7)	422	29.1 (7)	1,815	7.7 (11)	1,974	1.8 (8)
Agency 39	538	21.2 (2)	452	13.3 (3)	103	22.3 (15)	535	3.2 (2)	439	1.8 (8)
Agency 41	1,152	31.6 (13)	1,405	19.4 (15)	367	26.4 (8)	1,134	6.2 (8)	1,324	1.1 (4)
Agency 42	455	30.1 (10)	485	16.9 (9)	132	32.6 (6)	463	15.8 (20)	475	11.8 (20)
Agency 43	1,789	33.8 (14)	1,360	19.3 (14)	532	23.9 (12)	1,816	9.0 (13)	1,413	1.8 (8)
Agency 45	1,450	35.1 (16)	1,775	19.5 (16)	497	34.0 (5)	1,445	12.6 (19)	1,742	2.9 (16)
Agency 46	1,074	29.3 (8)	699	18.2 (12)	274	13.5 (20)	1,065	11.2 (17)	657	2.4 (14)
Agency 47	785	23.1 (4)	975	10.6 (2)	170	43.5 (1)	787	9.8 (15)	945	7.8 (19)
Agency 48	684	29.5 (9)	517	17.6 (11)	177	21.5 (16)	683	4.2 (3)	513	0.0 (1)
Agency 49	1,387	25.6 (7)	1,078	13.9 (5)	303	25.1 (10)	1,397	5.5 (5)	1,157	1.7 (7)
Agency 50	911	33.9 (15)	1,017	19.0 (13)	311	39.5 (2)	908	7.4 (10)	989	1.9 (11)
Agency 51	832	16.0 (1)	677	8.7 (1)	99	26.3 (9)	828	4.2 (3)	662	0.8 (3)
Agency 52	1,028	37.2 (18)	907	23.7 (19)	360	23.1 (13)	1,023	6.1 (7)	921	1.1 (4)
Agency 53	412	36.2 (17)	467	24.0 (20)	142	21.1 (17)	428	9.6 (14)	474	3.4 (17)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Local Agency

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 54	1,253	30.6 (12)	932	17.4 (10)	315	20.3 (18)	1,253	11.2 (17)	920	3.6 (18)
Iowa	**23,222**	**28.5**	**22,919**	**16.8**	**6,329**	**27.0**	**23,756**	**7.4**	**23,482**	**2.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Local Agency

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31	6,486	8.3 (15)	8.2 (9)	5,932	9.9 (11)	5,266	3.4 (15)	6,583	78.7 (3)
Agency 33	539	8.7 (18)	8.5 (14)	531	10.4 (14)	476	3.4 (15)	539	70.1 (11)
Agency 34	2,418	7.3 (9)	8.9 (16)	2,154	9.2 (2)	1,916	2.8 (9)	2,471	68.9 (14)
Agency 35	1,775	8.4 (17)	8.2 (9)	1,449	9.2 (2)	1,314	3.2 (12)	1,781	59.6 (20)
Agency 37	814	7.0 (7)	7.4 (5)	653	9.6 (6)	590	3.1 (10)	813	66.5 (16)
Agency 38	2,290	7.4 (11)	9.4 (18)	1,973	9.3 (4)	1,777	3.2 (12)	2,305	73.1 (7)
Agency 39	754	6.8 (5)	7.3 (4)	698	10.9 (17)	618	3.2 (12)	758	80.6 (2)
Agency 41	1,397	6.9 (6)	10.2 (20)	1,196	10.4 (14)	1,068	2.7 (6)	1,403	68.9 (14)
Agency 42	574	6.6 (2)	9.2 (17)	487	7.8 (1)	449	3.1 (10)	574	76.1 (5)
Agency 43	2,527	7.7 (14)	8.4 (13)	2,237	9.7 (8)	2,002	3.4 (15)	2,547	70.0 (12)
Agency 45	1,794	7.5 (12)	7.8 (7)	1,484	9.7 (8)	1,337	3.6 (18)	1,798	65.7 (18)
Agency 46	1,386	6.4 (1)	8.2 (9)	1,260	9.5 (5)	1,140	2.2 (4)	1,388	70.4 (10)
Agency 47	972	6.6 (2)	8.6 (15)	841	10.6 (16)	749	2.4 (5)	977	71.5 (8)
Agency 48	945	7.1 (8)	7.1 (1)	922	11.9 (20)	812	1.6 (1)	945	66.1 (17)
Agency 49	1,812	6.6 (2)	8.3 (12)	1,718	10.3 (12)	1,513	2.0 (2)	1,844	71.1 (9)
Agency 50	1,018	7.6 (13)	8.0 (8)	925	10.3 (12)	830	2.7 (6)	1,019	76.2 (4)
Agency 51	1,247	10.7 (20)	7.1 (1)	1,131	9.8 (10)	1,008	4.2 (19)	1,263	83.7 (1)
Agency 52	1,370	8.3 (15)	7.5 (6)	1,293	11.2 (19)	1,144	2.7 (6)	1,378	75.3 (6)
Agency 53	642	7.3 (9)	10.0 (19)	521	9.6 (6)	471	2.1 (3)	641	69.4 (13)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8F

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Local Agency

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Local Agency	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 54	1,828	10.1 (19)	7.1 (1)	1,547	11.1 (18)	1,373	4.7 (20)	1,832	64.8 (19)
Iowa	**32,588**	**7.8**	**8.3**	**28,952**	**10.0**	**25,853**	**3.1**	**32,859**	**67.7**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.