WIC PNSS & PedNSS

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Ackley	3	33.3	0.0	66.7	0.0	0.0	0.0	0.0
Adel	11	81.8	0.0	18.2	0.0	0.0	0.0	0.0
Albia	14	92.9	0.0	0.0	7.1	0.0	0.0	0.0
Algona	47	76.6	6.4	17.0	0.0	0.0	0.0	0.0
Allen WHC	102	51.0	30.4	10.8	0.0	4.9	2.9	0.0
Allison	23	95.7	0.0	0.0	0.0	0.0	4.3	0.0
Altoona	42	76.2	14.3	4.8	0.0	2.4	2.4	0.0
Ames	214	68.7	11.7	11.7	0.0	6.5	1.4	0.0
Anamosa	26	96.2	0.0	3.8	0.0	0.0	0.0	0.0
Ankeny	54	66.7	11.1	13.0	0.0	3.7	5.6	0.0
Atlantic	36	94.4	0.0	5.6	0.0	0.0	0.0	0.0
Audubon	9	77.8	0.0	22.2	0.0	0.0	0.0	0.0
Avoca	7	85.7	14.3	0.0	0.0	0.0	0.0	0.0
Bedford	9	88.9	0.0	0.0	0.0	0.0	11.1	0.0
Belle Plaine	8	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Bloomfield	9	88.9	11.1	0.0	0.0	0.0	0.0	0.0
Boone	27	88.9	3.7	3.7	0.0	0.0	3.7	0.0
Brdlwns RP	1,171	38.9	26.1	18.5	0.3	11.3	4.8	0.0
Brooklyn	3	33.3	0.0	66.7	0.0	0.0	0.0	0.0
Burlington	171	76.6	19.3	3.5	0.0	0.0	0.6	0.0
CHC	268	57.5	21.6	14.2	0.4	2.2	4.1	0.0
CR Urban	483	47.8	40.2	4.6	0.2	1.0	6.2	0.0
Carlisle	13	100.0	0.0	0.0	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Carroll	70	70.0	5.7	24.3	0.0	0.0	0.0	0.0
Cedar Falls	11	63.6	0.0	18.2	0.0	18.2	0.0	0.0
Centervill	30	93.3	3.3	3.3	0.0	0.0	0.0	0.0
Chariton	25	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Charles City	65	76.9	7.7	9.2	0.0	6.2	0.0	0.0
Cherokee	39	89.7	0.0	10.3	0.0	0.0	0.0	0.0
Clarinda	27	88.9	0.0	3.7	0.0	0.0	7.4	0.0
Clarion	50	40.0	0.0	60.0	0.0	0.0	0.0	0.0
Clinton	182	81.3	8.2	8.2	0.0	0.0	2.2	0.0
Colfax	11	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Columbus J	17	17.6	0.0	41.2	5.9	35.3	0.0	0.0
Corning	15	93.3	0.0	6.7	0.0	0.0	0.0	0.0
Corydon	17	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Council Blfs	420	77.4	3.6	17.1	0.2	0.2	1.4	0.0
Cresco	28	89.3	0.0	10.7	0.0	0.0	0.0	0.0
Creston	78	87.2	0.0	11.5	0.0	0.0	1.3	0.0
Dallas Ctr	3	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Decorah	22	81.8	4.5	13.6	0.0	0.0	0.0	0.0
Denison	104	26.9	1.9	62.5	0.0	8.7	0.0	0.0
Dewitt	6	83.3	0.0	0.0	0.0	0.0	16.7	0.0
Dubuque	377	68.7	16.7	7.2	0.3	5.3	1.9	0.0
Dyersville	7	85.7	14.3	0.0	0.0	0.0	0.0	0.0
Eagle Grove	29	34.5	0.0	65.5	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Edgerton	382	53.9	26.7	9.2	1.0	3.1	6.0	0.0
Eldora	7	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Elkader	29	96.6	0.0	3.4	0.0	0.0	0.0	0.0
Emmetsburg	13	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Esthervill	39	59.0	0.0	41.0	0.0	0.0	0.0	0.0
Fairfiel	17	94.1	0.0	0.0	0.0	5.9	0.0	0.0
Forest City	60	83.3	8.3	8.3	0.0	0.0	0.0	0.0
Fort Dodge	227	73.6	11.0	12.3	0.9	0.4	1.8	0.0
Ft Madison	56	89.3	1.8	8.9	0.0	0.0	0.0	0.0
Garner	27	81.5	0.0	14.8	0.0	0.0	3.7	0.0
Glenwood	20	90.0	0.0	10.0	0.0	0.0	0.0	0.0
Grand Junction	3	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Greenfield	20	95.0	0.0	5.0	0.0	0.0	0.0	0.0
Grinnell	29	86.2	3.4	10.3	0.0	0.0	0.0	0.0
Grundy Ctr	9	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Guthrie Ct	15	73.3	6.7	13.3	0.0	0.0	6.7	0.0
Hampton	45	53.3	2.2	44.4	0.0	0.0	0.0	0.0
Harlan	39	82.1	2.6	15.4	0.0	0.0	0.0	0.0
Hawarden	17	47.1	0.0	52.9	0.0	0.0	0.0	0.0
Hispanic Cln	6	16.7	0.0	83.3	0.0	0.0	0.0	0.0
Humboldt	19	73.7	0.0	26.3	0.0	0.0	0.0	0.0
ILH	391	38.4	16.1	35.3	0.0	5.9	4.3	0.0
Ida Grove	26	76.9	0.0	19.2	0.0	0.0	3.8	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Independen	32	96.9	3.1	0.0	0.0	0.0	0.0	0.0
Indianola	51	88.2	5.9	3.9	0.0	0.0	2.0	0.0
Iowa City	368	34.5	43.2	17.1	0.5	1.9	2.7	0.0
Iowa Falls	15	73.3	0.0	20.0	0.0	0.0	6.7	0.0
Jamaica	3	66.7	0.0	33.3	0.0	0.0	0.0	0.0
Jefferson	17	82.4	0.0	17.6	0.0	0.0	0.0	0.0
KR	14	57.1	14.3	7.1	0.0	21.4	0.0	0.0
Keokuk	55	92.7	3.6	3.6	0.0	0.0	0.0	0.0
Keosauqua	7	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Knoxville	94	92.6	3.2	3.2	0.0	0.0	1.1	0.0
Lamoni	6	50.0	16.7	16.7	0.0	0.0	16.7	0.0
Le Mars	46	67.4	6.5	21.7	2.2	2.2	0.0	0.0
Lenox	13	69.2	0.0	30.8	0.0	0.0	0.0	0.0
Leon	22	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Logan	34	91.2	0.0	5.9	2.9	0.0	0.0	0.0
Manchester	16	93.8	0.0	6.3	0.0	0.0	0.0	0.0
Manning	2	50.0	0.0	0.0	0.0	50.0	0.0	0.0
Maquoketa	30	83.3	10.0	0.0	0.0	0.0	6.7	0.0
Marengo	11	90.9	0.0	0.0	0.0	0.0	9.1	0.0
Marion	130	83.1	2.3	10.8	0.0	0.0	3.8	0.0
Marshalltown	303	41.9	3.0	45.5	0.7	7.9	1.0	0.0
Mason City	216	75.5	6.0	10.6	0.5	3.2	4.2	0.0
Monticello	10	100.0	0.0	0.0	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Mt Ayr	12	83.3	0.0	16.7	0.0	0.0	0.0	0.0
Mt Pleasan	55	80.0	1.8	16.4	0.0	1.8	0.0	0.0
Mtown Dtown	2	0.0	0.0	50.0	0.0	50.0	0.0	0.0
Muscatine	231	56.7	6.9	32.9	0.0	2.6	0.9	0.0
Nevada	4	25.0	0.0	75.0	0.0	0.0	0.0	0.0
New Hampto	17	70.6	0.0	29.4	0.0	0.0	0.0	0.0
Newton	93	77.4	7.5	9.7	2.2	3.2	0.0	0.0
North Towne	150	63.3	22.7	8.7	0.0	2.7	2.7	0.0
Northwood	10	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Norwalk	11	81.8	0.0	9.1	0.0	9.1	0.0	0.0
Oakridge	8	25.0	50.0	12.5	0.0	0.0	12.5	0.0
Oelwein	34	85.3	2.9	11.8	0.0	0.0	0.0	0.0
Onawa	28	96.4	0.0	0.0	3.6	0.0	0.0	0.0
Orange Cty	37	43.2	2.7	51.4	0.0	2.7	0.0	0.0
Osage	31	83.9	0.0	12.9	0.0	0.0	3.2	0.0
Osceola	54	70.4	1.9	25.9	0.0	1.9	0.0	0.0
Oskaloosa	44	88.6	6.8	4.5	0.0	0.0	0.0	0.0
Ottumwa	415	70.4	8.4	14.9	0.0	4.8	1.2	0.2
Pella	14	85.7	0.0	0.0	0.0	14.3	0.0	0.0
Perry	40	37.5	10.0	52.5	0.0	0.0	0.0	0.0
Pocahontas	3	66.7	33.3	0.0	0.0	0.0	0.0	0.0
Postville	37	35.1	5.4	59.5	0.0	0.0	0.0	0.0
Red Oak	29	96.6	0.0	3.4	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Redfield	9	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Refugee Cln	13	7.7	92.3	0.0	0.0	0.0	0.0	0.0
Rock Rapid	14	92.9	0.0	0.0	0.0	7.1	0.0	0.0
Rock Valle	37	40.5	0.0	56.8	2.7	0.0	0.0	0.0
Rockwell C	20	90.0	0.0	10.0	0.0	0.0	0.0	0.0
SDM	403	43.4	15.4	28.0	0.5	8.7	4.0	0.0
Sac City	12	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Sheldon	37	67.6	8.1	21.6	2.7	0.0	0.0	0.0
Shenandoah	31	93.5	0.0	6.5	0.0	0.0	0.0	0.0
Sibley	16	56.3	6.3	37.5	0.0	0.0	0.0	0.0
Sidney	7	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Sigourney	10	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Sioux Cent	42	21.4	0.0	73.8	0.0	0.0	4.8	0.0
Sioux City	683	41.9	11.6	36.7	3.2	3.8	2.8	0.0
Spencer	59	79.7	1.7	16.9	0.0	1.7	0.0	0.0
Spirit Lak	21	85.7	4.8	4.8	0.0	0.0	4.8	0.0
Storm Lake	117	22.2	4.3	53.8	0.0	18.8	0.9	0.0
Stuart	11	81.8	9.1	0.0	0.0	0.0	9.1	0.0
Tama	40	40.0	0.0	52.5	5.0	0.0	2.5	0.0
Tama/Settl	2	50.0	0.0	0.0	50.0	0.0	0.0	0.0
Tipton	9	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Traer	2	100.0	0.0	0.0	0.0	0.0	0.0	0.0
UIHC	13	69.2	30.8	0.0	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Vinton	27	92.6	0.0	7.4	0.0	0.0	0.0	0.0
WDM	61	47.5	18.0	26.2	0.0	6.6	1.6	0.0
Washington	36	75.0	0.0	22.2	0.0	0.0	2.8	0.0
Waterloo	661	53.9	26.9	10.0	0.3	5.6	3.3	0.0
Waukee	37	70.3	2.7	16.2	0.0	2.7	8.1	0.0
Waukon	23	91.3	0.0	4.3	0.0	4.3	0.0	0.0
Waverly	17	88.2	11.8	0.0	0.0	0.0	0.0	0.0
Wbstr Ct	68	57.4	2.9	29.4	0.0	10.3	0.0	0.0
West Union	19	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Winterset	24	87.5	4.2	8.3	0.0	0.0	0.0	0.0
Iowa	**11,313**	**59.6**	**14.6**	**18.7**	**0.5**	**4.1**	**2.6**	**0.0**
National (Prior Year)	**133,781**	**53.3**	**21.7**	**17.1**	**1.1**	**2.8**	**3.5**	**0.4**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Ackley	3	0.0	0.0	0.0	66.7	33.3	0.0	0.0
Adel	11	0.0	0.0	0.0	81.8	18.2	0.0	0.0
Albia	14	0.0	0.0	7.1	71.4	21.4	0.0	0.0
Algona	47	0.0	2.1	2.1	53.2	42.6	0.0	0.0
Allen WHC	102	1.0	2.9	7.8	51.0	35.3	2.0	0.0
Allison	23	0.0	0.0	0.0	69.6	30.4	0.0	0.0
Altoona	42	0.0	4.8	7.1	42.9	42.9	2.4	0.0
Ames	214	0.0	1.9	6.5	59.8	30.8	0.9	0.0
Anamosa	26	0.0	3.8	0.0	80.8	15.4	0.0	0.0
Ankeny	54	0.0	0.0	3.7	64.8	29.6	1.9	0.0
Atlantic	36	0.0	2.8	0.0	72.2	22.2	2.8	0.0
Audubon	9	0.0	11.1	11.1	44.4	33.3	0.0	0.0
Avoca	7	0.0	0.0	0.0	71.4	28.6	0.0	0.0
Bedford	9	0.0	0.0	0.0	77.8	22.2	0.0	0.0
Belle Plaine	8	0.0	0.0	0.0	75.0	25.0	0.0	0.0
Bloomfield	9	0.0	0.0	0.0	77.8	11.1	11.1	0.0
Boone	27	0.0	0.0	7.4	70.4	22.2	0.0	0.0
Brdlwns RP	1,171	0.1	2.3	5.4	58.6	31.2	2.5	0.0
Brooklyn	3	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Burlington	171	0.0	2.3	5.8	66.1	25.7	0.0	0.0
CHC	268	0.0	1.9	8.6	61.9	25.0	2.6	0.0
CR Urban	483	0.0	1.9	6.6	58.0	31.3	2.3	0.0
Carlisle	13	0.0	0.0	7.7	61.5	23.1	7.7	0.0
Carroll	70	0.0	1.4	7.1	60.0	28.6	2.9	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cedar Falls	11	0.0	0.0	0.0	63.6	36.4	0.0	0.0
Centervill	30	0.0	0.0	3.3	63.3	33.3	0.0	0.0
Chariton	25	0.0	0.0	4.0	60.0	32.0	4.0	0.0
Charles City	65	0.0	1.5	6.2	67.7	24.6	0.0	0.0
Cherokee	39	0.0	2.6	17.9	59.0	20.5	0.0	0.0
Clarinda	27	0.0	0.0	11.1	77.8	11.1	0.0	0.0
Clarion	50	0.0	6.0	4.0	54.0	36.0	0.0	0.0
Clinton	182	0.0	3.8	8.8	62.6	22.5	2.2	0.0
Colfax	11	0.0	9.1	18.2	54.5	18.2	0.0	0.0
Columbus J	17	0.0	0.0	0.0	41.2	47.1	11.8	0.0
Corning	15	0.0	0.0	13.3	53.3	33.3	0.0	0.0
Corydon	17	0.0	5.9	0.0	47.1	47.1	0.0	0.0
Council Blfs	420	0.0	1.4	8.1	66.9	21.7	1.9	0.0
Cresco	28	0.0	0.0	0.0	57.1	39.3	3.6	0.0
Creston	78	1.3	1.3	11.5	65.4	19.2	1.3	0.0
Dallas Ctr	3	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Decorah	22	0.0	4.5	9.1	50.0	36.4	0.0	0.0
Denison	104	0.0	0.0	2.9	59.6	32.7	4.8	0.0
Dewitt	6	0.0	0.0	16.7	66.7	16.7	0.0	0.0
Dubuque	377	0.0	2.7	6.6	59.2	29.7	1.6	0.3
Dyersville	7	0.0	0.0	0.0	71.4	28.6	0.0	0.0
Eagle Grove	29	0.0	0.0	10.3	51.7	31.0	6.9	0.0
Edgerton	382	0.3	4.2	6.5	60.5	25.7	2.9	0.0
Eldora	7	0.0	0.0	0.0	57.1	42.9	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Elkader	29	0.0	0.0	0.0	75.9	24.1	0.0	0.0
Emmetsburg	13	0.0	0.0	7.7	69.2	23.1	0.0	0.0
Esthervill	39	0.0	2.6	5.1	64.1	25.6	2.6	0.0
Fairfiel	17	0.0	0.0	0.0	70.6	29.4	0.0	0.0
Forest City	60	0.0	0.0	5.0	66.7	28.3	0.0	0.0
Fort Dodge	227	0.0	0.9	6.6	66.1	25.1	1.3	0.0
Ft Madison	56	0.0	1.8	3.6	64.3	28.6	1.8	0.0
Garner	27	0.0	0.0	14.8	59.3	25.9	0.0	0.0
Glenwood	20	0.0	0.0	5.0	80.0	15.0	0.0	0.0
Grand Junction	3	0.0	0.0	100.0	0.0	0.0	0.0	0.0
Greenfield	20	0.0	0.0	0.0	70.0	30.0	0.0	0.0
Grinnell	29	0.0	6.9	3.4	58.6	31.0	0.0	0.0
Grundy Ctr	9	0.0	0.0	0.0	88.9	11.1	0.0	0.0
Guthrie Ct	15	0.0	6.7	6.7	66.7	20.0	0.0	0.0
Hampton	45	0.0	4.4	4.4	53.3	35.6	2.2	0.0
Harlan	39	0.0	5.1	2.6	61.5	28.2	2.6	0.0
Hawarden	17	0.0	0.0	5.9	52.9	41.2	0.0	0.0
Hispanic Cln	6	0.0	0.0	16.7	50.0	33.3	0.0	0.0
Humboldt	19	0.0	0.0	5.3	63.2	31.6	0.0	0.0
ILH	391	0.3	2.3	6.1	52.7	36.1	2.6	0.0
Ida Grove	26	0.0	0.0	11.5	65.4	19.2	3.8	0.0
Independen	32	0.0	0.0	0.0	65.6	28.1	6.3	0.0
Indianola	51	0.0	5.9	9.8	62.7	19.6	2.0	0.0
Iowa City	368	0.0	0.5	3.8	49.2	42.7	3.8	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Iowa Falls	15	0.0	6.7	6.7	60.0	26.7	0.0	0.0
Jamaica	3	0.0	0.0	0.0	33.3	66.7	0.0	0.0
Jefferson	17	0.0	0.0	5.9	82.4	11.8	0.0	0.0
KR	14	0.0	0.0	0.0	50.0	42.9	7.1	0.0
Keokuk	55	0.0	0.0	5.5	70.9	23.6	0.0	0.0
Keosauqua	7	0.0	14.3	0.0	28.6	57.1	0.0	0.0
Knoxville	94	0.0	0.0	11.7	62.8	23.4	2.1	0.0
Lamoni	6	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Le Mars	46	0.0	2.2	2.2	78.3	13.0	4.3	0.0
Lenox	13	0.0	0.0	0.0	69.2	30.8	0.0	0.0
Leon	22	0.0	0.0	18.2	59.1	18.2	4.5	0.0
Logan	34	0.0	2.9	11.8	52.9	32.4	0.0	0.0
Manchester	16	0.0	0.0	6.3	62.5	25.0	6.3	0.0
Manning	2	0.0	0.0	50.0	50.0	0.0	0.0	0.0
Maquoketa	30	0.0	0.0	3.3	70.0	26.7	0.0	0.0
Marengo	11	0.0	9.1	9.1	54.5	27.3	0.0	0.0
Marion	130	0.0	1.5	4.6	57.7	34.6	1.5	0.0
Marshalltown	303	0.0	3.6	6.3	58.4	27.7	4.0	0.0
Mason City	216	0.0	1.4	6.0	67.1	23.6	1.9	0.0
Monticello	10	0.0	0.0	10.0	40.0	50.0	0.0	0.0
Mt Ayr	12	0.0	0.0	8.3	75.0	16.7	0.0	0.0
Mt Pleasan	55	0.0	3.6	5.5	69.1	18.2	3.6	0.0
Mtown Dtown	2	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Muscatine	231	0.0	2.2	6.9	59.7	29.0	2.2	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Nevada	4	0.0	0.0	0.0	75.0	25.0	0.0	0.0
New Hampto	17	0.0	0.0	11.8	64.7	23.5	0.0	0.0
Newton	93	0.0	1.1	8.6	55.9	32.3	1.1	1.1
North Towne	150	0.0	3.3	8.7	56.7	29.3	2.0	0.0
Northwood	10	0.0	0.0	10.0	70.0	20.0	0.0	0.0
Norwalk	11	0.0	0.0	0.0	54.5	45.5	0.0	0.0
Oakridge	8	0.0	0.0	0.0	37.5	62.5	0.0	0.0
Oelwein	34	0.0	0.0	0.0	73.5	26.5	0.0	0.0
Onawa	28	0.0	0.0	10.7	57.1	32.1	0.0	0.0
Orange Cty	37	0.0	0.0	10.8	67.6	16.2	5.4	0.0
Osage	31	0.0	0.0	0.0	67.7	32.3	0.0	0.0
Osceola	54	0.0	1.9	1.9	64.8	29.6	1.9	0.0
Oskaloosa	44	0.0	0.0	0.0	61.4	34.1	4.5	0.0
Ottumwa	415	0.0	2.4	6.0	63.6	25.8	2.2	0.0
Pella	14	0.0	7.1	0.0	71.4	21.4	0.0	0.0
Perry	40	0.0	0.0	5.0	60.0	35.0	0.0	0.0
Pocahontas	3	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Postville	37	0.0	2.7	8.1	29.7	59.5	0.0	0.0
Red Oak	29	0.0	3.4	13.8	55.2	27.6	0.0	0.0
Redfield	9	0.0	0.0	0.0	77.8	22.2	0.0	0.0
Refugee Cln	13	0.0	7.7	0.0	30.8	53.8	7.7	0.0
Rock Rapid	14	0.0	0.0	0.0	42.9	50.0	7.1	0.0
Rock Valle	37	0.0	2.7	13.5	59.5	24.3	0.0	0.0
Rockwell C	20	0.0	0.0	0.0	80.0	20.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
SDM	403	0.0	1.7	6.0	60.5	30.0	1.7	0.0
Sac City	12	0.0	0.0	8.3	50.0	41.7	0.0	0.0
Sheldon	37	0.0	2.7	5.4	51.4	40.5	0.0	0.0
Shenandoah	31	0.0	9.7	6.5	61.3	19.4	3.2	0.0
Sibley	16	0.0	0.0	6.3	62.5	31.3	0.0	0.0
Sidney	7	0.0	0.0	0.0	57.1	42.9	0.0	0.0
Sigourney	10	0.0	0.0	10.0	50.0	40.0	0.0	0.0
Sioux Cent	42	0.0	4.8	9.5	54.8	31.0	0.0	0.0
Sioux City	683	0.0	1.9	6.3	59.4	30.3	2.0	0.0
Spencer	59	0.0	0.0	3.4	74.6	22.0	0.0	0.0
Spirit Lak	21	0.0	0.0	4.8	66.7	23.8	4.8	0.0
Storm Lake	117	0.0	1.7	3.4	55.6	35.0	4.3	0.0
Stuart	11	0.0	0.0	9.1	45.5	45.5	0.0	0.0
Tama	40	0.0	2.5	5.0	70.0	22.5	0.0	0.0
Tama/Settl	2	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Tipton	9	0.0	0.0	22.2	33.3	33.3	11.1	0.0
Traer	2	0.0	0.0	0.0	50.0	50.0	0.0	0.0
UIHC	13	0.0	7.7	0.0	53.8	38.5	0.0	0.0
Vinton	27	0.0	0.0	0.0	63.0	33.3	3.7	0.0
WDM	61	0.0	0.0	4.9	55.7	32.8	6.6	0.0
Washington	36	0.0	0.0	5.6	58.3	36.1	0.0	0.0
Waterloo	661	0.0	2.1	5.7	60.4	29.5	2.1	0.2
Waukee	37	0.0	0.0	5.4	59.5	35.1	0.0	0.0
Waukon	23	0.0	0.0	4.3	78.3	17.4	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Waverly	17	0.0	0.0	11.8	52.9	35.3	0.0	0.0
Wbstr Ct	68	0.0	4.4	4.4	64.7	25.0	1.5	0.0
West Union	19	0.0	5.3	10.5	78.9	5.3	0.0	0.0
Winterset	24	0.0	0.0	4.2	58.3	37.5	0.0	0.0
Iowa	**11,314**	**0.0**	**2.0**	**6.1**	**60.4**	**29.3**	**2.1**	**0.0**
National (Prior Year)	**133,790**	**0.1**	**2.7**	**8.0**	**62.0**	**25.2**	**1.9**	**0.1**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ackley	3	0.0 (1)	33.3 (8)	3	33.3 (120)	33.3 (28)	1	0.0 (1)	2	50.0 (134)	0	0.0 (1)	0	0.0 (1)
Adel	5	0.0 (1)	80.0 (135)	5	20.0 (52)	40.0 (52)	2	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Albia	12	0.0 (1)	16.7 (5)	12	33.3 (120)	41.7 (61)	1	0.0 (1)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Algona	34	2.9 (78)	73.5 (123)	34	23.5 (82)	47.1 (89)	3	0.0 (1)	11	9.1 (95)	4	100.0 (45)	1	100.0 (66)
Allen WHC	95	3.2 (81)	47.4 (14)	92	20.7 (60)	48.9 (98)	12	0.0 (1)	12	0.0 (1)	4	100.0 (45)	2	100.0 (66)
Allison	19	5.3 (106)	84.2 (142)	19	26.3 (101)	36.8 (40)	3	0.0 (1)	6	0.0 (1)	2	100.0 (45)	1	100.0 (66)
Altoona	30	6.7 (118)	56.7 (38)	29	41.4 (138)	24.1 (10)	7	0.0 (1)	5	0.0 (1)	5	100.0 (45)	2	100.0 (66)
Ames	152	5.3 (106)	59.2 (58)	144	32.6 (119)	41.7 (61)	35	2.9 (85)	33	15.2 (103)	13	100.0 (45)	1	100.0 (66)
Anamosa	22	0.0 (1)	63.6 (83)	21	14.3 (22)	52.4 (114)	5	20.0 (103)	3	33.3 (115)	1	100.0 (45)	1	100.0 (66)
Ankeny	36	8.3 (129)	66.7 (95)	34	17.6 (43)	41.2 (59)	17	0.0 (1)	10	0.0 (1)	3	100.0 (45)	0	0.0 (1)
Atlantic	31	3.2 (81)	77.4 (131)	30	20.0 (52)	70.0 (146)	8	12.5 (98)	10	0.0 (1)	0	0.0 (1)	3	100.0 (66)
Audubon	9	0.0 (1)	44.4 (12)	9	22.2 (67)	55.6 (121)	2	50.0 (134)	7	28.6 (112)	0	0.0 (1)	0	0.0 (1)
Avoca	5	0.0 (1)	60.0 (63)	3	33.3 (120)	66.7 (137)	2	0.0 (1)	0	0.0 (1)	0	0.0 (1)	1	100.0 (66)
Bedford	7	14.3 (145)	71.4 (115)	7	28.6 (108)	28.6 (21)	1	100.0 (148)	2	50.0 (134)	1	100.0 (45)	0	0.0 (1)
Belle Plaine	8	0.0 (1)	75.0 (126)	8	37.5 (133)	37.5 (43)	2	0.0 (1)	2	0.0 (1)	1	100.0 (45)	2	100.0 (66)
Bloomfield	9	11.1 (139)	55.6 (33)	8	25.0 (88)	50.0 (99)	3	33.3 (115)	4	0.0 (1)	0	0.0 (1)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Boone	22	0.0 (1)	72.7 (120)	22	31.8 (115)	50.0 (99)	5	20.0 (103)	5	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Brdlwns RP	914	4.2 (97)	53.9 (28)	883	25.4 (97)	39.9 (51)	267	3.0 (86)	85	7.1 (90)	66	100.0 (45)	35	100.0 (66)
Brooklyn	3	0.0 (1)	66.7 (95)	3	0.0 (1)	66.7 (137)	1	0.0 (1)	1	100.0 (145)	0	0.0 (1)	0	0.0 (1)
Burlington	129	6.2 (114)	54.3 (29)	125	20.0 (52)	51.2 (111)	46	8.7 (96)	18	5.6 (87)	14	100.0 (45)	15	100.0 (66)
CHC	198	4.5 (101)	64.6 (89)	185	23.8 (84)	45.4 (78)	64	43.8 (131)	32	34.4 (124)	2	100.0 (45)	11	100.0 (66)
CR Urban	352	3.7 (87)	59.4 (60)	343	25.1 (96)	46.6 (86)	113	49.6 (133)	47	31.9 (114)	43	100.0 (45)	17	100.0 (66)
Carlisle	6	0.0 (1)	50.0 (15)	6	16.7 (34)	33.3 (28)	2	0.0 (1)	0	0.0 (1)	0	0.0 (1)	1	100.0 (66)
Carroll	61	3.3 (84)	65.6 (92)	59	22.0 (65)	54.2 (119)	22	36.4 (122)	12	33.3 (115)	9	100.0 (45)	2	100.0 (66)
Cedar Falls	9	11.1 (139)	55.6 (33)	9	22.2 (67)	66.7 (137)	2	0.0 (1)	1	0.0 (1)	0	0.0 (1)	1	100.0 (66)
Centervill	24	4.2 (97)	79.2 (133)	23	26.1 (100)	52.2 (112)	9	33.3 (115)	4	50.0 (134)	1	100.0 (45)	3	100.0 (66)
Chariton	19	0.0 (1)	52.6 (23)	19	36.8 (132)	36.8 (40)	8	37.5 (126)	3	0.0 (1)	2	100.0 (45)	1	100.0 (66)
Charles City	44	2.3 (74)	68.2 (108)	43	16.3 (33)	41.9 (66)	13	15.4 (101)	11	0.0 (1)	7	100.0 (45)	3	100.0 (66)
Cherokee	26	11.5 (142)	57.7 (46)	26	15.4 (30)	42.3 (67)	4	0.0 (1)	18	0.0 (1)	4	100.0 (45)	1	100.0 (66)
Clarinda	20	0.0 (1)	60.0 (63)	19	21.1 (62)	52.6 (116)	4	50.0 (134)	6	33.3 (115)	0	0.0 (1)	0	0.0 (1)
Clarion	40	2.5 (76)	60.0 (63)	39	30.8 (112)	28.2 (19)	11	0.0 (1)	5	0.0 (1)	3	100.0 (45)	0	0.0 (1)
Clinton	126	2.4 (75)	61.9 (76)	121	19.0 (47)	57.0 (125)	44	36.4 (122)	13	15.4 (104)	7	100.0 (45)	11	100.0 (66)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Colfax	6	16.7 (146)	83.3 (139)	6	50.0 (141)	0.0 (1)	3	0.0 (1)	2	50.0 (134)	1	100.0 (45)	0	0.0 (1)
Columbus J	13	7.7 (126)	23.1 (7)	13	38.5 (134)	38.5 (49)	4	0.0 (1)	3	0.0 (1)	6	100.0 (45)	0	0.0 (1)
Corning	12	8.3 (129)	66.7 (95)	12	16.7 (34)	58.3 (130)	5	40.0 (128)	3	0.0 (1)	2	100.0 (45)	1	100.0 (66)
Corydon	15	6.7 (118)	66.7 (95)	15	33.3 (120)	46.7 (87)	4	0.0 (1)	3	0.0 (1)	0	0.0 (1)	1	100.0 (66)
Council Blfs	308	1.9 (70)	65.6 (92)	297	22.9 (78)	50.5 (109)	110	8.2 (94)	39	7.7 (93)	17	100.0 (45)	16	100.0 (66)
Cresco	21	9.5 (136)	66.7 (95)	20	20.0 (52)	65.0 (136)	12	8.3 (95)	11	36.4 (125)	3	100.0 (45)	2	100.0 (66)
Creston	60	10.0 (137)	56.7 (38)	59	22.0 (65)	47.5 (92)	14	35.7 (121)	12	25.0 (110)	11	100.0 (45)	2	100.0 (66)
Dallas Ctr	1	0.0 (1)	0.0 (1)	1	0.0 (1)	100.0 (147)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Decorah	16	12.5 (144)	43.8 (11)	16	6.3 (15)	50.0 (99)	5	0.0 (1)	4	0.0 (1)	3	100.0 (45)	1	100.0 (66)
Denison	93	1.1 (66)	60.2 (71)	92	39.1 (135)	28.3 (20)	40	2.5 (83)	22	0.0 (1)	0	0.0 (1)	6	100.0 (66)
Dewitt	5	0.0 (1)	80.0 (135)	4	50.0 (141)	25.0 (11)	1	0.0 (1)	1	0.0 (1)	1	100.0 (45)	0	0.0 (1)
Dubuque	312	3.5 (85)	62.2 (78)	301	23.3 (81)	44.9 (77)	105	46.7 (132)	33	48.5 (133)	22	100.0 (45)	7	100.0 (66)
Dyersville	6	0.0 (1)	83.3 (139)	6	16.7 (34)	16.7 (8)	2	50.0 (134)	2	0.0 (1)	1	100.0 (45)	0	0.0 (1)
Eagle Grove	27	3.7 (87)	59.3 (59)	27	25.9 (99)	37.0 (42)	11	0.0 (1)	1	0.0 (1)	3	100.0 (45)	1	100.0 (66)
Edgerton	304	3.6 (86)	55.3 (32)	291	21.3 (64)	46.0 (84)	91	27.5 (111)	67	22.4 (108)	6	100.0 (45)	13	100.0 (66)
Eldora	3	0.0 (1)	66.7 (95)	3	0.0 (1)	0.0 (1)	2	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Elkader	24	0.0 (1)	79.2 (133)	22	31.8 (115)	45.5 (79)	6	33.3 (115)	9	33.3 (115)	0	0.0 (1)	1	100.0 (66)
Emmetsburg	5	0.0 (1)	60.0 (63)	5	20.0 (52)	60.0 (132)	2	0.0 (1)	2	0.0 (1)	0	0.0 (1)	1	100.0 (66)
Esthervill	26	3.8 (91)	73.1 (122)	25	36.0 (130)	40.0 (52)	7	0.0 (1)	8	0.0 (1)	9	100.0 (45)	2	100.0 (66)
Fairfiel	12	0.0 (1)	58.3 (50)	12	25.0 (88)	25.0 (11)	4	0.0 (1)	3	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Forest City	50	4.0 (95)	76.0 (129)	46	17.4 (41)	47.8 (93)	15	0.0 (1)	12	8.3 (94)	3	100.0 (45)	6	100.0 (66)
Fort Dodge	190	3.7 (87)	67.4 (107)	184	21.2 (63)	50.5 (109)	65	1.5 (80)	9	0.0 (1)	7	100.0 (45)	5	100.0 (66)
Ft Madison	38	0.0 (1)	73.7 (124)	36	13.9 (21)	50.0 (99)	14	7.1 (92)	4	0.0 (1)	8	100.0 (45)	3	100.0 (66)
Garner	19	0.0 (1)	68.4 (110)	19	31.6 (114)	47.4 (91)	2	0.0 (1)	3	33.3 (115)	3	100.0 (45)	1	100.0 (66)
Glenwood	14	0.0 (1)	57.1 (41)	13	30.8 (112)	53.8 (117)	6	0.0 (1)	5	0.0 (1)	0	0.0 (1)	1	100.0 (66)
Grand Junction	3	0.0 (1)	0.0 (1)	2	0.0 (1)	50.0 (99)	1	0.0 (1)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Greenfield	16	0.0 (1)	68.8 (111)	16	12.5 (18)	37.5 (43)	1	0.0 (1)	3	0.0 (1)	1	100.0 (45)	0	0.0 (1)
Grinnell	24	8.3 (129)	66.7 (95)	21	14.3 (22)	61.9 (133)	6	33.3 (115)	5	40.0 (126)	3	100.0 (45)	2	100.0 (66)
Grundy Ctr	3	0.0 (1)	66.7 (95)	3	33.3 (120)	33.3 (28)	3	0.0 (1)	2	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Guthrie Ct	9	0.0 (1)	77.8 (132)	9	22.2 (67)	66.7 (137)	3	0.0 (1)	5	0.0 (1)	2	100.0 (45)	0	0.0 (1)
Hampton	40	0.0 (1)	85.0 (143)	39	17.9 (45)	53.8 (117)	7	0.0 (1)	8	0.0 (1)	6	100.0 (45)	2	100.0 (66)
Harlan	31	6.5 (116)	58.1 (48)	31	25.8 (98)	48.4 (96)	12	0.0 (1)	14	7.1 (90)	1	100.0 (45)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Hawarden	14	0.0 (1)	100.0 (144)	14	14.3 (22)	64.3 (135)	4	0.0 (1)	6	0.0 (1)	1	100.0 (45)	0	0.0 (1)
Hispanic Cln	6	0.0 (1)	100.0 (144)	6	33.3 (120)	33.3 (28)	1	0.0 (1)	1	100.0 (145)	0	0.0 (1)	0	0.0 (1)
Humboldt	17	0.0 (1)	64.7 (90)	17	35.3 (129)	35.3 (36)	3	0.0 (1)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)
ILH	324	4.3 (100)	59.6 (62)	314	26.8 (104)	40.4 (58)	79	2.5 (83)	30	3.3 (85)	21	100.0 (45)	8	100.0 (66)
Ida Grove	22	4.5 (101)	68.2 (108)	21	23.8 (84)	66.7 (137)	4	0.0 (1)	4	0.0 (1)	3	100.0 (45)	0	0.0 (1)
Independen	26	0.0 (1)	61.5 (74)	26	26.9 (105)	34.6 (34)	5	0.0 (1)	5	0.0 (1)	1	100.0 (45)	2	100.0 (66)
Indianola	34	5.9 (111)	58.8 (52)	34	32.4 (117)	44.1 (74)	6	0.0 (1)	7	0.0 (1)	1	100.0 (45)	0	0.0 (1)
Iowa City	269	4.1 (96)	59.1 (56)	252	22.2 (67)	39.7 (50)	99	36.4 (122)	114	22.8 (109)	15	100.0 (45)	15	100.0 (66)
Iowa Falls	12	8.3 (129)	66.7 (95)	11	54.5 (145)	27.3 (18)	2	50.0 (134)	5	20.0 (106)	1	100.0 (45)	1	100.0 (66)
Jamaica	3	0.0 (1)	100.0 (144)	3	0.0 (1)	66.7 (137)	2	50.0 (134)	0	0.0 (1)	1	100.0 (45)	1	100.0 (66)
Jefferson	14	7.1 (124)	71.4 (115)	14	14.3 (22)	57.1 (126)	2	0.0 (1)	5	0.0 (1)	3	100.0 (45)	5	100.0 (66)
KR	11	9.1 (134)	45.5 (13)	10	30.0 (110)	30.0 (26)	4	0.0 (1)	2	0.0 (1)	1	100.0 (45)	1	100.0 (66)
Keokuk	46	6.5 (116)	71.7 (119)	44	25.0 (88)	36.4 (38)	17	0.0 (1)	11	9.1 (95)	3	100.0 (45)	8	100.0 (66)
Keosauqua	7	0.0 (1)	57.1 (41)	7	0.0 (1)	28.6 (21)	2	50.0 (134)	3	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Knoxville	69	7.2 (125)	60.9 (73)	67	13.4 (20)	47.8 (93)	17	5.9 (89)	7	14.3 (101)	2	100.0 (45)	3	100.0 (66)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lamoni	4	0.0 (1)	50.0 (15)	3	0.0 (1)	66.7 (137)	1	0.0 (1)	0	0.0 (1)	1	100.0 (45)	0	0.0 (1)
Le Mars	35	5.7 (109)	54.3 (29)	34	17.6 (43)	38.2 (48)	9	0.0 (1)	10	10.0 (98)	2	100.0 (45)	0	0.0 (1)
Lenox	9	11.1 (139)	55.6 (33)	9	22.2 (67)	44.4 (75)	3	66.7 (144)	2	50.0 (134)	3	100.0 (45)	0	0.0 (1)
Leon	17	5.9 (111)	58.8 (52)	17	47.1 (140)	23.5 (9)	7	14.3 (100)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Logan	27	0.0 (1)	63.0 (80)	27	22.2 (67)	55.6 (121)	11	0.0 (1)	10	0.0 (1)	0	0.0 (1)	1	100.0 (66)
Manchester	15	6.7 (118)	66.7 (95)	15	33.3 (120)	33.3 (28)	3	66.7 (144)	5	40.0 (126)	2	100.0 (45)	0	0.0 (1)
Manning	2	0.0 (1)	50.0 (15)	2	0.0 (1)	100.0 (147)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Maquoketa	22	9.1 (134)	72.7 (120)	22	22.7 (75)	54.5 (120)	4	25.0 (107)	3	66.7 (144)	0	0.0 (1)	2	100.0 (66)
Marengo	7	0.0 (1)	71.4 (115)	7	14.3 (22)	28.6 (21)	2	0.0 (1)	4	25.0 (110)	0	0.0 (1)	0	0.0 (1)
Marion	95	1.1 (66)	64.2 (88)	92	17.4 (41)	56.5 (124)	29	27.6 (112)	16	0.0 (1)	12	100.0 (45)	5	100.0 (66)
Marshalltown	227	2.6 (77)	59.0 (54)	223	22.9 (78)	41.7 (61)	52	13.5 (99)	16	12.5 (100)	19	100.0 (45)	2	100.0 (66)
Mason City	169	4.7 (104)	63.9 (86)	163	19.0 (47)	46.0 (84)	45	6.7 (91)	22	9.1 (95)	18	100.0 (45)	4	100.0 (66)
Monticello	7	0.0 (1)	57.1 (41)	7	14.3 (22)	42.9 (68)	2	50.0 (134)	3	0.0 (1)	2	100.0 (45)	0	0.0 (1)
Mt Ayr	11	0.0 (1)	54.5 (31)	11	36.4 (131)	36.4 (38)	5	40.0 (128)	2	0.0 (1)	2	100.0 (45)	0	0.0 (1)
Mt Pleasan	38	0.0 (1)	57.9 (47)	37	29.7 (109)	45.9 (82)	15	0.0 (1)	10	0.0 (1)	6	100.0 (45)	5	100.0 (66)
Mtown Dtown	2	50.0 (148)	0.0 (1)	2	100.0 (147)	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Muscatine	176	4.5 (101)	59.1 (56)	166	19.3 (49)	52.4 (114)	54	1.9 (81)	19	5.3 (86)	19	100.0 (45)	5	100.0 (66)
Nevada	4	0.0 (1)	50.0 (15)	4	0.0 (1)	25.0 (11)	2	0.0 (1)	2	0.0 (1)	0	0.0 (1)	0	0.0 (1)
New Hampto	16	6.3 (115)	56.3 (37)	16	25.0 (88)	43.8 (71)	3	0.0 (1)	3	0.0 (1)	2	100.0 (45)	1	100.0 (66)
Newton	72	5.6 (108)	56.9 (40)	67	28.4 (107)	41.8 (65)	26	23.1 (106)	10	50.0 (134)	14	100.0 (45)	2	100.0 (66)
North Towne	105	2.9 (78)	60.0 (63)	101	14.9 (29)	57.4 (128)	43	32.6 (114)	15	40.0 (126)	14	100.0 (45)	2	100.0 (66)
Northwood	7	0.0 (1)	57.1 (41)	7	0.0 (1)	42.9 (68)	1	0.0 (1)	4	0.0 (1)	1	100.0 (45)	1	100.0 (66)
Norwalk	9	0.0 (1)	55.6 (33)	9	33.3 (120)	33.3 (28)	3	0.0 (1)	0	0.0 (1)	1	100.0 (45)	1	100.0 (66)
Oakridge	6	0.0 (1)	16.7 (5)	4	50.0 (141)	25.0 (11)	2	0.0 (1)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Oelwein	26	0.0 (1)	50.0 (15)	26	3.8 (14)	57.7 (129)	11	36.4 (122)	6	50.0 (134)	3	100.0 (45)	3	100.0 (66)
Onawa	24	4.2 (97)	58.3 (50)	24	16.7 (34)	37.5 (43)	9	0.0 (1)	6	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Orange Cty	26	3.8 (91)	50.0 (15)	25	16.0 (31)	12.0 (6)	9	0.0 (1)	12	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Osage	21	0.0 (1)	76.2 (130)	21	14.3 (22)	57.1 (126)	5	0.0 (1)	8	0.0 (1)	4	100.0 (45)	1	100.0 (66)
Osceola	39	10.3 (138)	59.0 (54)	37	32.4 (117)	27.0 (17)	15	26.7 (109)	5	40.0 (126)	3	100.0 (45)	3	100.0 (66)
Oskaloosa	30	6.7 (118)	63.3 (81)	29	24.1 (86)	41.4 (60)	12	25.0 (107)	9	33.3 (115)	0	0.0 (1)	0	0.0 (1)
Ottumwa	348	5.7 (109)	59.5 (61)	326	19.9 (50)	44.5 (76)	133	27.1 (110)	51	29.4 (113)	5	100.0 (45)	13	100.0 (66)
Pella	12	0.0 (1)	75.0 (126)	12	8.3 (16)	66.7 (137)	3	33.3 (115)	3	33.3 (115)	0	0.0 (1)	0	0.0 (1)
Perry	31	3.2 (81)	58.1 (48)	29	17.2 (40)	37.9 (47)	10	0.0 (1)	6	0.0 (1)	3	100.0 (45)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Pocahontas	2	0.0 (1)	100.0 (144)	2	50.0 (141)	50.0 (99)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Postville	30	6.7 (118)	60.0 (63)	29	24.1 (86)	48.3 (95)	8	62.5 (143)	15	40.0 (126)	4	100.0 (45)	0	0.0 (1)
Red Oak	23	0.0 (1)	73.9 (125)	22	22.7 (75)	45.5 (79)	4	0.0 (1)	6	16.7 (105)	0	0.0 (1)	2	100.0 (66)
Redfield	9	0.0 (1)	66.7 (95)	9	22.2 (67)	55.6 (121)	4	0.0 (1)	0	0.0 (1)	1	100.0 (45)	1	100.0 (66)
Refugee Cln	10	0.0 (1)	70.0 (112)	10	30.0 (110)	30.0 (26)	2	50.0 (134)	7	42.9 (131)	1	100.0 (45)	0	0.0 (1)
Rock Rapid	7	0.0 (1)	42.9 (10)	7	57.1 (146)	14.3 (7)	2	0.0 (1)	4	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Rock Valle	26	3.8 (91)	53.8 (25)	25	16.0 (31)	44.0 (73)	6	0.0 (1)	14	7.1 (90)	3	100.0 (45)	1	100.0 (66)
Rockwell C	17	5.9 (111)	64.7 (90)	17	23.5 (82)	47.1 (89)	6	16.7 (102)	6	33.3 (115)	1	100.0 (45)	0	0.0 (1)
SDM	322	5.0 (105)	64.0 (87)	307	22.8 (77)	45.9 (82)	102	2.0 (82)	34	5.9 (88)	18	100.0 (45)	14	100.0 (66)
Sac City	12	8.3 (129)	75.0 (126)	12	16.7 (34)	50.0 (99)	2	0.0 (1)	2	0.0 (1)	1	100.0 (45)	1	100.0 (66)
Sheldon	26	3.8 (91)	53.8 (25)	24	25.0 (88)	58.3 (130)	7	0.0 (1)	7	0.0 (1)	2	100.0 (45)	0	0.0 (1)
Shenandoah	24	0.0 (1)	70.8 (114)	24	25.0 (88)	37.5 (43)	8	37.5 (126)	9	44.4 (132)	2	100.0 (45)	0	0.0 (1)
Sibley	6	0.0 (1)	83.3 (139)	5	20.0 (52)	40.0 (52)	3	0.0 (1)	4	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Sidney	5	0.0 (1)	60.0 (63)	4	25.0 (88)	25.0 (11)	2	0.0 (1)	0	0.0 (1)	1	100.0 (45)	0	0.0 (1)
Sigourney	8	0.0 (1)	37.5 (9)	8	25.0 (88)	25.0 (11)	3	66.7 (144)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Sioux Cent	28	0.0 (1)	57.1 (41)	28	46.4 (139)	28.6 (21)	16	6.3 (90)	13	0.0 (1)	3	100.0 (45)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Sioux City	525	1.3 (68)	63.8 (85)	510	22.4 (74)	45.7 (81)	190	4.7 (88)	164	6.1 (89)	37	100.0 (45)	15	100.0 (66)
Spencer	45	2.2 (73)	53.3 (24)	39	23.1 (80)	48.7 (97)	13	0.0 (1)	10	0.0 (1)	6	100.0 (45)	3	100.0 (66)
Spirit Lak	16	0.0 (1)	81.3 (138)	15	20.0 (52)	66.7 (137)	6	0.0 (1)	2	0.0 (1)	2	100.0 (45)	2	100.0 (66)
Storm Lake	71	1.4 (69)	63.4 (82)	67	26.9 (105)	35.8 (37)	29	0.0 (1)	10	20.0 (106)	13	100.0 (45)	7	100.0 (66)
Stuart	8	0.0 (1)	62.5 (79)	8	0.0 (1)	62.5 (134)	5	20.0 (103)	5	60.0 (141)	0	0.0 (1)	0	0.0 (1)
Tama	33	3.0 (80)	63.6 (83)	30	26.7 (102)	43.3 (70)	10	40.0 (128)	1	0.0 (1)	2	100.0 (45)	1	100.0 (66)
Tama/Settl	2	0.0 (1)	50.0 (15)	2	0.0 (1)	0.0 (1)	2	0.0 (1)	1	100.0 (145)	1	100.0 (45)	0	0.0 (1)
Tipton	6	16.7 (146)	50.0 (15)	5	40.0 (136)	40.0 (52)	2	50.0 (134)	4	0.0 (1)	1	100.0 (45)	0	0.0 (1)
Traer	2	0.0 (1)	100.0 (144)	2	0.0 (1)	50.0 (99)	1	0.0 (1)	0	0.0 (1)	1	100.0 (45)	0	0.0 (1)
UIHC	1	0.0 (1)	0.0 (1)	1	100.0 (147)	0.0 (1)	0	0.0 (1)	1	100.0 (145)	2	100.0 (45)	1	100.0 (66)
Vinton	21	0.0 (1)	71.4 (115)	20	40.0 (136)	35.0 (35)	4	0.0 (1)	5	0.0 (1)	2	100.0 (45)	2	100.0 (66)
WDM	50	2.0 (71)	66.0 (94)	48	20.8 (61)	43.8 (71)	19	0.0 (1)	10	10.0 (98)	2	100.0 (45)	5	100.0 (66)
Washington	26	7.7 (126)	53.8 (25)	24	33.3 (120)	29.2 (25)	10	10.0 (97)	7	14.3 (101)	3	100.0 (45)	0	0.0 (1)
Waterloo	491	3.7 (87)	61.9 (76)	472	19.9 (50)	52.3 (113)	136	4.4 (87)	61	1.6 (84)	32	100.0 (45)	29	100.0 (66)
Waukee	25	12.0 (143)	60.0 (63)	22	9.1 (17)	50.0 (99)	8	0.0 (1)	4	0.0 (1)	2	100.0 (45)	2	100.0 (66)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Waukon	17	0.0 (1)	70.6 (113)	16	18.8 (46)	50.0 (99)	6	33.3 (115)	3	33.3 (115)	1	100.0 (45)	0	0.0 (1)
Waverly	15	6.7 (118)	66.7 (95)	15	13.3 (19)	46.7 (87)	2	0.0 (1)	6	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Wbstr Ct	51	2.0 (71)	60.8 (72)	50	20.0 (52)	40.0 (52)	14	7.1 (92)	2	0.0 (1)	5	100.0 (45)	4	100.0 (66)
West Union	13	7.7 (126)	61.5 (74)	12	16.7 (34)	41.7 (61)	7	28.6 (113)	5	60.0 (141)	2	100.0 (45)	2	100.0 (66)
Winterset	15	0.0 (1)	80.0 (135)	15	26.7 (102)	40.0 (52)	3	66.7 (144)	5	60.0 (141)	4	100.0 (45)	0	0.0 (1)
Iowa	**8,677**	**3.8**	**61.0**	**8,349**	**23.1**	**45.1**	**2,701**	**15.5**	**1,557**	**14.7**	**655**	**100.0**	**376**	**100.0**
National (Prior Year)	**112,231**	**4.0**	**60.3**	**89,229**	**22.3**	**47.7**	**23,550**	**28.2**	**52,310**	**18.4**	**96,728**	**10.0**	**95,873**	**9.4**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley	3	33.3 (49)	33.3 (144)	3	0.0 (37)	2	0.0 (114)	3	66.7 (137)
Adel	5	60.0 (8)	0.0 (1)	7	0.0 (37)	3	66.7 (7)	3	100.0 (1)
Albia	10	60.0 (8)	0.0 (1)	10	0.0 (37)	4	25.0 (62)	4	100.0 (1)
Algona	40	50.0 (12)	7.5 (76)	42	0.0 (37)	36	19.4 (77)	38	97.4 (69)
Allen WHC	58	36.2 (41)	31.0 (140)	96	0.0 (37)	48	18.8 (81)	68	88.2 (111)
Allison	18	27.8 (83)	5.6 (60)	19	0.0 (37)	14	14.3 (99)	18	88.9 (106)
Altoona	26	30.8 (65)	7.7 (77)	32	0.0 (37)	15	33.3 (43)	16	100.0 (1)
Ames	107	31.8 (63)	11.2 (93)	159	2.5 (16)	83	25.3 (60)	82	90.2 (98)
Anamosa	20	25.0 (99)	0.0 (1)	24	4.2 (7)	13	46.2 (23)	20	95.0 (73)
Ankeny	30	26.7 (94)	6.7 (72)	46	0.0 (37)	10	20.0 (75)	12	91.7 (94)
Atlantic	25	36.0 (42)	4.0 (48)	34	0.0 (37)	3	66.7 (7)	4	100.0 (1)
Audubon	7	57.1 (10)	0.0 (1)	9	0.0 (37)	2	0.0 (114)	2	100.0 (1)
Avoca	5	0.0 (137)	40.0 (147)	7	0.0 (37)	4	0.0 (114)	4	100.0 (1)
Bedford	5	20.0 (116)	0.0 (1)	7	0.0 (37)	2	0.0 (114)	2	100.0 (1)
Belle Plaine	4	25.0 (99)	0.0 (1)	8	0.0 (37)	2	0.0 (114)	5	80.0 (129)
Bloomfield	5	40.0 (27)	0.0 (1)	9	11.1 (2)	1	0.0 (114)	1	100.0 (1)
Boone	17	29.4 (75)	0.0 (1)	20	0.0 (37)	9	44.4 (26)	9	88.9 (106)
Brdlwns RP	798	25.8 (97)	14.9 (110)	1,036	0.3 (35)	399	15.0 (97)	474	86.9 (113)
Brooklyn	2	50.0 (12)	50.0 (148)	3	0.0 (37)	1	100.0 (1)	2	100.0 (1)
Burlington	132	27.3 (88)	3.0 (45)	146	1.4 (25)	107	25.2 (61)	131	94.7 (76)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
CHC	171	25.1 (98)	19.3 (123)	224	2.7 (15)	7	0.0 (114)	8	100.0 (1)
CR Urban	335	26.9 (91)	10.1 (89)	407	0.2 (36)	211	10.9 (106)	285	89.5 (105)
Carlisle	4	50.0 (12)	25.0 (130)	7	0.0 (37)	2	0.0 (114)	3	100.0 (1)
Carroll	53	22.6 (113)	5.7 (64)	70	0.0 (37)	19	10.5 (107)	20	90.0 (99)
Cedar Falls	4	75.0 (3)	25.0 (130)	9	0.0 (37)	3	0.0 (114)	4	75.0 (135)
Centervill	15	53.3 (11)	26.7 (135)	25	4.0 (10)	2	0.0 (114)	2	100.0 (1)
Chariton	18	33.3 (49)	5.6 (60)	21	0.0 (37)	14	35.7 (42)	14	64.3 (141)
Charles City	46	39.1 (32)	13.0 (102)	54	0.0 (37)	38	21.1 (74)	48	89.6 (103)
Cherokee	18	38.9 (33)	5.6 (60)	35	0.0 (37)	11	36.4 (40)	13	100.0 (1)
Clarinda	20	50.0 (12)	5.0 (55)	24	0.0 (37)	8	50.0 (14)	7	100.0 (1)
Clarion	35	17.1 (128)	31.4 (142)	41	0.0 (37)	30	3.3 (113)	33	81.8 (126)
Clinton	127	37.0 (39)	8.7 (83)	136	0.7 (31)	57	36.8 (38)	109	86.2 (117)
Colfax	6	33.3 (49)	16.7 (113)	7	0.0 (37)	4	25.0 (62)	4	100.0 (1)
Columbus J	14	35.7 (43)	7.1 (75)	14	0.0 (37)	11	36.4 (40)	11	81.8 (126)
Corning	9	33.3 (49)	0.0 (1)	14	0.0 (37)	6	0.0 (114)	7	100.0 (1)
Corydon	13	46.2 (20)	7.7 (77)	15	0.0 (37)	9	44.4 (26)	9	100.0 (1)
Council Blfs	283	35.7 (43)	5.7 (64)	366	0.0 (37)	114	17.5 (84)	116	88.8 (110)
Cresco	20	30.0 (72)	15.0 (111)	26	0.0 (37)	22	18.2 (82)	22	81.8 (126)
Creston	47	44.7 (24)	8.5 (81)	65	0.0 (37)	23	21.7 (73)	27	92.6 (86)
Dallas Ctr	0	0.0 (137)	0.0 (1)	2	0.0 (37)	0	0.0 (114)	0	0.0 (142)
Decorah	13	23.1 (110)	15.4 (112)	17	0.0 (37)	13	30.8 (50)	13	100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Denison	76	25.0 (99)	7.9 (79)	99	0.0 (37)	19	26.3 (58)	28	92.9 (85)
Dewitt	5	20.0 (116)	0.0 (1)	5	0.0 (37)	3	66.7 (7)	5	100.0 (1)
Dubuque	293	37.5 (35)	11.9 (98)	317	1.9 (19)	266	16.2 (91)	269	93.3 (82)
Dyersville	7	28.6 (78)	14.3 (104)	7	0.0 (37)	7	14.3 (99)	7	100.0 (1)
Eagle Grove	26	23.1 (110)	19.2 (122)	28	0.0 (37)	21	19.0 (79)	25	92.0 (93)
Edgerton	249	32.5 (60)	8.8 (84)	345	1.4 (25)	17	11.8 (105)	17	100.0 (1)
Eldora	4	25.0 (99)	0.0 (1)	5	0.0 (37)	2	0.0 (114)	2	100.0 (1)
Elkader	23	30.4 (68)	13.0 (102)	27	0.0 (37)	23	39.1 (35)	24	100.0 (1)
Emmetsburg	8	37.5 (35)	12.5 (100)	8	0.0 (37)	7	71.4 (6)	8	100.0 (1)
Esthervill	24	29.2 (76)	8.3 (80)	31	0.0 (37)	21	47.6 (22)	22	90.9 (96)
Fairfiel	9	33.3 (49)	11.1 (92)	13	0.0 (37)	1	0.0 (114)	2	100.0 (1)
Forest City	51	35.3 (46)	11.8 (96)	56	1.8 (20)	51	23.5 (64)	52	84.6 (121)
Fort Dodge	167	27.5 (86)	24.0 (129)	201	2.0 (18)	125	16.0 (93)	165	82.4 (125)
Ft Madison	37	27.0 (90)	0.0 (1)	41	0.0 (37)	36	16.7 (87)	40	92.5 (87)
Garner	20	40.0 (27)	25.0 (130)	21	0.0 (37)	19	15.8 (95)	19	78.9 (131)
Glenwood	9	11.1 (134)	0.0 (1)	18	0.0 (37)	1	100.0 (1)	2	100.0 (1)
Grand Junction	2	50.0 (12)	0.0 (1)	3	0.0 (37)	3	33.3 (43)	3	66.7 (137)
Greenfield	14	35.7 (43)	0.0 (1)	18	0.0 (37)	7	14.3 (99)	6	83.3 (122)
Grinnell	21	33.3 (49)	19.0 (120)	24	8.3 (3)	13	15.4 (96)	13	92.3 (88)
Grundy Ctr	4	0.0 (137)	25.0 (130)	8	0.0 (37)	4	50.0 (14)	4	100.0 (1)
Guthrie Ct	8	62.5 (6)	0.0 (1)	14	0.0 (37)	2	0.0 (114)	2	100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Hampton	29	27.6 (85)	31.0 (140)	42	0.0 (37)	29	27.6 (54)	36	83.3 (122)
Harlan	21	19.0 (124)	9.5 (86)	35	0.0 (37)	1	100.0 (1)	3	100.0 (1)
Hawarden	11	18.2 (126)	18.2 (116)	17	0.0 (37)	5	40.0 (31)	6	100.0 (1)
Hispanic Cln	6	16.7 (129)	0.0 (1)	6	0.0 (37)	4	0.0 (114)	5	100.0 (1)
Humboldt	15	26.7 (94)	26.7 (135)	16	0.0 (37)	16	43.8 (28)	16	100.0 (1)
ILH	253	33.2 (59)	11.9 (98)	350	0.6 (32)	100	10.0 (108)	195	89.7 (102)
Ida Grove	20	45.0 (23)	5.0 (55)	24	4.2 (7)	9	22.2 (70)	9	100.0 (1)
Independen	19	47.4 (17)	10.5 (90)	26	0.0 (37)	14	28.6 (51)	18	88.9 (106)
Indianola	29	24.1 (107)	34.5 (146)	34	0.0 (37)	22	27.3 (55)	23	87.0 (112)
Iowa City	225	18.7 (125)	19.1 (121)	317	2.2 (17)	80	22.5 (68)	91	92.3 (88)
Iowa Falls	8	62.5 (6)	0.0 (1)	12	0.0 (37)	5	0.0 (114)	5	100.0 (1)
Jamaica	3	0.0 (137)	0.0 (1)	3	0.0 (37)	1	0.0 (114)	1	100.0 (1)
Jefferson	12	25.0 (99)	25.0 (130)	15	0.0 (37)	3	33.3 (43)	3	100.0 (1)
KR	8	25.0 (99)	12.5 (100)	14	0.0 (37)	7	42.9 (29)	10	90.0 (99)
Keokuk	43	27.9 (82)	2.3 (44)	49	0.0 (37)	37	16.2 (91)	43	97.7 (68)
Keosauqua	6	33.3 (49)	0.0 (1)	6	0.0 (37)	0	0.0 (114)	0	0.0 (142)
Knoxville	56	26.8 (93)	8.9 (85)	72	1.4 (25)	41	26.8 (57)	43	93.0 (84)
Lamoni	4	25.0 (99)	0.0 (1)	5	0.0 (37)	2	0.0 (114)	2	100.0 (1)
Le Mars	26	15.4 (131)	3.8 (47)	41	0.0 (37)	15	20.0 (75)	15	93.3 (82)
Lenox	10	40.0 (27)	10.0 (88)	12	0.0 (37)	5	60.0 (12)	5	80.0 (129)
Leon	17	23.5 (108)	5.9 (66)	18	0.0 (37)	9	22.2 (70)	8	100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Logan	21	33.3 (49)	4.8 (52)	32	0.0 (37)	3	0.0 (114)	3	66.7 (137)
Manchester	14	21.4 (115)	14.3 (104)	15	0.0 (37)	14	7.1 (110)	14	100.0 (1)
Manning	1	100.0 (1)	0.0 (1)	2	0.0 (37)	0	0.0 (114)	0	0.0 (142)
Maquoketa	21	38.1 (34)	4.8 (52)	23	0.0 (37)	11	45.5 (25)	19	94.7 (76)
Marengo	4	25.0 (99)	0.0 (1)	10	0.0 (37)	1	0.0 (114)	1	100.0 (1)
Marion	94	28.7 (77)	8.5 (81)	115	0.0 (37)	73	17.8 (83)	78	94.9 (75)
Marshalltown	189	33.9 (48)	6.3 (68)	242	0.4 (34)	146	17.1 (86)	153	86.3 (116)
Mason City	159	39.6 (31)	14.5 (108)	183	0.5 (33)	134	17.2 (85)	162	85.8 (118)
Monticello	7	42.9 (25)	0.0 (1)	10	0.0 (37)	3	0.0 (114)	7	100.0 (1)
Mt Ayr	9	11.1 (134)	0.0 (1)	11	0.0 (37)	6	16.7 (87)	6	100.0 (1)
Mt Pleasan	36	30.6 (67)	5.6 (60)	42	0.0 (37)	31	12.9 (103)	37	86.5 (114)
Mtown Dtown	1	0.0 (137)	0.0 (1)	2	0.0 (37)	0	0.0 (114)	0	0.0 (142)
Muscatine	175	32.0 (62)	6.3 (68)	185	1.6 (22)	181	19.3 (78)	185	96.2 (71)
Nevada	3	66.7 (4)	33.3 (144)	4	0.0 (37)	2	50.0 (14)	2	100.0 (1)
New Hampto	11	45.5 (21)	0.0 (1)	14	0.0 (37)	11	72.7 (5)	13	92.3 (88)
Newton	61	23.0 (112)	11.5 (94)	80	0.0 (37)	43	14.0 (102)	48	89.6 (103)
North Towne	99	19.2 (123)	4.0 (48)	125	1.6 (22)	50	26.0 (59)	88	90.9 (96)
Northwood	6	33.3 (49)	16.7 (113)	8	0.0 (37)	5	40.0 (31)	7	100.0 (1)
Norwalk	7	28.6 (78)	14.3 (104)	8	0.0 (37)	5	40.0 (31)	5	100.0 (1)
Oakridge	5	20.0 (116)	0.0 (1)	8	0.0 (37)	2	50.0 (14)	3	100.0 (1)
Oelwein	21	28.6 (78)	4.8 (52)	27	0.0 (37)	25	28.0 (53)	25	96.0 (72)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Onawa	17	17.6 (127)	0.0 (1)	25	0.0 (37)	2	50.0 (14)	4	100.0 (1)
Orange Cty	22	31.8 (63)	4.5 (51)	36	2.8 (14)	6	33.3 (43)	7	100.0 (1)
Osage	23	34.8 (47)	21.7 (128)	25	0.0 (37)	19	36.8 (38)	23	91.3 (95)
Osceola	37	21.6 (114)	10.8 (91)	46	0.0 (37)	20	15.0 (97)	20	95.0 (73)
Oskaloosa	24	37.5 (35)	4.2 (50)	36	0.0 (37)	5	40.0 (31)	5	100.0 (1)
Ottumwa	255	32.5 (60)	28.6 (137)	367	1.6 (22)	30	23.3 (65)	30	76.7 (134)
Pella	7	42.9 (25)	28.6 (137)	12	0.0 (37)	5	0.0 (114)	6	100.0 (1)
Perry	30	20.0 (116)	0.0 (1)	34	2.9 (12)	18	5.6 (111)	18	100.0 (1)
Pocahontas	1	100.0 (1)	0.0 (1)	2	0.0 (37)	1	0.0 (114)	1	100.0 (1)
Postville	28	14.3 (133)	32.1 (143)	33	0.0 (37)	25	16.0 (93)	27	85.2 (120)
Red Oak	17	23.5 (108)	11.8 (96)	26	0.0 (37)	2	0.0 (114)	4	100.0 (1)
Redfield	7	28.6 (78)	0.0 (1)	8	0.0 (37)	3	66.7 (7)	4	100.0 (1)
Refugee Cln	9	0.0 (137)	0.0 (1)	12	0.0 (37)	7	0.0 (114)	8	75.0 (135)
Rock Rapid	5	20.0 (116)	20.0 (124)	12	0.0 (37)	4	50.0 (14)	5	100.0 (1)
Rock Valle	21	33.3 (49)	9.5 (86)	34	2.9 (12)	8	12.5 (104)	9	77.8 (132)
Rockwell C	17	47.1 (18)	5.9 (66)	20	0.0 (37)	6	50.0 (14)	6	100.0 (1)
SDM	235	27.2 (89)	14.5 (108)	351	1.1 (28)	79	19.0 (79)	104	86.5 (114)
Sac City	10	30.0 (72)	20.0 (124)	10	0.0 (37)	6	16.7 (87)	6	100.0 (1)
Sheldon	26	26.9 (91)	11.5 (94)	32	3.1 (11)	26	50.0 (14)	28	96.4 (70)
Shenandoah	27	37.0 (39)	3.7 (46)	28	0.0 (37)	16	37.5 (36)	14	100.0 (1)
Sibley	6	16.7 (129)	0.0 (1)	12	0.0 (37)	6	66.7 (7)	6	100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Sidney	3	0.0 (137)	0.0 (1)	5	20.0 (1)	0	0.0 (114)	0	0.0 (142)
Sigourney	5	40.0 (27)	20.0 (124)	7	0.0 (37)	1	0.0 (114)	1	100.0 (1)
Sioux Cent	20	10.0 (136)	5.0 (55)	35	0.0 (37)	10	0.0 (114)	11	100.0 (1)
Sioux City	460	30.2 (70)	6.5 (71)	614	0.8 (30)	79	31.6 (49)	107	94.4 (78)
Spencer	43	30.2 (70)	7.0 (74)	49	4.1 (9)	46	45.7 (24)	48	93.8 (79)
Spirit Lak	15	20.0 (116)	6.7 (72)	19	0.0 (37)	14	57.1 (13)	15	100.0 (1)
Storm Lake	60	26.7 (94)	5.0 (55)	86	0.0 (37)	60	41.7 (30)	63	93.7 (81)
Stuart	7	0.0 (137)	14.3 (104)	10	0.0 (37)	3	33.3 (43)	3	100.0 (1)
Tama	23	30.4 (68)	0.0 (1)	35	5.7 (6)	21	9.5 (109)	22	77.3 (133)
Tama/Settl	1	0.0 (137)	0.0 (1)	2	0.0 (37)	0	0.0 (114)	0	0.0 (142)
Tipton	2	0.0 (137)	0.0 (1)	8	0.0 (37)	1	0.0 (114)	1	100.0 (1)
Traer	1	0.0 (137)	0.0 (1)	2	0.0 (37)	1	100.0 (1)	1	100.0 (1)
UIHC	1	0.0 (137)	0.0 (1)	2	0.0 (37)	0	0.0 (114)	0	0.0 (142)
Vinton	15	46.7 (19)	0.0 (1)	23	0.0 (37)	9	22.2 (70)	10	100.0 (1)
WDM	40	30.0 (72)	5.0 (55)	56	0.0 (37)	11	27.3 (55)	16	93.8 (79)
Washington	27	14.8 (132)	18.5 (119)	30	0.0 (37)	12	16.7 (87)	14	85.7 (119)
Waterloo	314	27.7 (84)	16.9 (115)	533	1.1 (28)	202	23.3 (65)	291	90.0 (99)
Waukee	22	45.5 (21)	18.2 (116)	30	0.0 (37)	12	33.3 (43)	13	92.3 (88)
Waukon	16	37.5 (35)	6.3 (68)	17	5.9 (5)	18	5.6 (111)	18	88.9 (106)
Waverly	11	63.6 (5)	18.2 (116)	17	0.0 (37)	8	37.5 (36)	10	100.0 (1)
Wbstr Ct	51	27.5 (86)	29.4 (139)	56	1.8 (20)	40	22.5 (68)	46	82.6 (124)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
West Union	13	30.8 (65)	0.0 (1)	14	7.1 (4)	13	23.1 (67)	13	92.3 (88)
Winterset	15	20.0 (116)	20.0 (124)	17	0.0 (37)	7	28.6 (51)	6	66.7 (137)
Iowa	**7,379**	**29.9**	**12.1**	**9,673**	**0.9**	**4,023**	**21.3**	**4,805**	**4805.0**
National (Prior Year)	**99,175**	**68.5**	**7.4**	**127,401**	**22.9**	**94,432**	**19.4**	**93,223**	**87.9**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley	3	33.3 (84)	3	33.3 (124)	1	0.0 (100)	3	0.0 (1)	3	0.0 (1)
Adel	3	0.0 (1)	3	33.3 (124)	0	0.0 (100)	3	0.0 (1)	4	0.0 (1)
Albia	8	50.0 (127)	7	42.9 (137)	4	0.0 (100)	8	12.5 (119)	7	0.0 (1)
Algona	41	9.8 (17)	46	10.9 (45)	4	50.0 (11)	40	5.0 (71)	40	0.0 (1)
Allen WHC	74	32.4 (82)	54	22.2 (102)	23	34.8 (38)	75	13.3 (126)	57	0.0 (1)
Allison	19	26.3 (63)	22	22.7 (108)	5	20.0 (76)	19	0.0 (1)	19	0.0 (1)
Altoona	22	27.3 (65)	27	11.1 (48)	5	20.0 (76)	23	0.0 (1)	30	0.0 (1)
Ames	135	24.4 (55)	134	14.2 (63)	30	36.7 (37)	141	7.1 (86)	125	0.8 (91)
Anamosa	19	57.9 (138)	17	41.2 (136)	11	0.0 (100)	21	4.8 (68)	24	8.3 (129)
Ankeny	24	20.8 (44)	33	9.1 (37)	6	16.7 (84)	25	4.0 (62)	33	0.0 (1)
Atlantic	25	40.0 (105)	11	45.5 (140)	10	20.0 (76)	25	0.0 (1)	10	0.0 (1)
Audubon	7	14.3 (25)	9	22.2 (102)	2	0.0 (100)	7	14.3 (127)	9	11.1 (136)
Avoca	6	33.3 (84)	3	33.3 (124)	3	33.3 (40)	6	16.7 (131)	3	0.0 (1)
Bedford	6	83.3 (145)	6	16.7 (72)	4	25.0 (64)	6	33.3 (146)	6	16.7 (143)
Belle Plaine	5	20.0 (39)	4	25.0 (112)	1	0.0 (100)	5	0.0 (1)	4	0.0 (1)
Bloomfield	5	0.0 (1)	5	20.0 (92)	1	0.0 (100)	5	0.0 (1)	3	33.3 (146)
Boone	21	57.1 (136)	20	55.0 (146)	12	8.3 (97)	21	0.0 (1)	19	0.0 (1)
Brdlwns RP	749	18.0 (35)	750	12.5 (54)	139	25.9 (61)	791	5.1 (73)	804	1.1 (97)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Brooklyn	1	100.0 (146)	3	33.3 (124)	1	0.0 (100)	2	0.0 (1)	2	0.0 (1)
Burlington	137	48.2 (126)	158	22.2 (102)	65	43.1 (26)	136	11.8 (115)	157	3.2 (113)
CHC	184	35.3 (97)	141	19.9 (91)	50	18.0 (82)	183	9.8 (106)	138	6.5 (126)
CR Urban	337	27.0 (64)	340	14.1 (62)	98	31.6 (51)	353	6.5 (84)	405	1.2 (98)
Carlisle	7	14.3 (25)	3	33.3 (124)	0	0.0 (100)	7	0.0 (1)	3	33.3 (146)
Carroll	55	41.8 (114)	67	19.4 (90)	22	45.5 (25)	54	1.9 (57)	66	3.0 (111)
Cedar Falls	6	16.7 (28)	9	0.0 (1)	0	0.0 (100)	6	16.7 (131)	10	0.0 (1)
Centervill	17	52.9 (133)	14	35.7 (133)	9	0.0 (100)	16	12.5 (119)	10	0.0 (1)
Chariton	20	25.0 (56)	17	11.8 (52)	5	40.0 (31)	20	5.0 (71)	16	0.0 (1)
Charles City	50	34.0 (94)	64	17.2 (77)	17	41.2 (29)	49	2.0 (58)	64	0.0 (1)
Cherokee	20	45.0 (121)	31	25.8 (115)	9	0.0 (100)	20	20.0 (138)	31	9.7 (134)
Clarinda	20	45.0 (121)	18	27.8 (121)	9	22.2 (73)	20	15.0 (129)	16	6.3 (124)
Clarion	38	7.9 (13)	47	12.8 (56)	4	25.0 (64)	38	7.9 (90)	46	2.2 (105)
Clinton	134	44.8 (120)	152	20.4 (97)	52	30.8 (52)	134	6.0 (80)	152	0.0 (1)
Colfax	4	50.0 (127)	7	14.3 (64)	2	0.0 (100)	4	0.0 (1)	7	0.0 (1)
Columbus J	14	0.0 (1)	15	0.0 (1)	0	0.0 (100)	14	0.0 (1)	15	0.0 (1)
Corning	9	33.3 (84)	13	30.8 (122)	4	0.0 (100)	9	11.1 (113)	12	8.3 (129)
Corydon	9	22.2 (48)	8	37.5 (134)	3	0.0 (100)	10	0.0 (1)	8	0.0 (1)
Council Blfs	294	39.5 (104)	279	22.2 (102)	111	26.1 (60)	293	6.1 (82)	269	0.4 (90)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Cresco	26	34.6 (96)	22	27.3 (119)	7	28.6 (55)	26	15.4 (130)	22	4.5 (120)
Creston	51	31.4 (78)	54	14.8 (66)	13	7.7 (98)	51	9.8 (106)	48	0.0 (1)
Dallas Ctr	0	0.0 (1)	2	0.0 (1)	0	0.0 (100)	0	0.0 (1)	2	0.0 (1)
Decorah	15	33.3 (84)	20	15.0 (67)	5	40.0 (31)	14	21.4 (142)	20	10.0 (135)
Denison	82	11.0 (20)	68	7.4 (33)	9	22.2 (73)	82	0.0 (1)	67	0.0 (1)
Dewitt	3	66.7 (141)	6	50.0 (141)	2	0.0 (100)	3	33.3 (146)	5	0.0 (1)
Dubuque	311	14.1 (24)	327	7.0 (31)	43	25.6 (63)	308	0.0 (1)	323	0.0 (1)
Dyersville	7	28.6 (67)	7	0.0 (1)	2	0.0 (100)	7	0.0 (1)	7	0.0 (1)
Eagle Grove	28	28.6 (67)	26	15.4 (68)	7	42.9 (27)	28	3.6 (61)	26	3.8 (118)
Edgerton	260	23.5 (53)	175	13.7 (60)	52	17.3 (83)	259	9.3 (102)	171	2.3 (107)
Eldora	5	40.0 (105)	5	60.0 (148)	2	0.0 (100)	5	0.0 (1)	5	0.0 (1)
Elkader	25	44.0 (119)	23	26.1 (117)	13	30.8 (52)	24	20.8 (141)	23	13.0 (138)
Emmetsburg	8	12.5 (22)	12	16.7 (72)	0	0.0 (100)	8	12.5 (119)	12	25.0 (144)
Esthervill	23	8.7 (15)	39	10.3 (43)	2	50.0 (11)	23	13.0 (125)	36	13.9 (139)
Fairfiel	11	54.5 (135)	7	14.3 (64)	7	28.6 (55)	11	9.1 (98)	6	0.0 (1)
Forest City	52	25.0 (56)	55	10.9 (45)	12	50.0 (11)	48	2.1 (59)	53	1.9 (103)
Fort Dodge	189	37.0 (100)	212	18.9 (86)	71	39.4 (34)	186	8.1 (93)	211	1.9 (103)
Ft Madison	39	41.0 (112)	53	18.9 (86)	15	40.0 (31)	39	5.1 (73)	53	0.0 (1)
Garner	20	20.0 (39)	26	15.4 (68)	4	50.0 (11)	19	10.5 (110)	23	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Glenwood	10	40.0 (105)	10	0.0 (1)	3	66.7 (5)	9	0.0 (1)	10	0.0 (1)
Grand Junction	3	100.0 (146)	2	0.0 (1)	2	100.0 (1)	3	0.0 (1)	2	0.0 (1)
Greenfield	17	23.5 (53)	11	9.1 (37)	3	33.3 (40)	16	0.0 (1)	11	0.0 (1)
Grinnell	21	19.0 (38)	23	13.0 (58)	3	33.3 (40)	21	19.0 (137)	22	0.0 (1)
Grundy Ctr	4	0.0 (1)	7	0.0 (1)	0	0.0 (100)	4	0.0 (1)	7	0.0 (1)
Guthrie Ct	8	37.5 (101)	15	20.0 (92)	3	66.7 (5)	8	12.5 (119)	15	0.0 (1)
Hampton	37	21.6 (46)	43	14.0 (61)	8	25.0 (64)	38	0.0 (1)	40	2.5 (109)
Harlan	23	43.5 (117)	23	34.8 (132)	12	16.7 (84)	23	8.7 (96)	23	0.0 (1)
Hawarden	13	23.1 (50)	15	6.7 (28)	3	66.7 (5)	13	0.0 (1)	15	0.0 (1)
Hispanic Cln	6	0.0 (1)	6	0.0 (1)	0	0.0 (100)	6	0.0 (1)	6	0.0 (1)
Humboldt	17	47.1 (125)	19	21.1 (100)	8	50.0 (11)	17	11.8 (115)	17	0.0 (1)
ILH	156	23.1 (50)	117	23.9 (111)	42	16.7 (84)	275	5.5 (76)	320	1.6 (101)
Ida Grove	22	31.8 (80)	22	18.2 (80)	6	16.7 (84)	21	9.5 (103)	22	9.1 (133)
Independen	22	50.0 (127)	22	22.7 (108)	10	10.0 (94)	21	4.8 (68)	19	0.0 (1)
Indianola	23	43.5 (117)	30	16.7 (72)	10	30.0 (54)	25	12.0 (117)	30	3.3 (114)
Iowa City	234	11.5 (21)	196	3.6 (24)	19	36.8 (36)	232	2.6 (60)	187	0.0 (1)
Iowa Falls	8	25.0 (56)	11	9.1 (37)	2	50.0 (11)	9	22.2 (143)	9	0.0 (1)
Jamaica	3	33.3 (84)	3	0.0 (1)	1	0.0 (100)	3	0.0 (1)	3	0.0 (1)
Jefferson	13	46.2 (123)	17	23.5 (110)	7	42.9 (27)	13	23.1 (144)	17	5.9 (122)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
KR	11	18.2 (36)	10	10.0 (41)	2	50.0 (11)	11	9.1 (98)	8	0.0 (1)
Keokuk	43	41.9 (115)	51	27.5 (120)	20	25.0 (64)	43	11.6 (114)	50	0.0 (1)
Keosauqua	6	33.3 (84)	3	0.0 (1)	2	0.0 (100)	6	16.7 (131)	3	0.0 (1)
Knoxville	47	46.8 (124)	42	21.4 (101)	18	16.7 (84)	56	10.7 (111)	41	0.0 (1)
Lamoni	5	40.0 (105)	2	0.0 (1)	1	0.0 (100)	5	0.0 (1)	2	0.0 (1)
Le Mars	27	29.6 (73)	37	8.1 (36)	7	71.4 (4)	27	0.0 (1)	36	2.8 (110)
Lenox	8	25.0 (56)	13	0.0 (1)	2	50.0 (11)	9	0.0 (1)	11	0.0 (1)
Leon	17	35.3 (97)	16	12.5 (54)	5	20.0 (76)	17	0.0 (1)	14	7.1 (128)
Logan	21	28.6 (67)	21	19.0 (88)	4	25.0 (64)	21	9.5 (103)	20	0.0 (1)
Manchester	14	35.7 (99)	13	15.4 (68)	5	0.0 (100)	14	0.0 (1)	13	0.0 (1)
Manning	1	0.0 (1)	2	0.0 (1)	0	0.0 (100)	1	0.0 (1)	2	0.0 (1)
Maquoketa	23	65.2 (140)	27	25.9 (116)	14	28.6 (55)	22	9.1 (98)	28	3.6 (116)
Marengo	5	60.0 (139)	7	57.1 (147)	4	25.0 (64)	5	20.0 (138)	7	0.0 (1)
Marion	93	28.0 (66)	123	17.9 (79)	34	41.2 (29)	101	7.9 (90)	122	0.8 (91)
Marshalltown	208	10.1 (19)	221	10.9 (45)	22	4.5 (99)	214	4.2 (64)	250	0.8 (91)
Mason City	169	32.0 (81)	203	18.7 (84)	54	22.2 (73)	168	4.8 (68)	191	1.0 (95)
Monticello	7	14.3 (25)	8	0.0 (1)	1	0.0 (100)	7	0.0 (1)	9	0.0 (1)
Mt Ayr	8	37.5 (101)	8	25.0 (112)	3	33.3 (40)	8	0.0 (1)	8	0.0 (1)
Mt Pleasan	37	29.7 (74)	53	17.0 (76)	11	27.3 (58)	37	8.1 (93)	52	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Mtown Dtown	2	0.0 (1)	0	0.0 (1)	0	0.0 (100)	2	0.0 (1)	1	0.0 (1)
Muscatine	185	30.8 (76)	219	20.1 (96)	54	25.9 (61)	185	12.4 (118)	214	0.9 (94)
Nevada	2	50.0 (127)	4	0.0 (1)	1	0.0 (100)	2	0.0 (1)	4	0.0 (1)
New Hampto	14	21.4 (45)	9	0.0 (1)	2	50.0 (11)	13	7.7 (89)	7	0.0 (1)
Newton	71	32.4 (82)	76	18.4 (83)	22	27.3 (58)	72	9.7 (105)	75	1.3 (100)
North Towne	104	23.1 (50)	118	11.9 (53)	29	48.3 (24)	113	1.8 (56)	134	2.2 (105)
Northwood	6	33.3 (84)	9	0.0 (1)	2	50.0 (11)	6	0.0 (1)	9	0.0 (1)
Norwalk	7	42.9 (116)	6	50.0 (141)	4	0.0 (100)	7	0.0 (1)	7	14.3 (140)
Oakridge	5	20.0 (39)	5	40.0 (135)	2	0.0 (100)	6	0.0 (1)	5	0.0 (1)
Oelwein	24	33.3 (84)	29	17.2 (77)	9	33.3 (40)	26	23.1 (144)	30	6.7 (127)
Onawa	17	17.6 (33)	19	26.3 (118)	5	0.0 (100)	17	5.9 (78)	18	0.0 (1)
Orange Cty	25	8.0 (14)	22	4.5 (26)	2	0.0 (100)	25	4.0 (62)	22	0.0 (1)
Osage	23	26.1 (62)	31	12.9 (57)	6	50.0 (11)	22	4.5 (65)	27	3.7 (117)
Osceola	37	18.9 (37)	39	20.5 (98)	6	16.7 (84)	37	5.4 (75)	36	0.0 (1)
Oskaloosa	27	40.7 (111)	25	32.0 (123)	9	11.1 (93)	27	14.8 (128)	24	8.3 (129)
Ottumwa	293	29.0 (71)	192	18.8 (85)	73	9.6 (96)	284	6.3 (83)	169	1.2 (98)
Pella	8	75.0 (144)	7	42.9 (137)	6	33.3 (40)	8	12.5 (119)	7	0.0 (1)
Perry	27	25.9 (60)	21	4.8 (27)	5	60.0 (10)	30	0.0 (1)	19	0.0 (1)
Pocahontas	1	100.0 (146)	3	33.3 (124)	1	0.0 (100)	1	0.0 (1)	3	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Postville	30	10.0 (18)	33	9.1 (37)	3	0.0 (100)	30	10.0 (108)	32	3.1 (112)
Red Oak	16	68.8 (143)	16	50.0 (141)	10	10.0 (94)	16	0.0 (1)	16	0.0 (1)
Redfield	3	33.3 (84)	2	50.0 (141)	1	0.0 (100)	2	0.0 (1)	2	0.0 (1)
Refugee Cln	8	0.0 (1)	11	0.0 (1)	0	0.0 (100)	8	0.0 (1)	8	0.0 (1)
Rock Rapid	6	16.7 (28)	12	16.7 (72)	1	0.0 (100)	6	0.0 (1)	12	0.0 (1)
Rock Valle	22	4.5 (12)	25	4.0 (25)	1	0.0 (100)	22	4.5 (65)	24	0.0 (1)
Rockwell C	17	29.4 (72)	20	20.0 (92)	6	16.7 (84)	17	17.6 (135)	20	5.0 (121)
SDM	164	29.9 (75)	169	18.3 (82)	62	33.9 (39)	176	9.1 (98)	199	3.5 (115)
Sac City	10	50.0 (127)	12	33.3 (124)	6	33.3 (40)	10	10.0 (108)	12	8.3 (129)
Sheldon	27	25.9 (60)	30	6.7 (28)	8	62.5 (9)	28	7.1 (86)	27	0.0 (1)
Shenandoah	28	53.6 (134)	26	19.2 (89)	13	15.4 (92)	28	7.1 (86)	24	0.0 (1)
Sibley	6	16.7 (28)	15	6.7 (28)	1	0.0 (100)	6	16.7 (131)	15	0.0 (1)
Sidney	5	40.0 (105)	4	25.0 (112)	1	0.0 (100)	5	20.0 (138)	4	0.0 (1)
Sigourney	6	66.7 (141)	6	50.0 (141)	4	0.0 (100)	6	0.0 (1)	6	0.0 (1)
Sioux Cent	22	9.1 (16)	26	7.7 (34)	3	33.3 (40)	22	0.0 (1)	25	0.0 (1)
Sioux City	477	21.6 (46)	377	11.7 (51)	87	23.0 (72)	486	6.0 (80)	413	1.0 (95)
Spencer	49	38.8 (103)	55	18.2 (80)	16	37.5 (35)	49	8.2 (95)	52	15.4 (141)
Spirit Lak	17	17.6 (33)	19	15.8 (71)	3	66.7 (5)	17	5.9 (78)	19	31.6 (145)
Storm Lake	64	12.5 (22)	107	1.9 (23)	8	75.0 (3)	64	4.7 (67)	103	3.9 (119)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Stuart	7	57.1 (136)	6	33.3 (124)	4	25.0 (64)	7	0.0 (1)	7	0.0 (1)
Tama	29	20.7 (43)	29	10.3 (43)	5	20.0 (76)	31	6.5 (84)	34	5.9 (122)
Tama/Settl	1	0.0 (1)	1	0.0 (1)	0	0.0 (100)	1	0.0 (1)	1	0.0 (1)
Tipton	3	33.3 (84)	7	42.9 (137)	2	0.0 (100)	3	0.0 (1)	7	0.0 (1)
Traer	2	50.0 (127)	2	0.0 (1)	1	100.0 (1)	2	0.0 (1)	2	0.0 (1)
UIHC	1	0.0 (1)	9	22.2 (102)	0	0.0 (100)	1	0.0 (1)	9	11.1 (136)
Vinton	15	20.0 (39)	13	7.7 (34)	4	25.0 (64)	17	17.6 (135)	12	0.0 (1)
WDM	40	17.5 (32)	43	7.0 (31)	5	20.0 (76)	41	0.0 (1)	44	2.3 (107)
Washington	28	28.6 (67)	18	22.2 (102)	5	0.0 (100)	28	10.7 (111)	17	0.0 (1)
Waterloo	372	31.5 (79)	331	20.5 (98)	110	16.4 (91)	380	8.9 (97)	340	1.8 (102)
Waukee	23	17.4 (31)	22	0.0 (1)	3	33.3 (40)	23	0.0 (1)	26	0.0 (1)
Waukon	18	22.2 (48)	20	10.0 (41)	4	50.0 (11)	18	5.6 (77)	19	15.8 (142)
Waverly	12	41.7 (113)	9	11.1 (48)	2	0.0 (100)	12	0.0 (1)	9	0.0 (1)
Wbstr Ct	50	34.0 (94)	65	20.0 (92)	21	33.3 (40)	50	8.0 (92)	65	0.0 (1)
West Union	15	40.0 (105)	15	13.3 (59)	6	33.3 (40)	15	33.3 (146)	15	33.3 (146)
Winterset	16	31.3 (77)	18	11.1 (48)	4	50.0 (11)	16	12.5 (119)	16	6.3 (124)
Iowa	**7,560**	**27.2**	**7,538**	**15.8**	**1,987**	**26.8**	**7,788**	**6.6**	**7,829**	**1.9**
National (Prior Year)	**108,764**	**27.1**	**119,789**	**14.3**	**24,778**	**36.6**	**97,290**	**10.0**	**120,070**	**3.5**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley	3	0.0 (1)	33.3 (145)	3	0.0 (1)	3	0.0 (1)	3	66.7 (90)
Adel	10	10.0 (109)	10.0 (98)	6	0.0 (1)	6	0.0 (1)	10	90.0 (9)
Albia	13	7.7 (78)	7.7 (67)	11	18.2 (132)	9	0.0 (1)	13	61.5 (116)
Algona	45	4.4 (51)	22.2 (140)	41	9.8 (74)	37	0.0 (1)	45	77.8 (38)
Allen WHC	94	6.4 (67)	5.3 (40)	91	7.7 (54)	83	2.4 (90)	95	66.3 (97)
Allison	23	21.7 (142)	8.7 (83)	19	21.1 (134)	15	13.3 (144)	23	73.9 (64)
Altoona	41	4.9 (55)	4.9 (38)	33	6.1 (41)	31	0.0 (1)	41	70.7 (72)
Ames	202	7.9 (83)	10.9 (102)	152	9.2 (70)	137	2.9 (95)	202	81.7 (26)
Anamosa	24	8.3 (87)	4.2 (34)	22	9.1 (67)	20	0.0 (1)	24	62.5 (108)
Ankeny	53	5.7 (62)	5.7 (45)	46	4.3 (36)	44	2.3 (89)	53	79.2 (32)
Atlantic	35	11.4 (119)	14.3 (120)	33	18.2 (132)	27	3.7 (105)	35	48.6 (143)
Audubon	9	11.1 (115)	0.0 (1)	9	0.0 (1)	9	11.1 (139)	9	77.8 (38)
Avoca	6	16.7 (136)	0.0 (1)	6	0.0 (1)	6	16.7 (147)	6	66.7 (90)
Bedford	8	12.5 (126)	25.0 (141)	7	0.0 (1)	7	0.0 (1)	8	62.5 (108)
Belle Plaine	8	0.0 (1)	25.0 (141)	8	0.0 (1)	8	0.0 (1)	8	62.5 (108)
Bloomfield	8	0.0 (1)	0.0 (1)	8	12.5 (99)	7	0.0 (1)	8	62.5 (108)
Boone	25	8.0 (85)	12.0 (113)	21	9.5 (72)	19	5.3 (122)	25	64.0 (102)
Brdlwns RP	1,094	9.6 (106)	6.9 (58)	1,003	9.0 (63)	903	4.2 (110)	1,108	79.1 (34)
Brooklyn	3	0.0 (1)	0.0 (1)	3	33.3 (141)	2	0.0 (1)	3	66.7 (90)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Burlington	166	6.6 (69)	6.0 (49)	142	7.0 (48)	132	5.3 (122)	166	54.2 (132)
CHC	249	12.0 (124)	9.6 (95)	214	10.7 (82)	191	6.8 (129)	249	67.9 (86)
CR Urban	433	7.2 (74)	8.8 (84)	394	7.9 (56)	345	3.2 (101)	455	74.5 (61)
Carlisle	13	0.0 (1)	7.7 (67)	7	0.0 (1)	7	0.0 (1)	13	69.2 (80)
Carroll	67	9.0 (93)	9.0 (88)	67	9.0 (63)	61	0.0 (1)	67	74.6 (60)
Cedar Falls	11	9.1 (95)	9.1 (89)	9	22.2 (136)	7	0.0 (1)	11	90.9 (8)
Centervill	29	3.4 (43)	6.9 (58)	24	12.5 (99)	21	0.0 (1)	29	58.6 (121)
Chariton	22	9.1 (95)	13.6 (119)	21	14.3 (110)	18	0.0 (1)	22	68.2 (85)
Charles City	63	3.2 (41)	11.1 (104)	54	11.1 (83)	47	2.1 (87)	64	64.1 (101)
Cherokee	37	13.5 (131)	8.1 (75)	35	11.4 (87)	31	12.9 (143)	37	89.2 (10)
Clarinda	25	0.0 (1)	8.0 (74)	23	8.7 (62)	21	0.0 (1)	24	75.0 (53)
Clarion	47	10.6 (113)	14.9 (128)	40	12.5 (99)	35	2.9 (95)	47	76.6 (44)
Clinton	175	10.9 (114)	7.4 (64)	132	9.8 (74)	119	5.0 (119)	176	69.3 (79)
Colfax	11	0.0 (1)	9.1 (89)	7	14.3 (110)	6	0.0 (1)	11	54.5 (130)
Columbus J	17	5.9 (64)	5.9 (47)	15	6.7 (44)	14	7.1 (131)	17	76.5 (45)
Corning	15	0.0 (1)	13.3 (117)	14	28.6 (139)	10	0.0 (1)	15	60.0 (118)
Corydon	16	18.8 (140)	18.8 (137)	15	13.3 (108)	13	7.7 (134)	16	68.8 (82)
Council Blfs	388	10.1 (112)	7.7 (67)	356	11.5 (89)	313	4.8 (118)	393	72.8 (66)
Cresco	27	7.4 (75)	14.8 (127)	25	12.0 (94)	22	0.0 (1)	27	85.2 (14)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Creston	75	4.0 (48)	5.3 (40)	65	6.2 (42)	61	1.6 (85)	75	82.7 (20)
Dallas Ctr	3	0.0 (1)	0.0 (1)	2	0.0 (1)	2	0.0 (1)	3	100.0 (1)
Decorah	22	18.2 (139)	9.1 (89)	17	11.8 (91)	15	6.7 (128)	22	81.8 (23)
Denison	101	6.9 (71)	4.0 (33)	101	14.9 (120)	86	1.2 (82)	101	64.4 (99)
Dewitt	6	16.7 (136)	0.0 (1)	5	40.0 (143)	3	0.0 (1)	6	66.7 (90)
Dubuque	352	7.4 (75)	6.5 (56)	303	7.3 (52)	281	5.0 (119)	355	51.8 (134)
Dyersville	7	0.0 (1)	14.3 (120)	7	0.0 (1)	7	0.0 (1)	7	57.1 (124)
Eagle Grove	28	3.6 (45)	7.1 (61)	27	11.1 (83)	24	0.0 (1)	28	75.0 (53)
Edgerton	364	9.3 (102)	9.6 (95)	339	12.7 (106)	295	2.4 (90)	365	69.0 (81)
Eldora	7	28.6 (143)	0.0 (1)	5	60.0 (148)	2	0.0 (1)	7	42.9 (145)
Elkader	26	7.7 (78)	11.5 (108)	24	4.2 (35)	23	8.7 (135)	26	80.8 (29)
Emmetsburg	12	16.7 (136)	0.0 (1)	8	12.5 (99)	7	0.0 (1)	12	75.0 (53)
Esthervill	37	5.4 (58)	10.8 (101)	31	12.9 (107)	26	0.0 (1)	38	63.2 (105)
Fairfiel	17	0.0 (1)	11.8 (109)	14	7.1 (50)	13	0.0 (1)	17	64.7 (98)
Forest City	55	12.7 (129)	10.9 (102)	54	14.8 (118)	46	2.2 (88)	55	70.9 (70)
Fort Dodge	214	7.9 (83)	8.4 (81)	199	10.1 (79)	179	1.1 (81)	214	66.8 (88)
Ft Madison	54	9.3 (102)	13.0 (116)	40	10.0 (76)	36	2.8 (94)	54	50.0 (136)
Garner	27	3.7 (46)	11.1 (104)	22	4.5 (37)	21	0.0 (1)	27	74.1 (63)
Glenwood	19	0.0 (1)	5.3 (40)	17	0.0 (1)	17	0.0 (1)	19	57.9 (122)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Grand Junction	2	50.0 (146)	0.0 (1)	2	0.0 (1)	2	50.0 (148)	2	50.0 (136)
Greenfield	17	0.0 (1)	17.6 (134)	17	0.0 (1)	17	0.0 (1)	17	70.6 (73)
Grinnell	26	3.8 (47)	7.7 (67)	21	4.8 (38)	20	0.0 (1)	26	57.7 (123)
Grundy Ctr	9	11.1 (115)	0.0 (1)	8	37.5 (142)	5	0.0 (1)	9	55.6 (127)
Guthrie Ct	14	0.0 (1)	0.0 (1)	13	0.0 (1)	13	0.0 (1)	14	64.3 (100)
Hampton	42	4.8 (54)	14.3 (120)	41	14.6 (113)	35	0.0 (1)	42	71.4 (67)
Harlan	38	5.3 (56)	0.0 (1)	35	14.3 (110)	30	0.0 (1)	38	55.3 (129)
Hawarden	17	11.8 (120)	11.8 (109)	17	5.9 (40)	16	6.3 (125)	17	82.4 (21)
Hispanic Cln	6	0.0 (1)	16.7 (132)	6	0.0 (1)	6	0.0 (1)	6	50.0 (136)
Humboldt	16	12.5 (126)	6.3 (52)	14	21.4 (135)	11	0.0 (1)	16	68.8 (82)
ILH	361	9.4 (104)	9.7 (97)	339	9.4 (71)	307	4.6 (117)	361	75.6 (52)
Ida Grove	22	4.5 (52)	4.5 (35)	22	9.1 (67)	20	0.0 (1)	22	81.8 (23)
Independen	32	6.3 (65)	15.6 (130)	26	7.7 (54)	24	0.0 (1)	32	62.5 (108)
Indianola	50	12.0 (124)	6.0 (49)	34	14.7 (116)	29	0.0 (1)	49	73.5 (65)
Iowa City	331	9.4 (104)	8.2 (78)	300	9.0 (63)	269	5.2 (121)	335	85.7 (13)
Iowa Falls	15	13.3 (130)	6.7 (57)	12	16.7 (123)	10	0.0 (1)	15	66.7 (90)
Jamaica	3	0.0 (1)	33.3 (145)	3	0.0 (1)	3	0.0 (1)	3	100.0 (1)
Jefferson	16	0.0 (1)	6.3 (52)	16	12.5 (99)	14	0.0 (1)	16	75.0 (53)
KR	13	7.7 (78)	15.4 (129)	13	15.4 (121)	11	0.0 (1)	13	84.6 (15)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Keokuk	51	11.8 (120)	3.9 (32)	47	10.6 (81)	41	7.3 (133)	51	45.1 (144)
Keosauqua	7	0.0 (1)	42.9 (148)	7	0.0 (1)	7	0.0 (1)	7	42.9 (145)
Knoxville	93	9.7 (108)	8.6 (82)	71	7.0 (48)	66	4.5 (115)	93	62.4 (114)
Lamoni	5	0.0 (1)	40.0 (147)	4	0.0 (1)	4	0.0 (1)	5	80.0 (31)
Le Mars	44	9.1 (95)	2.3 (26)	41	14.6 (113)	34	0.0 (1)	45	84.4 (16)
Lenox	13	0.0 (1)	7.7 (67)	12	0.0 (1)	12	0.0 (1)	13	53.8 (133)
Leon	22	9.1 (95)	0.0 (1)	18	0.0 (1)	18	5.6 (124)	22	54.5 (130)
Logan	33	9.1 (95)	6.1 (51)	32	12.5 (99)	28	3.6 (103)	33	78.8 (35)
Manchester	16	18.8 (140)	6.3 (52)	15	26.7 (138)	11	9.1 (136)	16	25.0 (148)
Manning	2	0.0 (1)	0.0 (1)	2	0.0 (1)	2	0.0 (1)	2	50.0 (136)
Maquoketa	30	10.0 (109)	3.3 (29)	23	26.1 (137)	17	0.0 (1)	30	56.7 (126)
Marengo	11	9.1 (95)	18.2 (136)	10	10.0 (76)	9	11.1 (139)	11	81.8 (23)
Marion	123	5.7 (62)	14.6 (125)	111	7.2 (51)	103	3.9 (106)	123	62.6 (107)
Marshalltown	285	8.8 (91)	8.1 (75)	240	9.6 (73)	211	4.3 (112)	292	76.4 (49)
Mason City	198	8.6 (89)	7.6 (66)	175	10.3 (80)	156	4.5 (115)	199	66.8 (88)
Monticello	9	0.0 (1)	0.0 (1)	9	11.1 (83)	8	0.0 (1)	9	77.8 (38)
Mt Ayr	12	0.0 (1)	0.0 (1)	11	0.0 (1)	11	0.0 (1)	12	66.7 (90)
Mt Pleasan	53	11.3 (118)	5.7 (45)	41	14.6 (113)	35	2.9 (95)	54	75.9 (50)
Mtown Dtown	2	0.0 (1)	0.0 (1)	2	0.0 (1)	2	0.0 (1)	2	50.0 (136)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Muscatine	219	8.2 (86)	7.8 (73)	179	12.3 (98)	157	2.5 (92)	219	76.7 (43)
Nevada	4	0.0 (1)	25.0 (141)	4	0.0 (1)	4	0.0 (1)	4	75.0 (53)
New Hampto	16	6.3 (65)	6.3 (52)	15	6.7 (44)	14	0.0 (1)	16	81.3 (27)
Newton	85	4.7 (53)	17.6 (134)	75	5.3 (39)	70	1.4 (84)	86	70.9 (70)
North Towne	141	9.2 (101)	7.1 (61)	122	9.0 (63)	110	6.4 (126)	143	69.9 (77)
Northwood	9	11.1 (115)	11.1 (104)	8	0.0 (1)	8	12.5 (142)	9	77.8 (38)
Norwalk	10	0.0 (1)	10.0 (98)	8	12.5 (99)	7	0.0 (1)	10	50.0 (136)
Oakridge	6	0.0 (1)	0.0 (1)	6	16.7 (123)	5	0.0 (1)	6	50.0 (136)
Oelwein	34	8.8 (91)	14.7 (126)	27	11.1 (83)	24	4.2 (110)	34	61.8 (115)
Onawa	27	7.4 (75)	3.7 (31)	25	8.0 (57)	23	4.3 (112)	27	70.4 (75)
Orange Cty	36	2.8 (38)	8.3 (79)	35	11.4 (87)	31	0.0 (1)	36	91.7 (7)
Osage	30	3.3 (42)	3.3 (29)	25	8.0 (57)	23	0.0 (1)	30	83.3 (17)
Osceola	51	7.8 (82)	11.8 (109)	45	15.6 (122)	38	2.6 (93)	51	70.6 (73)
Oskaloosa	43	7.0 (72)	9.3 (94)	35	8.6 (60)	32	3.1 (99)	43	51.2 (135)
Ottumwa	380	5.5 (59)	7.4 (64)	344	6.7 (44)	321	1.2 (82)	380	74.5 (61)
Pella	14	14.3 (133)	21.4 (139)	12	16.7 (123)	10	10.0 (138)	14	57.1 (124)
Perry	36	5.6 (60)	2.8 (27)	31	3.2 (33)	30	3.3 (102)	36	80.6 (30)
Pocahontas	3	33.3 (144)	0.0 (1)	2	50.0 (144)	1	0.0 (1)	3	66.7 (90)
Postville	37	13.5 (131)	5.4 (43)	34	14.7 (116)	29	6.9 (130)	37	67.6 (87)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Red Oak	29	3.4 (43)	6.9 (58)	27	7.4 (53)	25	0.0 (1)	29	75.9 (50)
Redfield	9	0.0 (1)	0.0 (1)	9	0.0 (1)	9	0.0 (1)	9	55.6 (127)
Refugee Cln	7	0.0 (1)	28.6 (144)	12	16.7 (123)	7	0.0 (1)	13	92.3 (5)
Rock Rapid	13	7.7 (78)	7.7 (67)	12	8.3 (59)	11	0.0 (1)	13	61.5 (116)
Rock Valle	34	2.9 (39)	8.8 (84)	33	12.1 (95)	28	3.6 (103)	35	82.9 (19)
Rockwell C	20	10.0 (109)	5.0 (39)	20	10.0 (76)	18	11.1 (139)	20	60.0 (118)
SDM	364	8.5 (88)	9.1 (89)	340	11.5 (89)	290	4.1 (109)	380	75.0 (53)
Sac City	10	0.0 (1)	0.0 (1)	10	0.0 (1)	10	0.0 (1)	10	60.0 (118)
Sheldon	35	8.6 (89)	2.9 (28)	30	16.7 (123)	25	0.0 (1)	35	88.6 (11)
Shenandoah	30	6.7 (70)	10.0 (98)	29	13.8 (109)	25	4.0 (107)	30	63.3 (104)
Sibley	16	0.0 (1)	12.5 (114)	12	0.0 (1)	12	0.0 (1)	16	81.3 (27)
Sidney	6	0.0 (1)	16.7 (132)	4	50.0 (144)	2	0.0 (1)	6	83.3 (17)
Sigourney	8	12.5 (126)	12.5 (114)	6	16.7 (123)	5	0.0 (1)	8	62.5 (108)
Sioux Cent	42	7.1 (73)	4.8 (37)	35	8.6 (60)	32	3.1 (99)	42	88.1 (12)
Sioux City	635	6.5 (68)	7.2 (63)	601	12.1 (95)	512	1.8 (86)	653	68.3 (84)
Spencer	52	1.9 (37)	13.5 (118)	47	6.4 (43)	44	0.0 (1)	52	78.8 (35)
Spirit Lak	19	5.3 (56)	15.8 (131)	18	16.7 (123)	15	0.0 (1)	19	63.2 (105)
Storm Lake	111	9.0 (93)	8.1 (75)	88	14.8 (118)	75	4.0 (107)	111	70.3 (76)
Stuart	11	36.4 (145)	9.1 (89)	10	30.0 (140)	7	14.3 (146)	11	36.4 (147)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Tama	36	5.6 (60)	11.1 (104)	33	0.0 (1)	33	3.0 (98)	36	69.4 (78)
Tama/Settl	2	50.0 (146)	0.0 (1)	2	50.0 (144)	1	0.0 (1)	2	100.0 (1)
Tipton	8	0.0 (1)	0.0 (1)	8	0.0 (1)	8	0.0 (1)	8	75.0 (53)
Traer	2	0.0 (1)	0.0 (1)	2	0.0 (1)	2	0.0 (1)	2	100.0 (1)
UIHC	13	76.9 (148)	0.0 (1)	2	50.0 (144)	1	0.0 (1)	13	92.3 (5)
Vinton	25	4.0 (48)	20.0 (138)	22	9.1 (67)	20	0.0 (1)	25	64.0 (102)
WDM	56	16.1 (135)	5.4 (43)	51	17.6 (130)	42	9.5 (137)	56	82.1 (22)
Washington	34	2.9 (39)	8.8 (84)	29	6.9 (47)	27	0.0 (1)	34	76.5 (45)
Waterloo	622	9.6 (106)	8.8 (84)	521	12.1 (95)	455	4.4 (114)	627	71.0 (69)
Waukee	35	0.0 (1)	14.3 (120)	29	3.4 (34)	28	0.0 (1)	35	77.1 (42)
Waukon	21	0.0 (1)	14.3 (120)	17	0.0 (1)	17	0.0 (1)	21	71.4 (67)
Waverly	17	11.8 (120)	11.8 (109)	17	11.8 (91)	15	13.3 (144)	17	76.5 (45)
Wbstr Ct	66	15.2 (134)	4.5 (35)	56	17.9 (131)	46	6.5 (127)	66	78.8 (35)
West Union	17	11.8 (120)	5.9 (47)	14	0.0 (1)	14	7.1 (131)	17	76.5 (45)
Winterset	24	4.2 (50)	8.3 (79)	17	11.8 (91)	15	0.0 (1)	24	79.2 (32)
Iowa	**10,590**	**8.3**	**8.4**	**9,401**	**10.2**	**8,360**	**3.4**	**10,700**	**71.7**
National (Prior Year)	**117,623**	**9.3**	**6.6**	**117,628**	**11.8**	**100,072**	**3.9**	**116,293**	**57.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Ackley	14	71.4	0.0	28.6	0.0	0.0	0.0	0.0
Adel	29	86.2	0.0	13.8	0.0	0.0	0.0	0.0
Albia	62	85.5	11.3	0.0	1.6	0.0	1.6	0.0
Algona	112	79.5	4.5	15.2	0.0	0.0	0.9	0.0
Allen WHC	277	50.2	32.1	7.2	0.4	7.6	2.5	0.0
Allison	60	95.0	0.0	3.3	0.0	0.0	1.7	0.0
Altoona	107	77.6	13.1	6.5	0.0	1.9	0.9	0.0
Ames	685	69.5	8.5	10.5	0.3	9.8	1.5	0.0
Anamosa	68	94.1	1.5	4.4	0.0	0.0	0.0	0.0
Ankeny	147	74.8	9.5	8.8	0.0	2.0	4.8	0.0
Atlantic	127	89.8	0.8	8.7	0.0	0.0	0.8	0.0
Audubon	36	94.4	0.0	5.6	0.0	0.0	0.0	0.0
Avoca	22	86.4	4.5	9.1	0.0	0.0	0.0	0.0
Bedford	26	92.3	0.0	0.0	0.0	3.8	3.8	0.0
Belle Plaine	23	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Bloomfield	16	81.3	6.3	6.3	0.0	0.0	6.3	0.0
Boone	78	84.6	2.6	10.3	0.0	0.0	2.6	0.0
Brdlwns RP	3,791	40.4	24.0	19.9	0.3	10.0	5.4	0.0
Brooklyn	17	70.6	0.0	29.4	0.0	0.0	0.0	0.0
Burlington	527	75.0	18.6	4.7	0.0	0.4	1.3	0.0
CHC	675	58.8	21.3	13.5	0.1	2.1	4.1	0.0
CR Urban	1,547	50.5	35.6	6.3	0.2	1.0	6.2	0.1
Carlisle	42	92.9	2.4	2.4	0.0	2.4	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)								
Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Carroll	214	76.2	6.5	16.4	0.0	0.9	0.0	0.0
Cedar Falls	34	76.5	2.9	11.8	0.0	8.8	0.0	0.0
Centervill	73	91.8	1.4	6.8	0.0	0.0	0.0	0.0
Chariton	54	96.3	0.0	3.7	0.0	0.0	0.0	0.0
Charles City	188	78.2	9.0	8.0	0.0	4.3	0.5	0.0
Cherokee	116	89.7	1.7	8.6	0.0	0.0	0.0	0.0
Clarinda	59	94.9	0.0	1.7	0.0	0.0	3.4	0.0
Clarion	134	38.1	0.0	61.9	0.0	0.0	0.0	0.0
Clinton	535	79.4	10.3	7.9	0.0	0.2	2.2	0.0
Colfax	34	88.2	0.0	8.8	0.0	2.9	0.0	0.0
Columbus J	42	11.9	0.0	42.9	2.4	42.9	0.0	0.0
Corning	32	96.9	0.0	3.1	0.0	0.0	0.0	0.0
Correctionvl	6	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Corydon	48	95.8	0.0	4.2	0.0	0.0	0.0	0.0
Council Blfs	1,457	78.1	3.2	15.9	0.7	0.5	1.6	0.0
Cresco	72	91.7	1.4	5.6	0.0	1.4	0.0	0.0
Creston	246	86.2	2.8	10.6	0.0	0.0	0.4	0.0
Dallas Ctr	7	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Decorah	77	83.1	1.3	13.0	0.0	0.0	2.6	0.0
Denison	347	32.9	3.7	57.1	0.3	6.1	0.0	0.0
Dewitt	30	90.0	0.0	6.7	0.0	0.0	3.3	0.0
Dubuque	1,157	72.9	13.6	7.0	0.4	4.1	2.0	0.0
Dyersville	13	84.6	7.7	7.7	0.0	0.0	0.0	0.0
Eagle Grove	70	34.3	1.4	64.3	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Edgerton	1,279	58.6	24.2	9.0	0.7	1.8	5.6	0.0
Eldora	25	84.0	0.0	12.0	0.0	0.0	4.0	0.0
Elkader	103	91.3	0.0	6.8	0.0	1.0	1.0	0.0
Emmetsburg	67	89.6	3.0	6.0	0.0	0.0	1.5	0.0
Esthervill	136	61.0	2.9	33.8	0.0	0.7	1.5	0.0
Fairfiel	64	85.9	3.1	4.7	0.0	6.3	0.0	0.0
Fayette	20	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Forest City	134	83.6	4.5	11.2	0.7	0.0	0.0	0.0
Fort Dodge	615	77.4	8.3	11.4	0.5	0.2	2.3	0.0
Ft Madison	186	90.9	2.7	5.4	0.0	0.0	1.1	0.0
Garner	74	77.0	1.4	18.9	1.4	0.0	1.4	0.0
Glenwood	69	87.0	0.0	13.0	0.0	0.0	0.0	0.0
Grand Junction	6	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Greenfield	49	87.8	0.0	10.2	0.0	2.0	0.0	0.0
Grinnell	108	88.0	1.9	9.3	0.0	0.0	0.9	0.0
Grundy Ctr	26	96.2	0.0	3.8	0.0	0.0	0.0	0.0
Guthrie Ct	62	88.7	1.6	8.1	0.0	0.0	1.6	0.0
Hampton	131	53.4	0.8	45.8	0.0	0.0	0.0	0.0
Harlan	96	86.5	1.0	12.5	0.0	0.0	0.0	0.0
Hawarden	71	57.7	0.0	40.8	1.4	0.0	0.0	0.0
Hispanic Cln	20	10.0	0.0	90.0	0.0	0.0	0.0	0.0
Humboldt	73	65.8	2.7	30.1	0.0	0.0	1.4	0.0
ILH	1,450	39.6	15.7	35.4	0.3	6.1	3.0	0.0
Ida Grove	82	84.1	1.2	13.4	0.0	0.0	1.2	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Independen	103	96.1	1.0	2.9	0.0	0.0	0.0	0.0
Indianola	177	91.5	2.3	4.0	0.0	0.6	1.7	0.0
Iowa City	1,184	36.1	42.1	16.9	0.3	2.3	2.3	0.0
Iowa Falls	44	77.3	2.3	11.4	0.0	0.0	9.1	0.0
Jamaica	10	80.0	0.0	20.0	0.0	0.0	0.0	0.0
Jefferson	70	80.0	0.0	17.1	0.0	1.4	1.4	0.0
KR	46	69.6	13.0	4.3	0.0	10.9	2.2	0.0
Keokuk	178	88.2	4.5	5.1	0.0	0.6	1.7	0.0
Keosauqua	15	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Knoxville	235	91.5	2.6	3.8	0.4	0.4	1.3	0.0
Lamoni	20	70.0	5.0	20.0	0.0	0.0	5.0	0.0
Le Mars	137	66.4	3.6	24.1	1.5	2.9	1.5	0.0
Lenox	28	67.9	0.0	28.6	0.0	3.6	0.0	0.0
Leon	48	97.9	2.1	0.0	0.0	0.0	0.0	0.0
Logan	102	93.1	0.0	5.9	1.0	0.0	0.0	0.0
Manchester	61	90.2	3.3	6.6	0.0	0.0	0.0	0.0
Manning	2	50.0	0.0	0.0	0.0	50.0	0.0	0.0
Maquoketa	92	82.6	4.3	2.2	0.0	6.5	4.3	0.0
Marengo	32	87.5	0.0	9.4	0.0	0.0	3.1	0.0
Marion	422	82.0	6.4	8.5	0.0	0.9	2.1	0.0
Marshalltown	894	39.8	2.5	47.0	0.3	9.1	1.2	0.0
Mason City	686	77.0	6.0	10.5	0.4	3.1	3.1	0.0
Monticello	23	91.3	0.0	8.7	0.0	0.0	0.0	0.0
Mt Ayr	31	93.5	0.0	6.5	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Mt Pleasan	175	78.3	4.6	16.0	0.0	0.6	0.6	0.0
Mtown Dtown	2	0.0	0.0	50.0	0.0	50.0	0.0	0.0
Murray	1	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Muscatine	774	59.3	6.2	32.2	0.0	2.1	0.3	0.0
Nevada	12	58.3	0.0	41.7	0.0	0.0	0.0	0.0
New Hampto	58	69.0	0.0	27.6	0.0	0.0	3.4	0.0
Newton	292	82.2	4.1	10.3	0.7	1.7	1.0	0.0
North Towne	549	63.6	19.5	7.5	0.0	5.5	4.0	0.0
Northwood	24	79.2	4.2	12.5	0.0	4.2	0.0	0.0
Norwalk	43	86.0	0.0	11.6	0.0	2.3	0.0	0.0
Oakridge	8	25.0	50.0	12.5	0.0	0.0	12.5	0.0
Oelwein	97	87.6	3.1	8.2	0.0	0.0	1.0	0.0
Onawa	81	95.1	0.0	1.2	2.5	0.0	1.2	0.0
Orange Cty	82	45.1	1.2	51.2	0.0	2.4	0.0	0.0
Osage	79	89.9	1.3	6.3	0.0	1.3	1.3	0.0
Osceola	128	67.2	0.8	31.3	0.0	0.8	0.0	0.0
Oskaloosa	141	91.5	4.3	4.3	0.0	0.0	0.0	0.0
Ottumwa	1,221	72.3	7.4	15.0	0.1	3.9	1.2	0.1
Pella	45	91.1	0.0	4.4	0.0	4.4	0.0	0.0
Perry	149	39.6	5.4	49.7	0.0	2.0	3.4	0.0
Pocahontas	26	84.6	3.8	0.0	0.0	11.5	0.0	0.0
Pomeroy	1	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Postville	120	35.8	10.0	53.3	0.0	0.0	0.8	0.0
Red Oak	130	93.8	0.0	5.4	0.0	0.8	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Redfield	10	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Refugee Cln	16	6.3	93.8	0.0	0.0	0.0	0.0	0.0
Rock Rapid	54	83.3	1.9	9.3	0.0	5.6	0.0	0.0
Rock Valle	99	40.4	0.0	56.6	2.0	0.0	1.0	0.0
Rockwell C	52	94.2	0.0	5.8	0.0	0.0	0.0	0.0
SDM	762	46.1	13.6	27.6	0.4	8.1	4.2	0.0
Sac City	39	97.4	0.0	2.6	0.0	0.0	0.0	0.0
Sheldon	128	62.5	3.1	31.3	0.8	2.3	0.0	0.0
Shenandoah	73	87.7	0.0	12.3	0.0	0.0	0.0	0.0
Sibley	44	61.4	4.5	34.1	0.0	0.0	0.0	0.0
Sidney	30	96.7	0.0	3.3	0.0	0.0	0.0	0.0
Sigourney	33	97.0	0.0	3.0	0.0	0.0	0.0	0.0
Sioux Cent	151	27.2	0.0	70.9	0.0	0.0	2.0	0.0
Sioux City	1,934	43.0	10.9	36.7	3.5	3.5	2.5	0.0
Southridge	304	49.7	13.5	26.0	0.3	6.6	3.9	0.0
Spencer	197	79.2	0.5	17.3	0.0	2.5	0.5	0.0
Spirit Lak	60	81.7	1.7	13.3	0.0	1.7	1.7	0.0
Storm Lake	400	27.8	4.5	50.0	0.3	17.0	0.5	0.0
Stuart	20	90.0	5.0	0.0	0.0	0.0	5.0	0.0
Tama	156	41.7	0.6	43.6	10.9	0.6	2.6	0.0
Tama/Settl	11	27.3	0.0	9.1	63.6	0.0	0.0	0.0
Tipton	25	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Traer	14	100.0	0.0	0.0	0.0	0.0	0.0	0.0
UIHC	35	71.4	14.3	8.6	0.0	5.7	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Comparison of Racial and Ethnic Distribution (2)

Clinic	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Vinton	75	97.3	0.0	2.7	0.0	0.0	0.0	0.0
WDM	213	51.6	17.4	24.9	0.5	2.3	3.3	0.0
Wapello	1	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Washington	106	68.9	0.9	27.4	0.0	0.0	2.8	0.0
Waterloo	2,033	57.4	23.7	9.6	0.6	5.0	3.7	0.0
Waukee	93	75.3	4.3	10.8	0.0	4.3	5.4	0.0
Waukon	76	90.8	0.0	7.9	0.0	1.3	0.0	0.0
Waverly	65	93.8	3.1	3.1	0.0	0.0	0.0	0.0
Wbstr Ct	188	62.8	1.6	29.8	0.0	5.9	0.0	0.0
West Union	33	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Williamsburg	1	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Winterset	62	91.9	1.6	6.5	0.0	0.0	0.0	0.0
Iowa	**34,950**	**60.9**	**13.4**	**18.6**	**0.5**	**3.9**	**2.6**	**0.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Ackley	14	0.0	0.0	0.0	64.3	35.7	0.0	0.0
Adel	29	0.0	3.4	6.9	65.5	24.1	0.0	0.0
Albia	62	0.0	1.6	3.2	66.1	29.0	0.0	0.0
Algona	112	0.0	0.9	7.1	56.3	35.7	0.0	0.0
Allen WHC	277	0.4	3.2	5.8	57.4	31.8	1.4	0.0
Allison	60	0.0	0.0	1.7	56.7	41.7	0.0	0.0
Altoona	107	0.0	3.7	3.7	53.3	38.3	0.9	0.0
Ames	685	0.0	1.6	5.5	57.5	33.3	2.0	0.0
Anamosa	68	0.0	2.9	2.9	64.7	27.9	1.5	0.0
Ankeny	147	0.0	2.0	4.1	66.7	23.1	4.1	0.0
Atlantic	127	0.0	1.6	7.9	61.4	27.6	0.8	0.8
Audubon	36	0.0	5.6	11.1	63.9	19.4	0.0	0.0
Avoca	22	0.0	0.0	4.5	63.6	31.8	0.0	0.0
Bedford	26	0.0	0.0	7.7	69.2	23.1	0.0	0.0
Belle Plaine	23	0.0	0.0	0.0	69.6	30.4	0.0	0.0
Bloomfield	16	0.0	0.0	6.3	75.0	12.5	6.3	0.0
Boone	78	0.0	1.3	3.8	62.8	28.2	3.8	0.0
Brdlwns RP	3,791	0.1	2.2	6.1	57.4	31.1	3.1	0.0
Brooklyn	17	0.0	0.0	0.0	82.4	17.6	0.0	0.0
Burlington	527	0.0	3.6	7.6	61.3	26.8	0.8	0.0
CHC	675	0.1	2.2	7.1	64.1	24.9	1.5	0.0
CR Urban	1,547	0.0	1.2	6.2	61.2	29.3	2.1	0.0
Carlisle	42	0.0	0.0	9.5	61.9	26.2	2.4	0.0
Carroll	214	0.0	0.5	4.7	64.0	29.0	1.9	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cedar Falls	34	0.0	0.0	0.0	55.9	44.1	0.0	0.0
Centervill	73	0.0	1.4	6.8	64.4	26.0	1.4	0.0
Chariton	54	0.0	0.0	9.3	66.7	22.2	1.9	0.0
Charles City	188	0.0	0.5	4.3	68.6	26.6	0.0	0.0
Cherokee	116	0.0	1.7	12.1	68.1	17.2	0.9	0.0
Clarinda	59	0.0	0.0	6.8	72.9	16.9	3.4	0.0
Clarion	134	0.7	4.5	6.7	56.0	30.6	1.5	0.0
Clinton	535	0.0	2.8	6.7	65.4	23.2	1.7	0.2
Colfax	34	0.0	2.9	11.8	58.8	20.6	5.9	0.0
Columbus J	42	0.0	0.0	0.0	40.5	52.4	7.1	0.0
Corning	32	0.0	0.0	9.4	62.5	28.1	0.0	0.0
Correctionvl	6	0.0	0.0	0.0	83.3	16.7	0.0	0.0
Corydon	48	0.0	2.1	2.1	66.7	29.2	0.0	0.0
Council Blfs	1,457	0.0	2.7	7.2	64.9	23.7	1.6	0.0
Cresco	72	0.0	4.2	5.6	56.9	31.9	1.4	0.0
Creston	246	0.4	3.3	10.6	61.0	23.2	1.2	0.4
Dallas Ctr	7	0.0	0.0	0.0	71.4	28.6	0.0	0.0
Decorah	77	0.0	1.3	7.8	57.1	32.5	1.3	0.0
Denison	347	0.0	1.4	2.6	62.5	30.3	3.2	0.0
Dewitt	30	0.0	3.3	10.0	46.7	40.0	0.0	0.0
Dubuque	1,157	0.0	2.7	6.7	62.4	26.8	1.3	0.2
Dyersville	13	0.0	0.0	0.0	61.5	38.5	0.0	0.0
Eagle Grove	70	0.0	1.4	8.6	58.6	27.1	4.3	0.0
Edgerton	1,279	0.2	3.6	8.1	61.8	24.7	1.6	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Eldora	25	0.0	0.0	8.0	68.0	24.0	0.0	0.0
Elkader	103	0.0	1.0	1.0	67.0	31.1	0.0	0.0
Emmetsburg	67	0.0	0.0	3.0	65.7	29.9	1.5	0.0
Esthervill	136	0.0	1.5	6.6	70.6	20.6	0.7	0.0
Fairfiel	64	0.0	0.0	1.6	71.9	23.4	3.1	0.0
Fayette	20	0.0	0.0	0.0	70.0	25.0	5.0	0.0
Forest City	134	0.0	0.7	3.7	68.7	26.1	0.7	0.0
Fort Dodge	615	0.2	3.1	6.7	63.3	26.0	0.8	0.0
Ft Madison	186	0.0	4.3	5.4	65.6	23.7	1.1	0.0
Garner	74	0.0	0.0	9.5	63.5	24.3	2.7	0.0
Glenwood	69	0.0	1.4	7.2	68.1	23.2	0.0	0.0
Grand Junction	6	0.0	0.0	50.0	33.3	16.7	0.0	0.0
Greenfield	49	0.0	0.0	4.1	61.2	34.7	0.0	0.0
Grinnell	108	0.0	2.8	5.6	63.9	26.9	0.9	0.0
Grundy Ctr	26	0.0	0.0	3.8	69.2	26.9	0.0	0.0
Guthrie Ct	62	0.0	1.6	6.5	66.1	25.8	0.0	0.0
Hampton	131	0.8	3.1	5.3	52.7	34.4	3.8	0.0
Harlan	96	0.0	3.1	3.1	69.8	20.8	3.1	0.0
Hawarden	71	0.0	1.4	5.6	60.6	29.6	2.8	0.0
Hispanic Cln	20	0.0	0.0	10.0	45.0	40.0	5.0	0.0
Humboldt	73	0.0	0.0	5.5	53.4	41.1	0.0	0.0
ILH	1,450	0.1	3.1	7.0	56.2	30.8	2.8	0.0
Ida Grove	82	0.0	1.2	7.3	57.3	31.7	2.4	0.0
Independen	103	0.0	1.9	1.0	64.1	29.1	3.9	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Indianola	177	0.0	1.7	7.3	65.0	24.3	1.7	0.0
Iowa City	1,184	0.0	0.6	3.7	52.1	40.2	3.4	0.0
Iowa Falls	44	0.0	2.3	4.5	70.5	22.7	0.0	0.0
Jamaica	10	0.0	10.0	0.0	60.0	30.0	0.0	0.0
Jefferson	70	0.0	0.0	5.7	78.6	14.3	1.4	0.0
KR	46	0.0	0.0	0.0	52.2	43.5	4.3	0.0
Keokuk	178	0.0	2.8	11.2	67.4	18.5	0.0	0.0
Keosauqua	15	0.0	13.3	0.0	33.3	53.3	0.0	0.0
Knoxville	235	0.0	2.1	7.7	63.8	23.4	3.0	0.0
Lamoni	20	0.0	0.0	10.0	85.0	5.0	0.0	0.0
Le Mars	137	0.0	1.5	8.8	61.3	24.8	3.6	0.0
Lenox	28	0.0	0.0	3.6	64.3	32.1	0.0	0.0
Leon	48	0.0	0.0	10.4	68.8	18.8	2.1	0.0
Logan	102	0.0	1.0	4.9	60.8	33.3	0.0	0.0
Manchester	61	0.0	0.0	4.9	65.6	27.9	1.6	0.0
Manning	2	0.0	0.0	50.0	50.0	0.0	0.0	0.0
Maquoketa	92	0.0	1.1	6.5	70.7	21.7	0.0	0.0
Marengo	32	0.0	3.1	3.1	75.0	18.8	0.0	0.0
Marion	422	0.0	1.9	5.0	63.0	28.4	1.7	0.0
Marshalltown	894	0.1	3.1	6.6	59.5	28.4	2.2	0.0
Mason City	686	0.0	1.5	7.6	65.6	23.6	1.7	0.0
Monticello	23	0.0	0.0	4.3	52.2	43.5	0.0	0.0
Mt Ayr	31	0.0	0.0	9.7	71.0	19.4	0.0	0.0
Mt Pleasan	175	0.0	2.9	6.3	65.7	22.9	2.3	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)								
Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Mtown Dtown	2	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Murray	1	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Muscatine	774	0.1	3.0	7.9	60.6	26.2	2.2	0.0
Nevada	12	0.0	0.0	0.0	66.7	33.3	0.0	0.0
New Hampto	58	0.0	0.0	8.6	60.3	31.0	0.0	0.0
Newton	292	0.0	1.4	7.9	61.6	27.7	1.0	0.3
North Towne	549	0.0	2.7	7.7	60.5	27.5	1.6	0.0
Northwood	24	0.0	0.0	4.2	66.7	25.0	4.2	0.0
Norwalk	43	0.0	2.3	2.3	53.5	39.5	2.3	0.0
Oakridge	8	0.0	0.0	0.0	37.5	62.5	0.0	0.0
Oelwein	97	0.0	2.1	8.2	60.8	28.9	0.0	0.0
Onawa	81	0.0	2.5	3.7	58.0	35.8	0.0	0.0
Orange Cty	82	0.0	2.4	6.1	61.0	26.8	3.7	0.0
Osage	79	0.0	0.0	3.8	60.8	35.4	0.0	0.0
Osceola	128	0.0	0.8	4.7	63.3	28.9	2.3	0.0
Oskaloosa	141	0.0	2.1	6.4	61.0	26.2	4.3	0.0
Ottumwa	1,221	0.0	2.3	7.1	63.7	25.1	1.6	0.1
Pella	45	0.0	4.4	4.4	62.2	28.9	0.0	0.0
Perry	149	0.7	2.0	5.4	61.1	28.2	2.7	0.0
Pocahontas	26	0.0	7.7	0.0	69.2	23.1	0.0	0.0
Pomeroy	1	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Postville	120	0.0	1.7	4.2	46.7	46.7	0.8	0.0
Red Oak	130	0.0	2.3	10.8	58.5	27.7	0.8	0.0
Redfield	10	0.0	0.0	0.0	70.0	30.0	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Refugee Cln	16	0.0	6.3	0.0	37.5	50.0	6.3	0.0
Rock Rapid	54	0.0	3.7	5.6	59.3	27.8	3.7	0.0
Rock Valle	99	0.0	3.0	9.1	59.6	27.3	1.0	0.0
Rockwell C	52	0.0	0.0	3.8	69.2	23.1	3.8	0.0
SDM	762	0.0	2.4	7.0	60.8	27.6	2.4	0.0
Sac City	39	0.0	2.6	2.6	48.7	43.6	2.6	0.0
Sheldon	128	0.0	3.9	5.5	57.8	31.3	1.6	0.0
Shenandoah	73	0.0	8.2	6.8	65.8	17.8	1.4	0.0
Sibley	44	0.0	0.0	2.3	59.1	36.4	2.3	0.0
Sidney	30	0.0	0.0	13.3	53.3	33.3	0.0	0.0
Sigourney	33	0.0	0.0	6.1	69.7	24.2	0.0	0.0
Sioux Cent	151	0.7	6.0	12.6	47.7	30.5	2.0	0.7
Sioux City	1,934	0.1	2.0	6.2	59.9	29.7	2.1	0.0
Southridge	304	0.3	3.0	5.6	59.5	28.9	2.6	0.0
Spencer	197	0.0	1.5	6.6	71.6	19.8	0.5	0.0
Spirit Lak	60	0.0	3.3	8.3	68.3	18.3	1.7	0.0
Storm Lake	400	0.0	2.0	5.3	57.5	31.5	3.8	0.0
Stuart	20	0.0	0.0	10.0	45.0	45.0	0.0	0.0
Tama	156	0.0	1.9	4.5	63.5	29.5	0.6	0.0
Tama/Settl	11	0.0	0.0	9.1	45.5	36.4	9.1	0.0
Tipton	25	0.0	4.0	16.0	64.0	12.0	4.0	0.0
Traer	14	0.0	0.0	7.1	85.7	7.1	0.0	0.0
UIHC	35	0.0	2.9	2.9	57.1	37.1	0.0	0.0
Vinton	75	0.0	2.7	4.0	65.3	25.3	2.7	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Comparison of Age Distribution (2)

Clinic	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
WDM	213	0.0	0.0	5.6	59.2	30.5	4.7	0.0
Wapello	1	0.0	0.0	0.0	0.0	100.0	0.0	0.0
Washington	106	0.0	0.9	2.8	60.4	35.8	0.0	0.0
Waterloo	2,033	0.1	1.7	5.2	63.8	27.8	1.3	0.0
Waukee	93	0.0	3.2	4.3	53.8	36.6	2.2	0.0
Waukon	76	0.0	2.6	1.3	75.0	21.1	0.0	0.0
Waverly	65	0.0	1.5	3.1	56.9	36.9	1.5	0.0
Wbstr Ct	188	0.0	4.3	6.9	59.6	26.1	3.2	0.0
West Union	33	0.0	3.0	9.1	75.8	9.1	3.0	0.0
Williamsburg	1	0.0	0.0	0.0	0.0	100.0	0.0	0.0
Winterset	62	0.0	0.0	4.8	59.7	33.9	1.6	0.0
Iowa	**34,950**	**0.1**	**2.2**	**6.3**	**61.0**	**28.4**	**2.0**	**0.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ackley	12	0.0 (1)	50.0 (12)	12	16.7 (28)	41.7 (55)	5	20.0 (101)	6	50.0 (155)	0	0.0 (1)	0	0.0 (1)
Adel	8	0.0 (1)	62.5 (81)	8	12.5 (22)	87.5 (159)	4	75.0 (162)	3	0.0 (1)	1	100.0 (34)	0	0.0 (1)
Adel	9	0.0 (1)	66.7 (113)	8	25.0 (108)	37.5 (29)	3	0.0 (1)	2	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Albia	49	4.1 (111)	53.1 (21)	48	27.1 (126)	41.7 (55)	12	0.0 (1)	12	8.3 (86)	3	100.0 (34)	1	100.0 (43)
Algona	85	5.9 (135)	65.9 (109)	82	26.8 (122)	48.8 (104)	15	0.0 (1)	28	17.9 (107)	6	100.0 (34)	3	100.0 (43)
Allen WHC	257	3.1 (81)	55.3 (28)	238	19.7 (48)	46.6 (81)	31	3.2 (69)	31	6.5 (80)	6	100.0 (34)	5	100.0 (43)
Allison	46	2.2 (63)	73.9 (144)	46	17.4 (34)	50.0 (111)	7	0.0 (1)	20	0.0 (1)	6	100.0 (34)	1	100.0 (43)
Altoona	73	4.1 (111)	63.0 (87)	71	32.4 (145)	36.6 (28)	19	0.0 (1)	11	0.0 (1)	12	100.0 (34)	3	100.0 (43)
Ames	506	5.3 (128)	56.5 (34)	487	24.8 (107)	40.9 (49)	120	9.2 (89)	148	12.2 (99)	61	100.0 (34)	6	100.0 (43)
Anamosa	9	0.0 (1)	77.8 (152)	9	11.1 (17)	66.7 (153)	1	0.0 (1)	1	0.0 (1)	0	0.0 (1)	1	100.0 (43)
Anamosa	46	0.0 (1)	54.3 (24)	45	17.8 (36)	55.6 (142)	13	38.5 (149)	7	28.6 (128)	2	100.0 (34)	3	100.0 (43)
Ankeny	101	7.9 (154)	60.4 (62)	95	20.0 (51)	45.3 (73)	41	0.0 (1)	21	0.0 (1)	12	100.0 (34)	3	100.0 (43)
Atlantic	105	6.7 (143)	61.9 (78)	102	25.5 (115)	56.9 (147)	29	6.9 (82)	34	2.9 (59)	0	0.0 (1)	3	100.0 (43)
Audubon	33	0.0 (1)	57.6 (40)	32	21.9 (75)	53.1 (135)	12	33.3 (125)	17	17.6 (104)	2	100.0 (34)	0	0.0 (1)
Avoca	19	0.0 (1)	57.9 (41)	17	23.5 (91)	47.1 (83)	4	0.0 (1)	2	0.0 (1)	1	100.0 (34)	2	100.0 (43)
Bedford	19	10.5 (160)	68.4 (130)	19	42.1 (157)	21.1 (3)	3	33.3 (125)	5	20.0 (109)	2	100.0 (34)	0	0.0 (1)
Belle Plaine	21	4.8 (123)	81.0 (158)	20	25.0 (108)	40.0 (39)	4	0.0 (1)	5	0.0 (1)	4	100.0 (34)	2	100.0 (43)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Bloomfield	15	6.7 (143)	60.0 (58)	14	35.7 (153)	50.0 (111)	5	40.0 (151)	7	28.6 (128)	0	0.0 (1)	0	0.0 (1)
Boone	53	1.9 (56)	66.0 (110)	52	38.5 (155)	44.2 (67)	13	30.8 (120)	13	7.7 (82)	3	100.0 (34)	1	100.0 (43)
Brdlwns RP	2,986	3.4 (89)	54.5 (25)	2,852	24.2 (102)	40.6 (46)	828	2.9 (67)	249	6.0 (77)	217	100.0 (34)	98	100.0 (43)
Brooklyn	15	0.0 (1)	66.7 (113)	15	20.0 (51)	60.0 (150)	2	50.0 (156)	9	33.3 (135)	0	0.0 (1)	0	0.0 (1)
Burlington	371	6.2 (139)	59.3 (52)	358	16.8 (31)	52.8 (129)	127	6.3 (80)	51	5.9 (76)	43	100.0 (34)	34	100.0 (43)
CHC	432	3.7 (101)	66.4 (112)	402	21.6 (72)	48.3 (95)	124	37.1 (140)	64	28.1 (126)	5	100.0 (34)	26	100.0 (43)
CR Urban	1,123	3.8 (105)	58.8 (49)	1,067	22.9 (86)	48.1 (94)	328	37.5 (142)	196	34.2 (143)	116	100.0 (34)	49	100.0 (43)
Carlisle	23	4.3 (115)	56.5 (34)	23	13.0 (23)	47.8 (91)	7	0.0 (1)	5	0.0 (1)	1	100.0 (34)	1	100.0 (43)
Carroll	174	2.3 (65)	65.5 (105)	168	16.7 (28)	53.0 (134)	60	31.7 (121)	32	18.8 (108)	19	100.0 (34)	16	100.0 (43)
Cedar Falls	20	5.0 (125)	60.0 (58)	20	35.0 (152)	45.0 (71)	4	0.0 (1)	5	0.0 (1)	2	100.0 (34)	2	100.0 (43)
Centervill	57	3.5 (95)	73.7 (143)	55	29.1 (135)	49.1 (109)	16	37.5 (142)	17	35.3 (145)	1	100.0 (34)	3	100.0 (43)
Chariton	8	0.0 (1)	37.5 (8)	8	25.0 (108)	37.5 (29)	3	0.0 (1)	4	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Chariton	25	8.0 (155)	48.0 (11)	24	45.8 (160)	33.3 (18)	8	37.5 (142)	3	0.0 (1)	3	100.0 (34)	1	100.0 (43)
Charles City	132	2.3 (65)	62.9 (86)	129	20.2 (58)	41.1 (52)	28	7.1 (83)	33	3.0 (60)	18	100.0 (34)	5	100.0 (43)
Cherokee	91	4.4 (118)	60.4 (62)	88	20.5 (60)	48.9 (106)	19	0.0 (1)	37	0.0 (1)	4	100.0 (34)	2	100.0 (43)
Clarinda	45	4.4 (118)	55.6 (29)	40	20.0 (51)	55.0 (141)	8	62.5 (159)	12	25.0 (118)	0	0.0 (1)	0	0.0 (1)
Clarion	111	2.7 (70)	58.6 (46)	103	31.1 (141)	35.0 (21)	28	3.6 (71)	14	0.0 (1)	9	100.0 (34)	2	100.0 (43)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Clinton	388	4.1 (111)	63.1 (90)	366	21.0 (66)	54.4 (139)	111	32.4 (123)	43	34.9 (144)	24	100.0 (34)	23	100.0 (43)
Colfax	22	9.1 (158)	72.7 (140)	22	18.2 (39)	36.4 (27)	9	11.1 (92)	5	60.0 (161)	1	100.0 (34)	2	100.0 (43)
Columbus J	35	2.9 (77)	22.9 (6)	35	34.3 (150)	25.7 (9)	10	0.0 (1)	8	0.0 (1)	7	100.0 (34)	0	0.0 (1)
Corning	26	3.8 (105)	65.4 (104)	26	26.9 (123)	50.0 (111)	9	55.6 (158)	4	25.0 (118)	2	100.0 (34)	1	100.0 (43)
Correctionvl	5	0.0 (1)	80.0 (157)	5	20.0 (51)	60.0 (150)	3	33.3 (125)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Corydon	35	2.9 (77)	77.1 (149)	34	32.4 (145)	50.0 (111)	10	0.0 (1)	11	9.1 (87)	0	0.0 (1)	1	100.0 (43)
Council Blfs	1,112	3.4 (89)	62.1 (80)	1,070	22.2 (80)	49.0 (107)	353	7.1 (83)	169	4.7 (70)	49	100.0 (34)	49	100.0 (43)
Cresco	58	8.6 (157)	50.0 (12)	52	17.3 (32)	46.2 (79)	24	12.5 (96)	22	31.8 (134)	7	100.0 (34)	3	100.0 (43)
Creston	172	5.2 (127)	59.3 (52)	163	19.0 (46)	51.5 (121)	41	26.8 (113)	29	20.7 (113)	16	100.0 (34)	7	100.0 (43)
Dallas Ctr	3	0.0 (1)	0.0 (1)	3	0.0 (1)	66.7 (153)	1	0.0 (1)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Decorah	59	3.4 (89)	55.9 (30)	59	10.2 (16)	54.2 (138)	18	33.3 (125)	15	26.7 (125)	6	100.0 (34)	2	100.0 (43)
Denison	295	2.0 (60)	61.4 (72)	291	32.0 (143)	34.4 (20)	116	1.7 (63)	62	3.2 (62)	1	100.0 (34)	13	100.0 (43)
Dewitt	22	4.5 (121)	72.7 (140)	20	30.0 (139)	40.0 (39)	8	37.5 (142)	6	0.0 (1)	2	100.0 (34)	1	100.0 (43)
Dubuque	921	3.9 (110)	59.8 (56)	873	21.4 (69)	48.7 (100)	253	49.8 (155)	86	38.4 (149)	64	100.0 (34)	14	100.0 (43)
Dyersville	9	0.0 (1)	77.8 (152)	9	11.1 (17)	22.2 (5)	3	33.3 (125)	3	0.0 (1)	1	100.0 (34)	0	0.0 (1)
Eagle Grove	63	3.2 (83)	52.4 (20)	62	32.3 (144)	27.4 (12)	20	5.0 (77)	6	33.3 (135)	4	100.0 (34)	3	100.0 (43)
Edgerton	986	4.3 (115)	56.1 (32)	935	22.0 (76)	45.3 (73)	279	32.6 (124)	170	20.0 (109)	10	100.0 (34)	49	100.0 (43)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Eldora	18	0.0 (1)	61.1 (70)	17	11.8 (21)	41.2 (54)	5	0.0 (1)	0	0.0 (1)	1	100.0 (34)	0	0.0 (1)
Elkader	77	1.3 (48)	64.9 (100)	73	20.5 (60)	54.8 (140)	27	33.3 (125)	25	24.0 (116)	3	100.0 (34)	3	100.0 (43)
Emmetsburg	45	0.0 (1)	71.1 (138)	41	9.8 (15)	46.3 (80)	14	0.0 (1)	11	9.1 (87)	2	100.0 (34)	1	100.0 (43)
Esthervill	101	3.0 (80)	67.3 (122)	95	21.1 (67)	51.6 (123)	33	3.0 (68)	27	3.7 (68)	16	100.0 (34)	4	100.0 (43)
Fairfiel	53	0.0 (1)	73.6 (142)	50	22.0 (76)	48.0 (93)	21	38.1 (147)	17	17.6 (104)	0	0.0 (1)	2	100.0 (43)
Fayette	15	6.7 (143)	53.3 (23)	14	50.0 (161)	35.7 (24)	3	0.0 (1)	5	20.0 (109)	0	0.0 (1)	1	100.0 (43)
Forest City	107	2.8 (75)	68.2 (128)	102	16.7 (28)	49.0 (107)	28	7.1 (83)	20	10.0 (93)	10	100.0 (34)	10	100.0 (43)
Fort Dodge	501	2.8 (75)	65.3 (103)	482	22.0 (76)	48.3 (95)	148	3.4 (70)	44	9.1 (87)	24	100.0 (34)	12	100.0 (43)
Ft Madison	131	2.3 (65)	68.7 (132)	123	24.4 (105)	48.8 (104)	47	4.3 (75)	23	4.3 (69)	12	100.0 (34)	5	100.0 (43)
Garner	53	3.8 (105)	67.9 (126)	53	26.4 (119)	47.2 (86)	9	11.1 (92)	10	10.0 (93)	7	100.0 (34)	1	100.0 (43)
Glenwood	54	3.7 (101)	57.4 (39)	50	22.0 (76)	54.0 (137)	21	0.0 (1)	18	0.0 (1)	0	0.0 (1)	1	100.0 (43)
Grand Junction	6	0.0 (1)	33.3 (7)	5	0.0 (1)	40.0 (39)	1	0.0 (1)	2	50.0 (155)	1	100.0 (34)	0	0.0 (1)
Greenfield	36	0.0 (1)	75.0 (145)	35	20.0 (51)	40.0 (39)	8	37.5 (142)	6	16.7 (102)	2	100.0 (34)	1	100.0 (43)
Grinnell	83	7.2 (152)	66.3 (111)	78	17.9 (37)	48.7 (100)	21	38.1 (147)	25	24.0 (116)	5	100.0 (34)	5	100.0 (43)
Grundy Ctr	16	0.0 (1)	75.0 (145)	14	28.6 (131)	28.6 (13)	6	0.0 (1)	3	0.0 (1)	0	0.0 (1)	1	100.0 (43)
Guthrie Ct	44	0.0 (1)	77.3 (150)	38	28.9 (134)	50.0 (111)	10	30.0 (117)	16	25.0 (118)	6	100.0 (34)	3	100.0 (43)
Hampton	106	1.9 (56)	78.3 (155)	102	21.6 (72)	47.1 (83)	17	0.0 (1)	21	4.8 (72)	11	100.0 (34)	3	100.0 (43)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Harlan	79	3.8 (105)	62.0 (79)	77	18.2 (39)	50.6 (117)	24	4.2 (74)	30	10.0 (93)	2	100.0 (34)	2	100.0 (43)
Hawarden	58	1.7 (54)	77.6 (151)	56	21.4 (69)	51.8 (124)	18	0.0 (1)	20	0.0 (1)	3	100.0 (34)	1	100.0 (43)
Hispanic Cln	19	0.0 (1)	78.9 (156)	19	42.1 (157)	26.3 (10)	4	0.0 (1)	3	33.3 (135)	1	100.0 (34)	0	0.0 (1)
Humboldt	66	0.0 (1)	66.7 (113)	62	24.2 (102)	40.3 (43)	10	10.0 (90)	8	25.0 (118)	7	100.0 (34)	0	0.0 (1)
ILH	1,170	6.0 (138)	58.5 (45)	1,132	26.3 (117)	39.9 (38)	327	2.4 (66)	97	6.2 (78)	83	100.0 (34)	33	100.0 (43)
Ida Grove	69	5.8 (133)	69.6 (134)	67	29.9 (138)	47.8 (91)	14	0.0 (1)	22	0.0 (1)	5	100.0 (34)	1	100.0 (43)
Independen	83	0.0 (1)	67.5 (125)	79	21.5 (71)	40.5 (45)	19	0.0 (1)	12	0.0 (1)	3	100.0 (34)	4	100.0 (43)
Indianola	122	4.9 (124)	63.1 (90)	120	25.0 (108)	47.5 (88)	42	7.1 (83)	28	3.6 (66)	10	100.0 (34)	2	100.0 (43)
Iowa City	860	3.7 (101)	57.2 (38)	807	24.7 (106)	39.2 (36)	287	35.9 (138)	337	26.4 (124)	51	100.0 (34)	32	100.0 (43)
Iowa Falls	37	8.1 (156)	56.8 (36)	36	38.9 (156)	30.6 (16)	6	33.3 (125)	14	35.7 (146)	3	100.0 (34)	1	100.0 (43)
Jamaica	7	14.3 (162)	85.7 (159)	7	14.3 (24)	71.4 (158)	3	33.3 (125)	2	0.0 (1)	1	100.0 (34)	1	100.0 (43)
Jefferson	57	7.0 (150)	63.2 (92)	57	19.3 (47)	47.4 (87)	17	47.1 (154)	17	17.6 (104)	7	100.0 (34)	7	100.0 (43)
KR	34	5.9 (135)	55.9 (30)	32	34.4 (151)	37.5 (29)	12	0.0 (1)	3	0.0 (1)	4	100.0 (34)	1	100.0 (43)
Keokuk	142	5.6 (130)	60.6 (66)	136	26.5 (120)	37.5 (29)	49	0.0 (1)	27	11.1 (97)	21	100.0 (34)	14	100.0 (43)
Keosauqua	12	0.0 (1)	41.7 (9)	12	0.0 (1)	25.0 (6)	6	33.3 (125)	6	0.0 (1)	0	0.0 (1)	2	100.0 (43)
Knoxville	164	6.7 (143)	56.1 (32)	157	23.6 (94)	40.8 (47)	38	7.9 (87)	32	6.3 (79)	7	100.0 (34)	6	100.0 (43)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Lamoni	11	9.1 (158)	45.5 (10)	9	33.3 (147)	33.3 (18)	3	66.7 (160)	1	100.0 (162)	2	100.0 (34)	0	0.0 (1)
Le Mars	109	5.5 (129)	59.6 (55)	107	20.6 (62)	48.6 (98)	39	0.0 (1)	28	3.6 (66)	4	100.0 (34)	1	100.0 (43)
Lenox	17	5.9 (135)	58.8 (49)	17	23.5 (91)	52.9 (131)	4	75.0 (162)	3	33.3 (135)	3	100.0 (34)	0	0.0 (1)
Leon	32	6.3 (140)	53.1 (21)	30	36.7 (154)	30.0 (15)	10	20.0 (101)	3	0.0 (1)	0	0.0 (1)	1	100.0 (43)
Logan	76	1.3 (48)	60.5 (64)	74	18.9 (44)	52.7 (128)	23	0.0 (1)	31	0.0 (1)	0	0.0 (1)	3	100.0 (43)
Manchester	51	2.0 (60)	66.7 (113)	51	23.5 (91)	52.9 (131)	13	38.5 (149)	10	20.0 (109)	6	100.0 (34)	1	100.0 (43)
Manning	2	0.0 (1)	50.0 (12)	2	0.0 (1)	100.0 (160)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Maquoketa	75	6.7 (143)	68.0 (127)	74	23.0 (88)	48.6 (98)	10	30.0 (117)	12	50.0 (155)	2	100.0 (34)	3	100.0 (43)
Marengo	19	0.0 (1)	68.4 (130)	18	11.1 (17)	44.4 (68)	5	20.0 (101)	9	33.3 (135)	0	0.0 (1)	1	100.0 (43)
Marion	310	1.3 (48)	65.5 (105)	302	18.9 (44)	52.3 (125)	78	32.1 (122)	60	28.3 (127)	40	100.0 (34)	17	100.0 (43)
Marshalltown	748	3.6 (97)	59.9 (57)	729	24.1 (98)	41.0 (50)	143	14.0 (97)	99	15.2 (101)	61	100.0 (34)	4	100.0 (43)
Mason City	526	3.6 (97)	64.1 (97)	502	18.1 (38)	50.8 (118)	120	5.0 (77)	61	8.2 (85)	52	100.0 (34)	18	100.0 (43)
Monticello	15	6.7 (143)	66.7 (113)	15	26.7 (121)	26.7 (11)	2	50.0 (156)	4	0.0 (1)	3	100.0 (34)	1	100.0 (43)
Monticello	3	0.0 (1)	100.0 (160)	3	33.3 (147)	66.7 (153)	1	0.0 (1)	1	100.0 (162)	0	0.0 (1)	0	0.0 (1)
Mt Ayr	28	7.1 (151)	50.0 (12)	27	29.6 (137)	55.6 (142)	6	33.3 (125)	3	0.0 (1)	2	100.0 (34)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mt Pleasan	121	0.0 (1)	59.5 (54)	115	24.3 (104)	48.7 (100)	44	4.5 (76)	23	0.0 (1)	15	100.0 (34)	8	100.0 (43)
Mtown Dtown	2	50.0 (163)	0.0 (1)	2	100.0 (163)	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Murray	1	0.0 (1)	100.0 (160)	1	0.0 (1)	0.0 (1)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Muscatine	600	3.3 (87)	60.5 (64)	576	18.8 (43)	49.1 (109)	162	0.6 (61)	64	3.1 (61)	54	100.0 (34)	15	100.0 (43)
Nevada	9	0.0 (1)	77.8 (152)	9	0.0 (1)	44.4 (68)	2	0.0 (1)	2	0.0 (1)	1	100.0 (34)	0	0.0 (1)
New Hampto	49	4.1 (111)	63.3 (93)	48	22.9 (86)	41.7 (55)	9	0.0 (1)	12	0.0 (1)	3	100.0 (34)	1	100.0 (43)
Newton	222	3.6 (97)	58.6 (46)	208	25.0 (108)	47.1 (83)	69	26.1 (112)	48	31.3 (133)	25	100.0 (34)	4	100.0 (43)
North Towne	409	3.2 (83)	62.8 (85)	396	20.2 (58)	51.0 (120)	115	37.4 (141)	84	42.9 (153)	49	100.0 (34)	8	100.0 (43)
Northwood	15	0.0 (1)	66.7 (113)	15	0.0 (1)	53.3 (136)	2	0.0 (1)	5	0.0 (1)	3	100.0 (34)	2	100.0 (43)
Norwalk	27	0.0 (1)	63.0 (87)	27	14.8 (27)	66.7 (153)	13	0.0 (1)	4	0.0 (1)	3	100.0 (34)	1	100.0 (43)
Oakridge	6	0.0 (1)	16.7 (4)	4	50.0 (161)	25.0 (6)	2	0.0 (1)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Oelwein	68	1.5 (51)	60.3 (61)	67	20.9 (65)	43.3 (63)	18	33.3 (125)	26	30.8 (131)	9	100.0 (34)	7	100.0 (43)
Onawa	64	1.6 (52)	67.2 (121)	63	14.3 (24)	50.8 (118)	24	0.0 (1)	18	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Orange Cty	60	3.3 (87)	58.3 (44)	57	26.3 (117)	21.1 (3)	17	0.0 (1)	30	3.3 (64)	3	100.0 (34)	1	100.0 (43)
Osage	53	1.9 (56)	69.8 (135)	52	17.3 (32)	40.4 (44)	12	8.3 (88)	10	0.0 (1)	10	100.0 (34)	1	100.0 (43)
Osceola	88	5.7 (132)	56.8 (36)	85	27.1 (126)	35.3 (23)	23	34.8 (137)	12	33.3 (135)	3	100.0 (34)	5	100.0 (43)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Oskaloosa	75	2.7 (70)	58.7 (48)	71	23.9 (97)	40.8 (47)	21	33.3 (125)	24	25.0 (118)	0	0.0 (1)	2	100.0 (43)
Oskaloosa	38	2.6 (69)	57.9 (41)	36	27.8 (129)	58.3 (149)	9	22.2 (105)	7	14.3 (100)	5	100.0 (34)	1	100.0 (43)
Ottumwa	1,039	4.5 (121)	60.8 (67)	977	22.2 (80)	45.4 (75)	309	28.2 (114)	210	25.2 (123)	10	100.0 (34)	27	100.0 (43)
Pella	34	11.8 (161)	61.8 (77)	34	20.6 (62)	52.9 (131)	9	11.1 (92)	6	16.7 (102)	4	100.0 (34)	1	100.0 (43)
Perry	87	3.4 (89)	65.5 (105)	83	24.1 (98)	41.0 (50)	25	4.0 (73)	14	0.0 (1)	6	100.0 (34)	1	100.0 (43)
Perry	39	5.1 (126)	61.5 (74)	37	27.0 (125)	35.1 (22)	14	14.3 (98)	9	11.1 (97)	0	0.0 (1)	2	100.0 (43)
Pocahontas	14	0.0 (1)	64.3 (98)	14	28.6 (131)	35.7 (24)	5	0.0 (1)	0	0.0 (1)	0	0.0 (1)	1	100.0 (43)
Pocahontas	6	0.0 (1)	50.0 (12)	5	0.0 (1)	60.0 (150)	0	0.0 (1)	1	0.0 (1)	2	100.0 (34)	0	0.0 (1)
Pomeroy	1	0.0 (1)	100.0 (160)	1	0.0 (1)	100.0 (160)	1	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Postville	93	2.2 (63)	65.6 (108)	92	26.1 (116)	44.6 (70)	29	41.4 (153)	35	37.1 (147)	14	100.0 (34)	3	100.0 (43)
Red Oak	107	3.7 (101)	68.2 (128)	103	22.3 (83)	52.4 (126)	30	0.0 (1)	20	5.0 (73)	0	0.0 (1)	4	100.0 (43)
Redfield	10	0.0 (1)	70.0 (136)	10	20.0 (51)	50.0 (111)	5	0.0 (1)	1	0.0 (1)	1	100.0 (34)	1	100.0 (43)
Refugee Cln	13	0.0 (1)	76.9 (147)	13	23.1 (89)	38.5 (34)	4	25.0 (107)	7	42.9 (153)	1	100.0 (34)	0	0.0 (1)
Rock Rapid	37	2.7 (70)	70.3 (137)	35	31.4 (142)	57.1 (148)	9	0.0 (1)	17	0.0 (1)	1	100.0 (34)	0	0.0 (1)
Rock Valle	73	2.7 (70)	54.8 (26)	71	19.7 (48)	42.3 (60)	11	0.0 (1)	31	3.2 (62)	5	100.0 (34)	1	100.0 (43)
Rockwell C	41	7.3 (153)	61.0 (69)	40	17.5 (35)	45.0 (71)	13	23.1 (106)	13	23.1 (115)	1	100.0 (34)	3	100.0 (43)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
SDM	611	4.4 (118)	60.9 (68)	586	23.2 (90)	45.7 (76)	178	2.2 (65)	61	3.3 (64)	35	100.0 (34)	23	100.0 (43)
Sac City	34	2.9 (77)	64.7 (99)	33	18.2 (39)	48.5 (97)	11	36.4 (139)	13	30.8 (131)	6	100.0 (34)	2	100.0 (43)
Sheldon	93	3.2 (83)	60.2 (60)	89	23.6 (94)	52.8 (129)	24	0.0 (1)	20	0.0 (1)	5	100.0 (34)	3	100.0 (43)
Shenandoah	57	3.5 (95)	64.9 (100)	56	25.0 (108)	41.1 (52)	15	40.0 (151)	15	40.0 (152)	5	100.0 (34)	2	100.0 (43)
Sibley	26	3.8 (105)	76.9 (147)	25	20.0 (51)	36.0 (26)	10	0.0 (1)	9	0.0 (1)	2	100.0 (34)	2	100.0 (43)
Sidney	22	0.0 (1)	63.6 (95)	21	14.3 (24)	52.4 (126)	10	0.0 (1)	8	0.0 (1)	1	100.0 (34)	0	0.0 (1)
Sigourney	27	0.0 (1)	63.0 (87)	24	25.0 (108)	45.8 (77)	10	30.0 (117)	6	33.3 (135)	0	0.0 (1)	0	0.0 (1)
Sioux Cent	104	0.0 (1)	51.9 (19)	101	44.6 (159)	29.7 (14)	48	2.1 (64)	39	2.6 (58)	4	100.0 (34)	1	100.0 (43)
Sioux City	1,547	2.7 (70)	62.6 (82)	1,498	21.6 (72)	43.7 (64)	524	3.8 (72)	593	4.7 (70)	130	100.0 (34)	52	100.0 (43)
Southridge	254	3.1 (81)	55.1 (27)	245	24.1 (98)	43.7 (64)	67	0.0 (1)	25	8.0 (84)	15	100.0 (34)	9	100.0 (43)
Spencer	156	1.9 (56)	61.5 (74)	146	29.5 (136)	43.2 (62)	43	0.0 (1)	30	0.0 (1)	13	100.0 (34)	7	100.0 (43)
Spirit Lak	49	2.0 (60)	63.3 (93)	44	22.7 (85)	56.8 (146)	12	0.0 (1)	9	22.2 (114)	5	100.0 (34)	2	100.0 (43)
Storm Lake	296	3.4 (89)	58.8 (49)	282	24.1 (98)	39.7 (37)	109	0.9 (62)	39	7.7 (82)	30	100.0 (34)	15	100.0 (43)
Stuart	13	0.0 (1)	61.5 (74)	13	0.0 (1)	69.2 (157)	8	25.0 (107)	7	57.1 (160)	1	100.0 (34)	0	0.0 (1)
Tama	129	2.3 (65)	67.4 (124)	118	30.5 (140)	44.1 (66)	27	25.9 (111)	18	38.9 (151)	16	100.0 (34)	4	100.0 (43)
Tama/Settl	9	0.0 (1)	66.7 (113)	9	11.1 (17)	55.6 (142)	7	0.0 (1)	2	50.0 (155)	2	100.0 (34)	0	0.0 (1)
Tipton	16	6.3 (140)	50.0 (12)	14	28.6 (131)	42.9 (61)	7	28.6 (115)	11	9.1 (87)	2	100.0 (34)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Traer	12	0.0 (1)	66.7 (113)	12	8.3 (14)	25.0 (6)	4	25.0 (107)	3	0.0 (1)	4	100.0 (34)	0	0.0 (1)
UIHC	9	0.0 (1)	22.2 (5)	9	22.2 (80)	55.6 (142)	6	66.7 (160)	2	50.0 (155)	4	100.0 (34)	2	100.0 (43)
Vinton	62	3.2 (83)	61.3 (71)	60	33.3 (147)	41.7 (55)	9	0.0 (1)	15	6.7 (81)	6	100.0 (34)	5	100.0 (43)
WDM	163	1.8 (55)	62.6 (82)	156	22.4 (84)	46.8 (82)	57	5.3 (79)	18	5.6 (75)	9	100.0 (34)	7	100.0 (43)
Wapello	1	0.0 (1)	0.0 (1)	1	0.0 (1)	100.0 (160)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Washington	75	6.7 (143)	62.7 (84)	66	21.2 (68)	51.5 (121)	23	21.7 (104)	19	10.5 (96)	4	100.0 (34)	2	100.0 (43)
Waterloo	1,576	3.6 (97)	61.4 (72)	1,494	20.6 (62)	48.7 (100)	369	6.8 (81)	159	5.0 (73)	101	100.0 (34)	68	100.0 (43)
Waukee	18	5.6 (130)	50.0 (12)	16	6.3 (13)	31.3 (17)	6	16.7 (99)	8	37.5 (148)	1	100.0 (34)	0	0.0 (1)
Waukee	47	6.4 (142)	63.8 (96)	44	18.2 (39)	47.7 (90)	14	0.0 (1)	10	0.0 (1)	3	100.0 (34)	3	100.0 (43)
Waukon	61	1.6 (52)	68.9 (133)	57	28.1 (130)	42.1 (59)	17	29.4 (116)	21	28.6 (128)	7	100.0 (34)	3	100.0 (43)
Waverly	52	5.8 (133)	67.3 (122)	52	26.9 (123)	38.5 (34)	9	11.1 (92)	11	9.1 (87)	1	100.0 (34)	0	0.0 (1)
Wbstr Ct	148	3.4 (89)	58.1 (43)	141	19.9 (50)	46.1 (78)	38	10.5 (91)	22	9.1 (87)	16	100.0 (34)	8	100.0 (43)
West Union	23	4.3 (115)	65.2 (102)	21	23.8 (96)	47.6 (89)	11	18.2 (100)	9	33.3 (135)	5	100.0 (34)	2	100.0 (43)
Williamsburg	1	0.0 (1)	100.0 (160)	1	0.0 (1)	100.0 (160)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)	0	0.0 (1)
Winterset	42	0.0 (1)	71.4 (139)	40	27.5 (128)	37.5 (29)	8	25.0 (107)	13	38.5 (150)	5	100.0 (34)	0	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Iowa	**26,971**	**3.7**	**60.4**	**25,791**	**22.9**	**45.2**	**7,639**	**14.7**	**4,950**	**14.6**	**1,902**	**100.0**	**973**	**100.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley	13	46.2 (9)	7.7 (65)	14	0.0 (77)	11	45.5 (24)	12	91.7 (76)
Adel	9	55.6 (5)	11.1 (94)	14	0.0 (77)	6	66.7 (4)	6	100.0 (1)
Adel	7	0.0 (156)	14.3 (120)	8	0.0 (77)	5	40.0 (27)	5	100.0 (1)
Albia	44	31.8 (77)	18.2 (140)	54	0.0 (77)	31	35.5 (36)	32	93.8 (57)
Algona	91	44.0 (14)	9.9 (86)	99	1.0 (63)	90	21.1 (93)	92	97.8 (33)
Allen WHC	181	37.6 (35)	28.7 (161)	255	2.4 (29)	129	20.9 (98)	153	85.0 (121)
Allison	47	42.6 (18)	6.4 (48)	51	0.0 (77)	41	24.4 (73)	46	82.6 (135)
Altoona	61	27.9 (104)	11.5 (100)	84	1.2 (56)	24	20.8 (99)	27	100.0 (1)
Ames	436	30.7 (89)	14.2 (119)	529	1.5 (54)	362	21.3 (91)	362	87.6 (106)
Anamosa	6	33.3 (56)	16.7 (131)	9	0.0 (77)	8	0.0 (152)	8	100.0 (1)
Anamosa	42	23.8 (133)	7.1 (52)	51	2.0 (40)	27	33.3 (39)	41	90.2 (88)
Ankeny	84	26.2 (117)	8.3 (71)	127	1.6 (48)	43	23.3 (77)	49	93.9 (54)
Atlantic	83	37.3 (36)	3.6 (30)	124	0.8 (68)	10	60.0 (6)	20	95.0 (44)
Audubon	27	37.0 (39)	7.4 (59)	35	0.0 (77)	9	22.2 (80)	10	80.0 (141)
Avoca	17	11.8 (153)	23.5 (157)	22	0.0 (77)	17	23.5 (76)	17	88.2 (103)
Bedford	16	31.3 (81)	0.0 (1)	20	0.0 (77)	5	40.0 (27)	5	80.0 (141)
Belle Plaine	15	33.3 (56)	0.0 (1)	23	4.3 (8)	5	20.0 (106)	9	88.9 (96)
Bloomfield	10	40.0 (24)	10.0 (88)	15	6.7 (2)	2	50.0 (15)	2	100.0 (1)
Boone	47	31.9 (76)	17.0 (135)	56	1.8 (44)	32	28.1 (58)	33	84.8 (124)
Brdlwns RP	2,393	27.7 (107)	13.1 (113)	3,353	1.0 (63)	1,195	13.2 (139)	1,404	85.6 (118)
Brooklyn	12	41.7 (20)	25.0 (158)	15	0.0 (77)	9	55.6 (10)	13	100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Burlington	377	31.3 (81)	5.3 (38)	425	0.5 (75)	337	21.4 (90)	370	94.1 (53)
CHC	370	29.2 (97)	20.3 (150)	485	2.5 (28)	29	13.8 (137)	30	83.3 (127)
CR Urban	1,011	25.0 (124)	10.3 (92)	1,298	1.2 (56)	724	11.7 (142)	930	89.5 (92)
Carlisle	19	26.3 (114)	15.8 (126)	28	0.0 (77)	16	18.8 (113)	18	88.9 (96)
Carroll	153	24.8 (130)	7.2 (56)	210	0.5 (75)	68	20.6 (104)	73	84.9 (123)
Cedar Falls	20	40.0 (24)	15.0 (123)	27	0.0 (77)	10	40.0 (27)	11	81.8 (137)
Centervill	44	31.8 (77)	13.6 (117)	61	3.3 (14)	6	33.3 (39)	7	85.7 (115)
Chariton	8	12.5 (152)	12.5 (107)	8	0.0 (77)	8	37.5 (33)	8	87.5 (108)
Chariton	23	39.1 (27)	8.7 (74)	26	0.0 (77)	18	33.3 (39)	18	66.7 (157)
Charles City	136	45.6 (12)	16.2 (128)	154	0.0 (77)	124	27.4 (60)	136	89.0 (95)
Cherokee	75	42.7 (17)	2.7 (24)	104	1.9 (42)	29	37.9 (31)	31	87.1 (111)
Clarinda	37	54.1 (6)	2.7 (24)	49	6.1 (4)	20	45.0 (25)	19	94.7 (47)
Clarion	103	14.6 (151)	29.1 (162)	118	3.4 (13)	100	6.0 (148)	105	74.3 (152)
Clinton	380	37.1 (37)	8.9 (77)	410	1.0 (63)	226	30.1 (49)	360	86.7 (113)
Colfax	25	36.0 (47)	12.0 (105)	24	0.0 (77)	14	28.6 (52)	16	93.8 (57)
Columbus J	35	25.7 (123)	14.3 (120)	37	0.0 (77)	31	19.4 (110)	33	87.9 (104)
Corning	21	38.1 (33)	4.8 (34)	27	0.0 (77)	11	18.2 (116)	11	100.0 (1)
Correctionvl	5	20.0 (142)	0.0 (1)	6	0.0 (77)	4	0.0 (152)	4	75.0 (148)
Corydon	35	37.1 (37)	2.9 (26)	40	0.0 (77)	31	29.0 (51)	31	93.5 (60)
Council Blfs	936	36.5 (43)	7.1 (52)	1,325	0.7 (72)	452	19.2 (111)	461	87.0 (112)
Cresco	53	41.5 (23)	9.4 (81)	64	0.0 (77)	57	21.1 (93)	58	87.9 (104)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Creston	141	46.1 (11)	7.1 (52)	186	2.7 (24)	72	26.4 (63)	73	93.2 (64)
Dallas Ctr	2	0.0 (156)	0.0 (1)	4	0.0 (77)	0	0.0 (152)	0	0.0 (159)
Decorah	53	20.8 (140)	15.1 (124)	63	0.0 (77)	52	21.2 (92)	53	90.6 (85)
Denison	242	28.9 (101)	6.2 (47)	337	0.0 (77)	78	30.8 (47)	98	92.9 (65)
Dewitt	21	38.1 (33)	19.0 (142)	21	0.0 (77)	14	28.6 (52)	20	80.0 (141)
Dubuque	861	36.0 (47)	13.8 (118)	941	2.8 (21)	793	14.1 (135)	797	91.0 (84)
Dyersville	11	36.4 (44)	9.1 (79)	11	0.0 (77)	10	10.0 (146)	10	100.0 (1)
Eagle Grove	58	19.0 (146)	15.5 (125)	66	1.5 (54)	50	26.0 (65)	55	87.3 (109)
Edgerton	818	33.4 (55)	9.0 (78)	1,124	2.2 (33)	80	18.8 (113)	82	95.1 (43)
Eldora	15	26.7 (112)	13.3 (115)	20	0.0 (77)	12	33.3 (39)	12	100.0 (1)
Elkader	77	26.0 (119)	9.1 (79)	88	0.0 (77)	80	33.8 (38)	81	97.5 (34)
Emmetsburg	48	29.2 (97)	14.6 (122)	50	0.0 (77)	42	59.5 (7)	50	96.0 (37)
Esthervill	94	28.7 (102)	7.4 (59)	112	0.0 (77)	86	50.0 (15)	99	93.9 (54)
Fairfiel	36	27.8 (106)	11.1 (94)	56	0.0 (77)	9	22.2 (80)	11	81.8 (137)
Fayette	13	46.2 (9)	0.0 (1)	16	0.0 (77)	16	31.3 (46)	16	100.0 (1)
Forest City	106	32.1 (70)	9.4 (81)	118	0.8 (68)	107	17.8 (118)	109	87.2 (110)
Fort Dodge	464	30.6 (90)	21.8 (152)	535	2.2 (33)	410	16.1 (128)	464	79.5 (146)
Ft Madison	130	27.7 (107)	3.1 (29)	145	0.7 (72)	123	18.7 (115)	132	93.9 (54)
Garner	54	37.0 (39)	22.2 (153)	60	0.0 (77)	54	20.4 (105)	54	83.3 (127)
Glenwood	41	29.3 (96)	4.9 (36)	65	4.6 (6)	6	16.7 (122)	12	75.0 (148)
Grand Junction	5	60.0 (4)	0.0 (1)	6	0.0 (77)	4	25.0 (68)	4	75.0 (148)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Greenfield	34	32.4 (68)	5.9 (45)	40	0.0 (77)	21	28.6 (52)	19	94.7 (47)
Grinnell	77	31.2 (83)	16.9 (134)	92	3.3 (14)	55	12.7 (140)	57	82.5 (136)
Grundy Ctr	13	30.8 (87)	23.1 (155)	22	4.5 (7)	7	42.9 (26)	7	100.0 (1)
Guthrie Ct	38	50.0 (8)	5.3 (38)	60	1.7 (46)	20	5.0 (150)	21	71.4 (155)
Hampton	88	29.5 (94)	30.7 (163)	110	0.9 (66)	88	22.7 (78)	97	80.4 (140)
Harlan	59	35.6 (49)	5.1 (37)	90	0.0 (77)	6	83.3 (3)	12	100.0 (1)
Hawarden	50	32.0 (73)	6.0 (46)	67	3.0 (19)	13	15.4 (131)	14	92.9 (65)
Hispanic Cln	18	22.2 (138)	11.1 (94)	17	0.0 (77)	12	0.0 (152)	16	93.8 (57)
Humboldt	63	33.3 (56)	19.0 (142)	67	3.0 (19)	64	28.1 (58)	64	89.1 (94)
ILH	966	33.1 (63)	11.7 (101)	1,286	0.8 (68)	319	11.3 (144)	646	88.5 (99)
Ida Grove	61	36.1 (45)	13.1 (113)	72	2.8 (21)	23	21.7 (85)	23	91.3 (80)
Independen	62	38.7 (30)	16.1 (127)	89	0.0 (77)	37	37.8 (32)	45	95.6 (41)
Indianola	116	25.9 (120)	19.0 (142)	139	0.0 (77)	106	21.7 (85)	108	92.6 (69)
Iowa City	705	17.2 (149)	19.7 (146)	1,053	1.7 (46)	256	21.1 (93)	278	88.5 (99)
Iowa Falls	33	42.4 (19)	3.0 (27)	37	0.0 (77)	26	15.4 (131)	26	92.3 (72)
Jamaica	7	42.9 (16)	0.0 (1)	9	0.0 (77)	2	50.0 (15)	2	100.0 (1)
Jefferson	52	25.0 (124)	9.6 (83)	67	0.0 (77)	17	17.6 (119)	18	94.4 (51)
KR	28	32.1 (70)	17.9 (138)	41	0.0 (77)	15	40.0 (27)	19	94.7 (47)
Keokuk	136	22.8 (136)	7.4 (59)	159	0.6 (74)	121	20.7 (100)	136	95.6 (41)
Keosauqua	12	41.7 (20)	0.0 (1)	14	0.0 (77)	1	0.0 (152)	1	100.0 (1)
Knoxville	140	27.9 (104)	17.9 (138)	185	1.6 (48)	125	21.6 (88)	132	89.4 (93)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Lamoni	11	27.3 (110)	18.2 (140)	15	0.0 (77)	7	14.3 (134)	7	100.0 (1)
Le Mars	86	24.4 (132)	7.0 (51)	119	0.0 (77)	33	24.2 (74)	33	97.0 (35)
Lenox	17	29.4 (95)	11.8 (103)	21	0.0 (77)	6	50.0 (15)	6	83.3 (127)
Leon	30	30.0 (91)	13.3 (115)	33	0.0 (77)	15	13.3 (138)	14	85.7 (115)
Logan	65	29.2 (97)	10.8 (93)	95	1.1 (59)	13	30.8 (47)	15	66.7 (157)
Manchester	49	22.4 (137)	8.2 (69)	52	0.0 (77)	43	11.6 (143)	46	95.7 (39)
Manning	1	100.0 (1)	0.0 (1)	2	0.0 (77)	0	0.0 (152)	0	0.0 (159)
Maquoketa	72	36.1 (45)	5.6 (42)	77	0.0 (77)	43	25.6 (66)	66	86.4 (114)
Marengo	13	38.5 (31)	0.0 (1)	30	3.3 (14)	6	16.7 (122)	7	71.4 (155)
Marion	303	31.7 (79)	12.9 (111)	367	0.0 (77)	246	22.4 (79)	262	92.4 (70)
Marshalltown	598	39.0 (29)	7.2 (56)	793	0.9 (66)	522	12.5 (141)	568	85.2 (119)
Mason City	493	36.9 (42)	20.1 (149)	566	1.6 (48)	459	21.1 (93)	500	85.2 (119)
Monticello	15	53.3 (7)	0.0 (1)	18	0.0 (77)	10	30.0 (50)	14	92.9 (65)
Monticello	4	25.0 (124)	0.0 (1)	4	0.0 (77)	4	50.0 (15)	4	100.0 (1)
Mt Ayr	23	39.1 (27)	4.3 (32)	28	3.6 (11)	11	27.3 (61)	11	100.0 (1)
Mt Pleasan	119	27.7 (107)	7.6 (64)	141	2.8 (21)	111	20.7 (100)	121	87.6 (106)
Mtown Dtown	1	0.0 (156)	0.0 (1)	2	0.0 (77)	0	0.0 (152)	0	0.0 (159)
Murray	1	0.0 (156)	0.0 (1)	1	0.0 (77)	0	0.0 (152)	0	0.0 (159)
Muscatine	594	35.4 (50)	7.1 (52)	625	1.8 (44)	595	16.3 (127)	614	92.3 (72)
Nevada	9	44.4 (13)	11.1 (94)	11	0.0 (77)	6	50.0 (15)	6	83.3 (127)
New Hampto	35	31.4 (80)	17.1 (136)	51	2.0 (40)	29	51.7 (14)	31	93.5 (60)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Newton	186	25.8 (122)	7.5 (62)	246	1.6 (48)	156	15.4 (131)	165	89.7 (90)
North Towne	374	24.6 (131)	8.8 (76)	474	1.1 (59)	233	21.9 (84)	354	91.2 (81)
Northwood	12	41.7 (20)	25.0 (158)	15	0.0 (77)	12	25.0 (68)	14	100.0 (1)
Norwalk	23	17.4 (148)	8.7 (74)	33	0.0 (77)	19	15.8 (130)	20	85.0 (121)
Oakridge	5	20.0 (142)	0.0 (1)	8	0.0 (77)	2	50.0 (15)	3	100.0 (1)
Oelwein	64	32.8 (65)	7.8 (67)	76	0.0 (77)	69	24.6 (72)	68	91.2 (81)
Onawa	52	25.0 (124)	5.8 (44)	74	0.0 (77)	5	20.0 (106)	9	100.0 (1)
Orange Cty	51	33.3 (56)	11.8 (103)	76	3.9 (9)	11	27.3 (61)	12	100.0 (1)
Osage	55	34.5 (53)	12.7 (109)	61	0.0 (77)	51	21.6 (88)	56	91.1 (83)
Osceola	72	29.2 (97)	12.5 (107)	101	0.0 (77)	39	17.9 (117)	40	90.0 (89)
Oskaloosa	33	33.3 (56)	3.0 (27)	37	2.7 (24)	30	16.7 (122)	35	91.4 (79)
Oskaloosa	61	32.8 (65)	8.2 (69)	85	2.4 (29)	25	20.0 (106)	26	92.3 (72)
Ottumwa	804	31.0 (85)	26.7 (160)	1,081	2.6 (27)	161	21.1 (93)	162	75.9 (147)
Pella	25	32.0 (73)	20.0 (147)	38	0.0 (77)	25	36.0 (35)	27	88.9 (96)
Perry	30	23.3 (134)	10.0 (88)	45	2.2 (33)	24	25.0 (68)	24	83.3 (127)
Perry	81	25.9 (120)	9.9 (86)	92	3.3 (14)	46	10.9 (145)	46	93.5 (60)
Pocahontas	6	16.7 (150)	16.7 (131)	9	0.0 (77)	7	57.1 (8)	7	85.7 (115)
Pocahontas	13	38.5 (31)	7.7 (65)	14	0.0 (77)	11	36.4 (34)	13	84.6 (125)
Pomeroy	1	0.0 (156)	0.0 (1)	1	0.0 (77)	1	0.0 (152)	1	100.0 (1)
Postville	87	19.5 (145)	23.0 (154)	101	0.0 (77)	81	22.2 (80)	84	84.5 (126)
Red Oak	84	32.1 (70)	4.8 (34)	122	0.0 (77)	19	26.3 (64)	30	73.3 (153)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Redfield	8	25.0 (124)	0.0 (1)	9	0.0 (77)	3	66.7 (4)	4	100.0 (1)
Refugee Cln	12	0.0 (156)	0.0 (1)	15	0.0 (77)	10	10.0 (146)	11	72.7 (154)
Rock Rapid	30	26.7 (112)	6.7 (49)	49	0.0 (77)	12	33.3 (39)	13	100.0 (1)
Rock Valle	62	32.3 (69)	11.3 (98)	90	2.2 (33)	17	17.6 (119)	18	83.3 (127)
Rockwell C	41	43.9 (15)	9.8 (84)	49	0.0 (77)	15	20.0 (106)	16	81.3 (139)
SDM	491	31.2 (83)	12.2 (106)	658	1.1 (59)	145	20.7 (100)	214	88.3 (102)
Sac City	30	30.0 (91)	10.0 (88)	37	2.7 (24)	9	22.2 (80)	10	80.0 (141)
Sheldon	92	32.6 (67)	4.3 (32)	108	1.9 (42)	77	55.8 (9)	93	94.6 (50)
Shenandoah	54	37.0 (39)	5.6 (42)	64	1.6 (48)	26	34.6 (37)	24	91.7 (76)
Sibley	26	23.1 (135)	3.8 (31)	34	0.0 (77)	21	52.4 (13)	28	92.9 (65)
Sidney	19	26.3 (114)	5.3 (38)	28	3.6 (11)	2	100.0 (1)	3	100.0 (1)
Sigourney	19	26.3 (114)	5.3 (38)	30	3.3 (14)	6	16.7 (122)	6	100.0 (1)
Sioux Cent	92	10.9 (154)	9.8 (84)	133	0.0 (77)	18	5.6 (149)	20	95.0 (44)
Sioux City	1,314	29.8 (93)	8.5 (73)	1,756	0.8 (68)	335	25.4 (67)	430	91.6 (78)
Southridge	195	27.2 (111)	12.8 (110)	280	2.1 (39)	37	2.7 (151)	62	91.9 (75)
Spencer	149	30.9 (86)	10.1 (91)	165	1.2 (56)	137	46.0 (23)	157	92.4 (70)
Spirit Lak	44	20.5 (141)	11.4 (99)	56	5.4 (5)	39	53.8 (12)	48	95.8 (38)
Storm Lake	279	18.3 (147)	7.5 (62)	330	0.0 (77)	226	48.2 (22)	280	94.3 (52)
Stuart	12	8.3 (155)	16.7 (131)	16	0.0 (77)	6	33.3 (39)	6	100.0 (1)
Tama	103	33.0 (64)	11.7 (101)	135	6.7 (2)	87	20.7 (100)	97	89.7 (90)
Tama/Settl	5	0.0 (156)	20.0 (147)	9	0.0 (77)	6	16.7 (122)	6	83.3 (127)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Tipton	12	25.0 (124)	8.3 (71)	21	0.0 (77)	1	0.0 (152)	2	100.0 (1)
Traer	7	28.6 (103)	0.0 (1)	13	0.0 (77)	6	33.3 (39)	6	100.0 (1)
UIHC	5	20.0 (142)	0.0 (1)	16	0.0 (77)	4	0.0 (152)	4	100.0 (1)
Vinton	51	33.3 (56)	21.6 (151)	65	0.0 (77)	25	16.0 (129)	28	96.4 (36)
WDM	125	32.0 (73)	8.0 (68)	180	2.2 (33)	43	14.0 (136)	59	94.9 (46)
Wapello	1	0.0 (156)	0.0 (1)	1	0.0 (77)	1	100.0 (1)	1	100.0 (1)
Washington	62	21.0 (139)	12.9 (111)	88	1.1 (59)	25	24.0 (75)	31	90.3 (87)
Waterloo	1,176	34.3 (54)	19.2 (145)	1,712	2.3 (32)	770	19.2 (111)	902	88.4 (101)
Waukee	43	39.5 (26)	16.3 (130)	57	0.0 (77)	28	28.6 (52)	29	82.8 (134)
Waukee	13	30.8 (87)	0.0 (1)	21	9.5 (1)	8	25.0 (68)	8	100.0 (1)
Waukon	60	33.3 (56)	6.7 (49)	64	1.6 (48)	63	28.6 (52)	63	90.5 (86)
Waverly	41	61.0 (3)	7.3 (58)	55	0.0 (77)	27	55.6 (10)	30	93.3 (63)
Wbstr Ct	146	34.9 (52)	23.3 (156)	165	2.4 (29)	136	17.6 (119)	142	79.6 (145)
West Union	23	26.1 (118)	17.4 (137)	26	3.8 (10)	23	21.7 (85)	23	95.7 (39)
Williamsburg	1	100.0 (1)	0.0 (1)	1	0.0 (77)	0	0.0 (152)	0	0.0 (159)
Winterset	37	35.1 (51)	16.2 (128)	46	2.2 (33)	14	28.6 (52)	12	75.0 (148)
Iowa	**23,093**	**30.9**	**12.6**	**30,046**	**1.4**	**13,402**	**20.6**	**15,456**	**88.7**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley	13	30.8 (82)	13	7.7 (27)	4	75.0 (5)	13	0.0 (1)	13	0.0 (1)
Adel	8	25.0 (46)	9	22.2 (112)	2	0.0 (136)	8	0.0 (1)	9	0.0 (1)
Adel	6	16.7 (30)	6	16.7 (71)	0	0.0 (136)	6	0.0 (1)	7	0.0 (1)
Albia	40	42.5 (135)	49	28.6 (141)	17	5.9 (134)	44	13.6 (136)	49	2.0 (95)
Algona	95	30.5 (80)	105	21.0 (103)	29	27.6 (68)	93	6.5 (79)	96	1.0 (75)
Allen WHC	212	26.4 (57)	147	14.3 (58)	49	34.7 (45)	212	8.0 (97)	153	2.0 (95)
Allison	48	29.2 (72)	58	19.0 (85)	14	21.4 (105)	47	2.1 (46)	52	1.9 (94)
Altoona	43	30.2 (79)	53	13.2 (51)	10	30.0 (58)	43	0.0 (1)	58	1.7 (90)
Ames	496	24.8 (45)	552	13.8 (53)	119	37.0 (35)	501	8.8 (109)	534	1.3 (80)
Anamosa	38	57.9 (158)	44	29.5 (148)	24	25.0 (79)	42	7.1 (84)	53	5.7 (137)
Anamosa	8	25.0 (46)	10	10.0 (35)	2	0.0 (136)	8	0.0 (1)	9	0.0 (1)
Ankeny	68	20.6 (36)	86	14.0 (57)	17	35.3 (43)	69	1.4 (41)	88	0.0 (1)
Atlantic	87	47.1 (148)	55	32.7 (151)	40	22.5 (97)	87	8.0 (97)	54	0.0 (1)
Audubon	28	46.4 (147)	32	28.1 (140)	14	35.7 (40)	28	14.3 (140)	32	9.4 (148)
Avoca	18	38.9 (115)	13	7.7 (27)	7	42.9 (22)	18	5.6 (68)	12	0.0 (1)
Bedford	18	66.7 (160)	14	28.6 (141)	9	44.4 (20)	18	11.1 (125)	14	7.1 (145)
Belle Plaine	17	23.5 (43)	9	22.2 (112)	3	0.0 (136)	17	0.0 (1)	9	0.0 (1)
Bloomfield	11	36.4 (102)	10	20.0 (93)	4	50.0 (11)	11	9.1 (111)	8	12.5 (154)
Boone	54	50.0 (152)	66	34.8 (153)	27	22.2 (98)	55	9.1 (111)	62	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Brdlwns RP	1,938	21.7 (40)	1,924	13.4 (52)	416	24.0 (90)	2,052	5.3 (64)	2,067	1.5 (84)
Brooklyn	13	38.5 (112)	14	14.3 (58)	5	40.0 (26)	13	0.0 (1)	12	0.0 (1)
Burlington	388	44.1 (140)	493	22.1 (111)	171	36.8 (36)	386	13.7 (137)	483	3.7 (121)
CHC	388	36.3 (101)	338	22.5 (115)	111	18.0 (113)	389	12.6 (134)	333	6.0 (141)
CR Urban	1,066	29.7 (73)	1,213	16.7 (71)	351	31.6 (54)	1,121	7.7 (92)	1,357	1.0 (75)
Carlisle	25	16.0 (28)	23	8.7 (31)	3	100.0 (1)	25	8.0 (97)	24	4.2 (125)
Carroll	159	42.1 (134)	199	21.6 (107)	70	38.6 (30)	158	7.6 (89)	198	1.5 (84)
Cedar Falls	20	10.0 (14)	21	0.0 (1)	1	100.0 (1)	21	4.8 (59)	22	4.5 (129)
Centervill	48	39.6 (122)	28	28.6 (141)	17	0.0 (136)	46	15.2 (148)	24	0.0 (1)
Chariton	8	25.0 (46)	8	37.5 (154)	2	50.0 (11)	8	0.0 (1)	11	0.0 (1)
Chariton	26	26.9 (62)	27	22.2 (112)	7	28.6 (61)	26	3.8 (52)	27	0.0 (1)
Charles City	144	30.6 (81)	182	21.4 (106)	48	25.0 (79)	142	4.9 (60)	174	0.6 (71)
Cherokee	81	39.5 (121)	68	23.5 (122)	27	14.8 (116)	81	8.6 (107)	66	4.5 (129)
Clarinda	41	46.3 (146)	40	27.5 (139)	18	22.2 (98)	41	12.2 (129)	38	5.3 (133)
Clarion	110	12.7 (19)	123	13.0 (50)	15	20.0 (107)	113	6.2 (76)	116	1.7 (90)
Clinton	408	40.7 (128)	461	23.0 (118)	152	25.7 (77)	408	5.1 (62)	450	1.8 (93)
Colfax	19	36.8 (106)	28	25.0 (128)	7	14.3 (117)	19	10.5 (120)	28	3.6 (120)
Columbus J	37	2.7 (9)	38	2.6 (15)	1	0.0 (136)	37	2.7 (48)	37	0.0 (1)
Corning	23	52.2 (155)	23	43.5 (159)	13	7.7 (129)	23	17.4 (153)	22	4.5 (129)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Correctionvl	5	40.0 (124)	4	25.0 (128)	2	50.0 (11)	5	0.0 (1)	4	0.0 (1)
Corydon	28	28.6 (70)	29	31.0 (150)	8	0.0 (136)	29	0.0 (1)	31	6.5 (143)
Council Blfs	1,010	37.1 (107)	894	23.9 (124)	353	22.7 (96)	1,005	6.1 (74)	909	1.1 (77)
Cresco	60	35.0 (93)	55	20.0 (93)	18	33.3 (47)	62	12.9 (135)	54	5.6 (136)
Creston	157	35.0 (93)	160	25.0 (128)	51	13.7 (119)	158	14.6 (145)	148	1.4 (82)
Dallas Ctr	0	0.0 (1)	4	0.0 (1)	0	0.0 (136)	0	0.0 (1)	4	0.0 (1)
Decorah	54	35.2 (96)	69	21.7 (108)	22	31.8 (53)	53	15.1 (147)	68	14.7 (157)
Denison	259	10.8 (17)	207	6.8 (22)	25	16.0 (115)	259	1.2 (39)	206	0.0 (1)
Dewitt	20	30.0 (76)	26	23.1 (119)	7	28.6 (61)	20	5.0 (61)	26	7.7 (146)
Dubuque	905	13.7 (22)	998	8.1 (29)	122	27.0 (73)	901	0.1 (37)	982	0.0 (1)
Dyersville	11	36.4 (102)	13	23.1 (119)	4	0.0 (136)	10	0.0 (1)	12	0.0 (1)
Eagle Grove	61	16.4 (29)	65	10.8 (40)	12	41.7 (25)	61	1.6 (43)	65	1.5 (84)
Edgerton	865	28.1 (66)	594	14.5 (63)	204	21.6 (104)	864	10.5 (120)	587	2.2 (100)
Eldora	20	55.0 (157)	19	52.6 (161)	10	0.0 (136)	20	30.0 (161)	19	5.3 (133)
Elkader	82	39.0 (116)	82	18.3 (80)	31	41.9 (24)	81	21.0 (158)	80	16.3 (158)
Emmetsburg	51	35.3 (97)	62	22.6 (117)	16	43.8 (21)	51	13.7 (137)	62	14.5 (156)
Esthervill	102	21.6 (39)	130	13.8 (53)	20	25.0 (79)	102	14.7 (146)	127	8.7 (147)
Fairfiel	42	38.1 (111)	26	19.2 (88)	16	12.5 (121)	42	9.5 (117)	24	0.0 (1)
Fayette	16	43.8 (139)	16	37.5 (154)	7	28.6 (61)	16	18.8 (155)	16	18.8 (159)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Forest City	110	30.0 (76)	126	12.7 (47)	31	35.5 (41)	106	3.8 (52)	119	0.8 (73)
Fort Dodge	505	39.6 (122)	571	21.7 (108)	197	38.1 (31)	499	8.4 (104)	560	2.3 (103)
Ft Madison	135	43.0 (137)	175	21.1 (104)	55	34.5 (46)	133	11.3 (127)	173	1.2 (78)
Garner	57	26.3 (56)	71	15.5 (66)	15	26.7 (74)	55	9.1 (111)	65	0.0 (1)
Glenwood	46	34.8 (91)	29	10.3 (36)	11	27.3 (69)	45	2.2 (47)	29	0.0 (1)
Grand Junction	6	66.7 (160)	5	20.0 (93)	3	66.7 (6)	6	0.0 (1)	5	0.0 (1)
Greenfield	36	25.0 (46)	31	19.4 (90)	8	12.5 (121)	35	2.9 (49)	29	0.0 (1)
Grinnell	73	30.1 (78)	89	19.1 (86)	24	29.2 (59)	74	18.9 (156)	88	5.7 (137)
Grundy Ctr	14	14.3 (23)	14	0.0 (1)	2	0.0 (136)	14	7.1 (84)	16	0.0 (1)
Guthrie Ct	43	34.9 (92)	53	20.8 (100)	11	18.2 (111)	43	9.3 (115)	53	5.7 (137)
Hampton	102	20.6 (36)	124	12.9 (48)	21	23.8 (91)	102	1.0 (38)	118	0.8 (73)
Harlan	64	39.1 (118)	52	21.2 (105)	23	26.1 (76)	64	6.3 (77)	51	0.0 (1)
Hawarden	53	17.0 (31)	35	14.3 (58)	11	36.4 (38)	53	1.9 (44)	35	5.7 (137)
Hispanic Cln	17	0.0 (1)	18	0.0 (1)	1	100.0 (1)	18	0.0 (1)	18	0.0 (1)
Humboldt	66	39.4 (120)	67	20.9 (102)	26	50.0 (11)	66	6.1 (74)	61	3.3 (115)
ILH	663	25.0 (46)	666	18.5 (82)	183	19.7 (110)	889	7.6 (89)	1,013	2.5 (105)
Ida Grove	63	31.7 (85)	51	29.4 (146)	19	5.3 (135)	61	8.2 (101)	51	3.9 (124)
Independen	66	40.9 (130)	48	33.3 (152)	26	7.7 (129)	65	9.2 (114)	44	0.0 (1)
Indianola	114	37.7 (109)	136	19.1 (86)	42	31.0 (56)	116	13.8 (139)	137	4.4 (127)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Iowa City	733	13.4 (21)	579	6.0 (20)	70	31.4 (55)	729	3.8 (52)	567	0.4 (70)
Iowa Falls	34	26.5 (58)	35	14.3 (58)	9	33.3 (47)	35	14.3 (140)	31	0.0 (1)
Jamaica	6	50.0 (152)	8	12.5 (45)	3	33.3 (47)	6	16.7 (149)	9	0.0 (1)
Jefferson	54	40.7 (128)	68	20.6 (98)	24	37.5 (33)	54	9.3 (115)	68	2.9 (111)
KR	32	25.0 (46)	22	18.2 (78)	7	28.6 (61)	32	6.3 (77)	19	0.0 (1)
Keokuk	141	45.4 (143)	166	30.1 (149)	64	21.9 (103)	140	19.3 (157)	163	3.7 (121)
Keosauqua	12	41.7 (132)	5	0.0 (1)	5	0.0 (136)	12	33.3 (162)	5	0.0 (1)
Knoxville	143	37.8 (110)	153	26.8 (136)	48	16.7 (114)	151	8.6 (107)	151	3.3 (115)
Lamoni	13	23.1 (41)	7	0.0 (1)	2	0.0 (136)	12	0.0 (1)	6	0.0 (1)
Le Mars	90	23.3 (42)	84	10.7 (38)	19	36.8 (36)	89	0.0 (1)	79	1.3 (80)
Lenox	15	26.7 (60)	23	4.3 (18)	4	25.0 (79)	16	0.0 (1)	20	0.0 (1)
Leon	30	33.3 (87)	28	14.3 (58)	10	10.0 (126)	30	10.0 (118)	27	11.1 (151)
Logan	66	28.8 (71)	52	25.0 (128)	17	11.8 (125)	66	4.5 (58)	51	0.0 (1)
Manchester	48	31.3 (84)	52	19.2 (88)	16	6.3 (133)	48	0.0 (1)	51	0.0 (1)
Manning	1	0.0 (1)	2	0.0 (1)	0	0.0 (136)	1	0.0 (1)	2	0.0 (1)
Maquoketa	75	42.7 (136)	78	23.1 (119)	30	26.7 (74)	73	5.5 (66)	77	2.6 (107)
Marengo	13	38.5 (112)	18	38.9 (156)	7	28.6 (61)	13	7.7 (92)	18	0.0 (1)
Marion	308	35.1 (95)	384	19.8 (92)	127	35.4 (42)	328	7.6 (89)	388	1.5 (84)
Marshalltown	683	15.1 (27)	737	10.7 (38)	102	24.5 (89)	684	5.1 (62)	749	1.2 (78)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Mason City	526	35.9 (98)	639	22.5 (115)	190	23.7 (93)	521	7.9 (96)	613	1.5 (84)
Monticello	4	0.0 (1)	5	20.0 (93)	0	0.0 (136)	4	0.0 (1)	5	0.0 (1)
Monticello	15	13.3 (20)	15	6.7 (21)	2	0.0 (136)	15	0.0 (1)	17	0.0 (1)
Mt Ayr	23	47.8 (149)	19	26.3 (134)	10	20.0 (107)	23	17.4 (153)	18	0.0 (1)
Mt Pleasan	123	34.1 (90)	162	17.3 (76)	42	38.1 (31)	123	7.3 (86)	157	0.0 (1)
Mtown Dtown	2	0.0 (1)	0	0.0 (1)	0	0.0 (136)	2	0.0 (1)	1	0.0 (1)
Murray	0	0.0 (1)	1	100.0 (162)	1	0.0 (136)	0	0.0 (1)	1	100.0 (163)
Muscatine	625	27.5 (65)	740	16.4 (70)	163	35.0 (44)	625	12.2 (129)	728	3.3 (115)
Nevada	8	37.5 (108)	11	0.0 (1)	3	33.3 (47)	9	11.1 (125)	11	0.0 (1)
New Hampto	42	33.3 (87)	37	24.3 (127)	11	18.2 (111)	41	17.1 (152)	34	2.9 (111)
Newton	222	32.0 (86)	247	21.9 (110)	69	23.2 (95)	223	8.1 (100)	240	2.5 (105)
North Towne	385	27.0 (64)	474	14.8 (64)	125	42.4 (23)	414	5.3 (64)	505	2.0 (95)
Northwood	13	38.5 (112)	22	9.1 (32)	5	40.0 (26)	13	7.7 (92)	21	0.0 (1)
Norwalk	24	41.7 (132)	35	25.7 (133)	11	9.1 (127)	24	8.3 (102)	37	2.7 (108)
Oakridge	5	20.0 (35)	5	40.0 (157)	2	0.0 (136)	6	0.0 (1)	5	0.0 (1)
Oelwein	69	36.2 (100)	82	18.3 (80)	24	29.2 (59)	72	23.6 (159)	83	12.0 (153)
Onawa	53	43.4 (138)	41	29.3 (145)	19	21.1 (106)	53	5.7 (70)	40	0.0 (1)
Orange Cty	56	14.3 (23)	40	7.5 (26)	7	28.6 (61)	55	3.6 (51)	39	0.0 (1)
Osage	57	24.6 (44)	78	15.4 (65)	14	50.0 (11)	55	7.3 (86)	66	1.5 (84)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Osceola	84	29.8 (74)	77	19.5 (91)	24	25.0 (79)	83	8.4 (104)	72	1.4 (82)
Oskaloosa	70	40.0 (124)	50	24.0 (125)	22	13.6 (120)	70	14.3 (140)	48	4.2 (125)
Oskaloosa	34	44.1 (140)	37	18.9 (84)	15	40.0 (26)	34	5.9 (72)	37	2.7 (108)
Ottumwa	869	26.9 (62)	563	16.9 (74)	201	13.9 (118)	861	10.7 (122)	531	2.4 (104)
Pella	30	36.7 (105)	34	23.5 (122)	11	27.3 (69)	31	6.5 (79)	34	0.0 (1)
Perry	72	18.1 (33)	68	7.4 (25)	10	50.0 (11)	76	1.3 (40)	64	0.0 (1)
Perry	35	25.7 (54)	43	9.3 (33)	7	57.1 (9)	35	5.7 (70)	43	0.0 (1)
Pocahontas	13	46.2 (144)	16	25.0 (128)	6	33.3 (47)	13	0.0 (1)	16	0.0 (1)
Pocahontas	8	25.0 (46)	9	0.0 (1)	2	50.0 (11)	8	12.5 (132)	9	22.2 (161)
Pomeroy	1	100.0 (162)	1	0.0 (1)	1	100.0 (1)	1	0.0 (1)	1	0.0 (1)
Postville	89	9.0 (13)	93	9.7 (34)	8	25.0 (79)	89	5.6 (68)	90	3.3 (115)
Red Oak	89	52.8 (156)	66	24.2 (126)	39	25.6 (78)	89	6.7 (82)	67	0.0 (1)
Redfield	4	50.0 (152)	2	50.0 (160)	1	0.0 (136)	3	0.0 (1)	2	0.0 (1)
Refugee Cln	11	0.0 (1)	14	0.0 (1)	0	0.0 (136)	11	0.0 (1)	11	0.0 (1)
Rock Rapid	33	36.4 (102)	36	13.9 (55)	11	27.3 (69)	33	3.0 (50)	36	0.0 (1)
Rock Valle	64	7.8 (12)	60	8.3 (30)	4	0.0 (136)	65	1.5 (42)	58	0.0 (1)
Rockwell C	42	45.2 (142)	52	26.9 (138)	20	20.0 (107)	42	16.7 (149)	51	2.0 (95)
SDM	316	28.5 (69)	274	20.8 (100)	103	23.3 (94)	356	8.4 (104)	326	3.7 (121)
Sac City	31	48.4 (151)	38	28.9 (144)	16	37.5 (33)	31	6.5 (79)	37	2.7 (108)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Sheldon	96	28.1 (66)	115	12.2 (44)	28	53.6 (10)	98	14.3 (140)	110	5.5 (135)
Shenandoah	56	48.2 (150)	56	26.8 (136)	25	12.0 (124)	56	10.7 (122)	52	0.0 (1)
Sibley	28	17.9 (32)	42	7.1 (23)	5	60.0 (8)	28	10.7 (122)	43	4.7 (132)
Sidney	20	15.0 (26)	15	26.7 (135)	3	0.0 (136)	20	10.0 (118)	15	0.0 (1)
Sigourney	22	40.9 (130)	17	29.4 (146)	9	22.2 (98)	23	4.3 (57)	17	0.0 (1)
Sioux Cent	98	7.1 (11)	78	2.6 (15)	5	40.0 (26)	98	0.0 (1)	75	0.0 (1)
Sioux City	1,382	25.5 (53)	1,074	13.9 (55)	301	24.9 (88)	1,392	5.5 (66)	1,153	1.7 (90)
Southridge	89	28.1 (66)	83	20.5 (97)	30	6.7 (132)	103	3.9 (56)	103	2.9 (111)
Spencer	160	40.0 (124)	188	17.0 (75)	61	36.1 (39)	159	11.9 (128)	183	9.8 (149)
Spirit Lak	49	26.5 (58)	51	15.7 (67)	9	22.2 (98)	49	14.3 (140)	50	20.0 (160)
Storm Lake	291	10.3 (16)	378	3.7 (17)	29	65.5 (7)	292	3.8 (52)	361	4.4 (127)
Stuart	13	46.2 (144)	12	16.7 (71)	6	33.3 (47)	13	0.0 (1)	13	0.0 (1)
Tama	114	21.1 (38)	124	10.5 (37)	26	30.8 (57)	116	6.0 (73)	130	3.1 (114)
Tama/Settl	7	14.3 (23)	9	11.1 (41)	0	0.0 (136)	7	0.0 (1)	9	11.1 (151)
Tipton	13	61.5 (159)	15	40.0 (157)	8	12.5 (121)	13	7.7 (92)	15	6.7 (144)
Traer	10	10.0 (14)	11	18.2 (78)	2	50.0 (11)	10	0.0 (1)	10	10.0 (150)
UIHC	5	40.0 (124)	19	15.8 (68)	2	0.0 (136)	5	0.0 (1)	16	6.3 (142)
Vinton	52	30.8 (82)	39	5.1 (19)	12	25.0 (79)	54	16.7 (149)	37	0.0 (1)
WDM	97	18.6 (34)	84	7.1 (23)	14	7.1 (131)	104	1.9 (44)	91	2.2 (100)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Wapello	1	100.0 (162)	1	100.0 (162)	1	0.0 (136)	1	100.0 (163)	1	0.0 (1)
Washington	67	29.9 (75)	45	17.8 (77)	12	8.3 (128)	67	7.5 (88)	45	2.2 (100)
Waterloo	1,289	36.0 (99)	991	20.7 (99)	412	23.8 (91)	1,315	9.0 (110)	1,045	2.0 (95)
Waukee	16	6.3 (10)	17	11.8 (43)	2	0.0 (136)	16	0.0 (1)	17	0.0 (1)
Waukee	44	11.4 (18)	44	0.0 (1)	4	25.0 (79)	44	0.0 (1)	46	0.0 (1)
Waukon	62	25.8 (55)	62	12.9 (48)	14	28.6 (61)	64	12.5 (132)	58	13.8 (155)
Waverly	45	26.7 (60)	32	12.5 (45)	9	22.2 (98)	45	6.7 (82)	34	0.0 (1)
Wbstr Ct	156	34.0 (89)	175	16.0 (69)	55	45.5 (19)	156	8.3 (102)	171	0.6 (71)
West Union	23	39.1 (118)	26	11.5 (42)	8	25.0 (79)	26	26.9 (160)	26	23.1 (162)
Williamsburg	1	0.0 (1)	1	0.0 (1)	0	0.0 (136)	1	0.0 (1)	1	0.0 (1)
Winterset	41	39.0 (116)	32	18.8 (83)	11	27.3 (69)	41	12.2 (129)	30	3.3 (115)
Iowa	**23,222**	**28.5**	**22,919**	**16.8**	**6,329**	**27.0**	**23,756**	**7.4**	**23,482**	**2.2**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley	14	0.0 (1)	14.3 (149)	14	14.3 (137)	12	0.0 (1)	14	64.3 (123)
Adel	9	0.0 (1)	33.3 (162)	9	0.0 (1)	9	0.0 (1)	9	66.7 (110)
Adel	18	5.6 (63)	11.1 (119)	12	0.0 (1)	12	0.0 (1)	18	72.2 (74)
Albia	61	4.9 (53)	11.5 (129)	55	9.1 (70)	50	0.0 (1)	61	65.6 (120)
Algona	105	6.7 (86)	17.1 (155)	95	9.5 (79)	85	2.4 (81)	106	67.0 (109)
Allen WHC	241	5.8 (67)	6.6 (48)	237	7.6 (53)	213	2.3 (79)	247	69.2 (94)
Allison	60	11.7 (153)	8.3 (79)	51	9.8 (81)	46	4.3 (128)	60	80.0 (32)
Altoona	102	2.9 (30)	6.9 (53)	84	3.6 (27)	81	0.0 (1)	103	70.9 (84)
Ames	650	8.3 (112)	10.0 (107)	510	10.0 (83)	458	3.1 (103)	650	80.5 (29)
Anamosa	54	5.6 (63)	7.4 (62)	49	10.2 (87)	44	0.0 (1)	54	72.2 (74)
Anamosa	10	20.0 (158)	0.0 (1)	9	0.0 (1)	9	11.1 (159)	10	80.0 (32)
Ankeny	140	4.3 (42)	11.4 (126)	121	4.1 (31)	116	1.7 (66)	140	81.4 (24)
Atlantic	122	7.4 (100)	10.7 (115)	119	10.9 (104)	106	2.8 (91)	122	63.9 (126)
Audubon	35	11.4 (151)	2.9 (17)	34	8.8 (66)	31	3.2 (106)	35	77.1 (46)
Avoca	21	4.8 (49)	0.0 (1)	21	4.8 (32)	20	5.0 (136)	21	71.4 (78)
Bedford	24	4.2 (39)	20.8 (160)	20	5.0 (35)	19	0.0 (1)	24	45.8 (159)
Belle Plaine	22	4.5 (46)	13.6 (147)	22	13.6 (132)	19	0.0 (1)	22	68.2 (104)
Bloomfield	14	0.0 (1)	0.0 (1)	14	21.4 (157)	11	0.0 (1)	14	57.1 (145)
Boone	72	8.3 (112)	12.5 (142)	55	3.6 (27)	53	9.4 (156)	72	58.3 (140)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Brdlwns RP	3,518	8.3 (112)	7.2 (58)	3,224	9.4 (77)	2,876	3.5 (117)	3,574	80.2 (31)
Brooklyn	15	0.0 (1)	6.7 (51)	15	33.3 (161)	10	0.0 (1)	16	62.5 (129)
Burlington	507	7.3 (97)	9.3 (99)	413	9.0 (69)	375	3.5 (117)	509	59.1 (139)
CHC	627	11.0 (149)	7.2 (58)	458	11.4 (116)	406	5.4 (145)	628	63.2 (128)
CR Urban	1,400	7.6 (102)	8.4 (83)	1,234	9.1 (70)	1,089	2.8 (91)	1,441	68.4 (99)
Carlisle	42	7.1 (90)	4.8 (33)	28	14.3 (137)	24	0.0 (1)	42	73.8 (61)
Carroll	205	6.8 (87)	7.8 (70)	204	10.3 (89)	183	2.2 (77)	205	70.7 (86)
Cedar Falls	32	6.3 (76)	3.1 (19)	29	13.8 (134)	24	0.0 (1)	33	87.9 (10)
Centervill	68	8.8 (121)	7.4 (62)	58	15.5 (150)	49	4.1 (126)	68	54.4 (149)
Chariton	38	5.3 (59)	13.2 (145)	26	15.4 (147)	22	0.0 (1)	38	68.4 (99)
Chariton	12	0.0 (1)	8.3 (79)	8	0.0 (1)	8	0.0 (1)	12	41.7 (161)
Charles City	180	8.9 (122)	11.7 (133)	153	13.1 (125)	132	5.3 (142)	181	65.7 (118)
Cherokee	112	9.8 (136)	6.3 (44)	105	6.7 (44)	98	8.2 (153)	112	77.7 (44)
Clarinda	53	5.7 (66)	5.7 (39)	44	9.1 (70)	40	2.5 (84)	52	61.5 (133)
Clarion	124	5.6 (63)	9.7 (104)	111	13.5 (131)	96	2.1 (74)	124	79.0 (39)
Clinton	500	11.4 (151)	8.8 (88)	390	13.1 (125)	339	3.5 (117)	501	69.5 (93)
Colfax	33	0.0 (1)	12.1 (140)	25	4.0 (29)	24	0.0 (1)	33	66.7 (110)
Columbus J	41	9.8 (136)	9.8 (106)	38	10.5 (95)	34	5.9 (147)	42	66.7 (110)
Corning	32	0.0 (1)	9.4 (100)	28	17.9 (155)	23	0.0 (1)	32	68.8 (97)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Correctionvl	6	0.0 (1)	16.7 (153)	6	0.0 (1)	6	0.0 (1)	6	100.0 (1)
Corydon	47	8.5 (117)	19.1 (158)	40	7.5 (50)	37	2.7 (89)	47	57.4 (143)
Council Blfs	1,349	8.4 (116)	7.6 (66)	1,272	11.3 (114)	1,124	2.7 (89)	1,357	75.3 (55)
Cresco	68	10.3 (144)	13.2 (145)	60	11.7 (119)	53	3.8 (122)	68	77.9 (42)
Creston	228	7.9 (108)	6.1 (41)	177	10.7 (98)	158	3.2 (106)	228	68.4 (99)
Dallas Ctr	7	0.0 (1)	14.3 (149)	4	0.0 (1)	4	0.0 (1)	7	100.0 (1)
Decorah	77	7.8 (106)	10.4 (113)	63	6.3 (41)	59	3.4 (112)	77	79.2 (37)
Denison	338	9.2 (128)	6.2 (42)	337	14.8 (144)	287	1.0 (57)	338	65.1 (122)
Dewitt	30	23.3 (160)	3.3 (20)	23	34.8 (162)	15	0.0 (1)	30	66.7 (110)
Dubuque	1,068	6.5 (79)	8.1 (75)	885	6.7 (44)	825	2.9 (94)	1,073	55.6 (146)
Dyersville	13	0.0 (1)	7.7 (67)	11	0.0 (1)	11	0.0 (1)	13	46.2 (158)
Eagle Grove	67	7.5 (101)	7.5 (65)	64	12.5 (124)	56	1.8 (70)	67	76.1 (50)
Edgerton	1,201	9.7 (134)	7.0 (55)	1,089	11.0 (106)	967	4.3 (128)	1,204	65.6 (120)
Eldora	23	17.4 (157)	4.3 (29)	19	21.1 (156)	15	6.7 (150)	23	52.2 (151)
Elkader	98	6.1 (73)	8.2 (77)	85	3.5 (26)	82	2.4 (81)	98	81.6 (23)
Emmetsburg	61	9.8 (136)	6.6 (48)	49	10.2 (87)	44	2.3 (79)	61	70.5 (89)
Esthervill	127	9.4 (131)	9.4 (100)	107	10.3 (89)	95	5.3 (142)	128	68.0 (105)
Fairfiel	58	5.2 (58)	10.3 (111)	54	11.1 (108)	48	2.1 (74)	59	57.6 (142)
Fayette	18	0.0 (1)	16.7 (153)	14	14.3 (137)	12	0.0 (1)	18	66.7 (110)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Forest City	123	8.9 (122)	11.4 (126)	112	10.7 (98)	100	1.0 (57)	123	65.9 (117)
Fort Dodge	567	7.1 (90)	8.1 (75)	515	7.8 (54)	475	2.1 (74)	567	72.5 (70)
Ft Madison	176	9.1 (124)	8.5 (85)	139	8.6 (64)	127	5.5 (146)	176	52.8 (150)
Garner	73	1.4 (28)	6.8 (52)	61	4.9 (34)	58	0.0 (1)	73	71.2 (80)
Glenwood	65	4.6 (48)	6.2 (42)	63	7.9 (57)	58	1.7 (66)	65	69.2 (94)
Grand Junction	5	20.0 (158)	0.0 (1)	5	0.0 (1)	5	20.0 (162)	5	60.0 (135)
Greenfield	44	2.3 (29)	11.4 (126)	39	2.6 (24)	38	2.6 (87)	44	72.7 (68)
Grinnell	102	5.9 (68)	6.9 (53)	86	10.5 (95)	77	2.6 (87)	102	72.5 (70)
Grundy Ctr	24	4.2 (39)	12.5 (142)	20	15.0 (146)	17	0.0 (1)	24	62.5 (129)
Guthrie Ct	55	7.3 (97)	5.5 (38)	53	7.5 (50)	49	2.0 (72)	55	74.5 (58)
Hampton	125	4.0 (37)	12.0 (139)	109	14.7 (143)	92	0.0 (1)	126	71.4 (78)
Harlan	92	5.4 (61)	4.3 (29)	89	10.1 (85)	80	1.3 (61)	92	64.1 (125)
Hawarden	67	6.0 (72)	11.9 (136)	63	11.1 (108)	56	1.8 (70)	67	79.1 (38)
Hispanic Cln	17	5.9 (68)	5.9 (40)	17	5.9 (38)	16	0.0 (1)	17	70.6 (87)
Humboldt	66	10.6 (148)	9.1 (92)	61	11.5 (117)	54	3.7 (120)	66	84.8 (11)
ILH	1,351	9.5 (132)	9.2 (96)	1,249	11.8 (120)	1,095	3.7 (120)	1,357	75.9 (51)
Ida Grove	73	6.8 (87)	4.1 (26)	70	17.1 (153)	58	0.0 (1)	73	72.6 (69)
Independen	97	6.2 (75)	11.3 (125)	85	5.9 (38)	80	1.3 (61)	97	61.9 (132)
Indianola	173	8.7 (120)	11.0 (118)	140	9.3 (76)	127	3.1 (103)	172	76.7 (47)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Iowa City	1,068	9.1 (124)	7.2 (58)	991	9.2 (75)	888	4.3 (128)	1,084	84.4 (13)
Iowa Falls	44	9.1 (124)	4.5 (32)	38	13.2 (127)	33	0.0 (1)	44	79.5 (36)
Jamaica	9	11.1 (150)	11.1 (119)	9	0.0 (1)	9	11.1 (159)	9	100.0 (1)
Jefferson	69	4.3 (42)	8.7 (87)	68	10.3 (89)	61	0.0 (1)	69	71.0 (82)
KR	42	4.8 (49)	19.0 (157)	39	15.4 (147)	33	3.0 (99)	42	73.8 (61)
Keokuk	168	7.7 (103)	3.6 (24)	152	7.9 (57)	138	5.1 (139)	168	44.6 (160)
Keosauqua	15	0.0 (1)	26.7 (161)	15	0.0 (1)	15	0.0 (1)	15	60.0 (135)
Knoxville	220	7.7 (103)	7.7 (67)	178	8.4 (63)	163	2.5 (84)	220	66.4 (116)
Lamoni	17	5.9 (68)	11.8 (134)	13	0.0 (1)	13	0.0 (1)	17	70.6 (87)
Le Mars	131	6.9 (89)	5.3 (37)	119	14.3 (137)	101	1.0 (57)	132	81.8 (21)
Lenox	28	7.1 (90)	7.1 (56)	21	4.8 (32)	20	5.0 (136)	28	50.0 (152)
Leon	48	10.4 (146)	4.2 (27)	34	2.9 (25)	33	3.0 (99)	48	62.5 (129)
Logan	98	10.2 (143)	8.2 (77)	94	9.6 (80)	85	4.7 (133)	98	72.4 (72)
Manchester	60	11.7 (153)	10.0 (107)	51	9.8 (81)	46	6.5 (149)	60	46.7 (157)
Manning	2	0.0 (1)	0.0 (1)	2	0.0 (1)	2	0.0 (1)	2	50.0 (152)
Maquoketa	89	7.9 (108)	7.9 (72)	76	14.5 (142)	65	3.1 (103)	89	57.3 (144)
Marengo	29	10.3 (144)	10.3 (111)	27	7.4 (49)	25	8.0 (152)	29	75.9 (51)
Marion	396	7.3 (97)	11.1 (119)	354	10.7 (98)	314	2.9 (94)	399	67.7 (107)
Marshalltown	850	6.6 (84)	8.4 (83)	781	8.2 (61)	708	3.0 (99)	861	72.1 (76)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Mason City	631	6.5 (79)	9.2 (96)	538	10.0 (83)	483	3.3 (109)	634	67.8 (106)
Monticello	17	0.0 (1)	0.0 (1)	17	5.9 (38)	16	0.0 (1)	17	82.4 (17)
Monticello	5	0.0 (1)	0.0 (1)	4	0.0 (1)	4	0.0 (1)	5	80.0 (32)
Mt Ayr	30	10.0 (140)	3.3 (20)	28	0.0 (1)	28	10.7 (158)	30	63.3 (127)
Mt Pleasan	165	9.7 (134)	9.1 (92)	135	8.1 (59)	124	4.8 (134)	166	73.5 (65)
Mtown Dtown	2	0.0 (1)	0.0 (1)	2	0.0 (1)	2	0.0 (1)	2	50.0 (152)
Murray	1	0.0 (1)	0.0 (1)	1	0.0 (1)	1	0.0 (1)	1	0.0 (162)
Muscatine	736	6.5 (79)	7.2 (58)	606	11.2 (113)	538	2.4 (81)	736	76.5 (48)
Nevada	11	0.0 (1)	9.1 (92)	11	0.0 (1)	10	0.0 (1)	12	75.0 (56)
New Hampto	55	5.5 (62)	10.9 (116)	51	7.8 (54)	47	4.3 (128)	55	74.5 (58)
Newton	271	9.2 (128)	11.8 (134)	233	10.3 (89)	208	2.9 (94)	272	68.4 (99)
North Towne	524	7.1 (90)	9.0 (90)	468	8.3 (62)	427	3.3 (109)	527	69.8 (91)
Northwood	22	9.1 (124)	9.1 (92)	15	6.7 (44)	14	7.1 (151)	22	77.3 (45)
Norwalk	42	7.1 (90)	9.5 (102)	34	17.6 (154)	28	0.0 (1)	42	73.8 (61)
Oakridge	6	0.0 (1)	0.0 (1)	6	16.7 (151)	5	0.0 (1)	6	50.0 (152)
Oelwein	93	6.5 (79)	12.9 (144)	75	10.7 (98)	67	3.0 (99)	93	68.8 (97)
Onawa	78	3.8 (33)	3.8 (25)	74	6.8 (47)	69	1.4 (64)	78	60.3 (134)
Orange Cty	79	5.1 (55)	15.2 (152)	73	11.0 (106)	65	1.5 (65)	79	88.6 (9)
Osage	78	7.7 (103)	6.4 (45)	62	6.5 (42)	58	0.0 (1)	78	75.6 (53)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Osceola	122	8.2 (110)	6.6 (48)	97	13.4 (129)	84	4.8 (134)	122	73.8 (61)
Oskaloosa	91	9.9 (139)	8.8 (88)	80	11.3 (114)	71	2.8 (91)	91	59.3 (138)
Oskaloosa	39	5.1 (55)	7.7 (67)	36	5.6 (37)	34	2.9 (94)	39	66.7 (110)
Ottumwa	1,114	6.3 (76)	7.9 (72)	1,015	8.8 (66)	926	2.2 (77)	1,115	73.3 (66)
Pella	45	4.4 (45)	11.1 (119)	38	10.5 (95)	34	2.9 (94)	45	73.3 (66)
Perry	97	4.1 (38)	5.2 (36)	88	9.1 (70)	80	1.3 (61)	97	84.5 (12)
Perry	43	9.3 (130)	11.6 (132)	43	11.6 (118)	38	5.3 (142)	43	72.1 (76)
Pocahontas	9	0.0 (1)	11.1 (119)	8	0.0 (1)	8	0.0 (1)	9	77.8 (43)
Pocahontas	16	6.3 (76)	18.8 (156)	14	21.4 (157)	11	0.0 (1)	16	81.3 (25)
Pomeroy	1	0.0 (1)	0.0 (1)	1	0.0 (1)	1	0.0 (1)	1	100.0 (1)
Postville	118	5.9 (68)	4.2 (27)	102	11.8 (120)	90	3.3 (109)	118	78.0 (41)
Red Oak	123	4.9 (53)	10.6 (114)	119	13.4 (129)	103	0.0 (1)	123	68.3 (103)
Redfield	10	0.0 (1)	0.0 (1)	10	0.0 (1)	10	0.0 (1)	10	60.0 (135)
Refugee Cln	10	0.0 (1)	20.0 (159)	15	13.3 (128)	10	0.0 (1)	16	93.8 (6)
Rock Rapid	51	7.8 (106)	7.8 (70)	47	14.9 (145)	40	5.0 (136)	51	74.5 (58)
Rock Valle	94	5.3 (59)	7.4 (62)	88	9.1 (70)	79	5.1 (139)	95	82.1 (19)
Rockwell C	52	15.4 (156)	11.5 (129)	49	14.3 (137)	42	9.5 (157)	52	57.7 (141)
SDM	696	8.2 (110)	9.2 (96)	639	10.8 (103)	553	3.4 (112)	719	75.5 (54)
Sac City	36	8.3 (112)	13.9 (148)	36	11.1 (108)	32	0.0 (1)	36	55.6 (146)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Sheldon	123	6.5 (79)	6.5 (46)	106	14.2 (136)	91	1.1 (60)	124	80.6 (27)
Shenandoah	70	2.9 (30)	4.3 (29)	64	9.4 (77)	58	1.7 (66)	70	64.3 (123)
Sibley	42	4.8 (49)	11.9 (136)	34	11.8 (120)	29	3.4 (112)	42	81.0 (26)
Sidney	29	0.0 (1)	3.4 (23)	27	11.1 (108)	24	0.0 (1)	29	72.4 (72)
Sigourney	26	3.8 (33)	11.5 (129)	24	16.7 (151)	20	0.0 (1)	26	69.2 (94)
Sioux Cent	147	6.1 (73)	4.8 (33)	133	7.5 (50)	121	2.5 (84)	149	83.2 (15)
Sioux City	1,806	6.6 (84)	8.3 (79)	1,712	10.3 (89)	1,507	2.0 (72)	1,838	71.0 (82)
Southridge	281	8.5 (117)	10.0 (107)	270	10.7 (98)	235	3.4 (112)	289	80.6 (27)
Spencer	181	3.9 (36)	8.3 (79)	160	8.1 (59)	147	0.0 (1)	182	74.7 (57)
Spirit Lak	52	3.8 (33)	9.6 (103)	51	13.7 (133)	44	0.0 (1)	52	82.7 (16)
Storm Lake	377	7.2 (96)	9.0 (90)	326	10.4 (94)	291	3.4 (112)	379	65.7 (118)
Stuart	20	30.0 (162)	10.0 (107)	16	25.0 (160)	12	8.3 (154)	20	50.0 (152)
Tama	140	7.1 (90)	12.1 (140)	126	8.7 (65)	114	4.4 (132)	141	67.4 (108)
Tama/Settl	10	10.0 (140)	0.0 (1)	9	11.1 (108)	8	0.0 (1)	10	90.0 (8)
Tipton	22	0.0 (1)	0.0 (1)	20	5.0 (35)	19	0.0 (1)	22	54.5 (148)
Traer	14	14.3 (155)	7.1 (56)	13	15.4 (147)	11	9.1 (155)	14	78.6 (40)
UIHC	34	88.2 (163)	2.9 (17)	15	73.3 (163)	4	50.0 (163)	34	91.2 (7)
Vinton	67	3.0 (32)	14.9 (151)	62	6.5 (42)	58	1.7 (66)	67	70.1 (90)
WDM	193	10.4 (146)	10.9 (116)	170	11.8 (120)	149	4.0 (124)	195	80.5 (29)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2017 - 2019 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Wapello	1	0.0 (1)	0.0 (1)	1	0.0 (1)	1	0.0 (1)	1	0.0 (162)
Washington	93	4.3 (42)	6.5 (46)	77	7.8 (54)	71	0.0 (1)	93	81.7 (22)
Waterloo	1,884	8.5 (117)	7.9 (72)	1,637	10.1 (85)	1,462	3.8 (122)	1,897	69.8 (91)
Waukee	18	27.8 (161)	0.0 (1)	18	22.2 (159)	14	14.3 (161)	18	83.3 (14)
Waukee	67	4.5 (46)	11.9 (136)	55	7.3 (48)	50	4.0 (124)	68	82.4 (17)
Waukon	72	4.2 (39)	9.7 (104)	63	1.6 (23)	62	3.2 (106)	72	70.8 (85)
Waverly	63	4.8 (49)	11.1 (119)	55	10.9 (104)	49	6.1 (148)	63	76.2 (49)
Wbstr Ct	178	9.6 (133)	5.1 (35)	160	13.8 (134)	138	5.1 (139)	179	82.1 (19)
West Union	30	10.0 (140)	3.3 (20)	25	4.0 (29)	24	4.2 (127)	30	80.0 (32)
Williamsburg	1	0.0 (1)	100.0 (163)	1	0.0 (1)	1	0.0 (1)	1	100.0 (1)
Winterset	59	5.1 (55)	8.5 (85)	45	8.9 (68)	41	0.0 (1)	59	71.2 (80)
Iowa	**32,588**	**7.8**	**8.3**	**28,952**	**10.0**	**25,853**	**3.1**	**32,859**	**67.7**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.