WIC PNSS & PedNSS

Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Ackley	66.7	0.0	33.3	0.0	0.0	0.0	0.0
Adel	76.9	7.7	15.4	0.0	0.0	0.0	0.0
Albia	94.1	0.0	5.9	0.0	0.0	0.0	0.0
Algona	80.6	0.0	13.9	0.0	0.0	5.6	0.0
Allen WHC	50.5	35.4	5.1	0.0	6.1	3.0	0.0
Allison	86.4	4.5	9.1	0.0	0.0	0.0	0.0
Altoona	72.7	12.1	12.1	0.0	3.0	0.0	0.0
Ames	66.7	10.4	13.1	1.1	6.6	2.2	0.0
Anamosa	92.0	0.0	8.0	0.0	0.0	0.0	0.0
Ankeny	76.5	7.8	7.8	0.0	5.9	2.0	0.0
Atlantic	84.8	0.0	15.2	0.0	0.0	0.0	0.0
Audubon	91.7	0.0	8.3	0.0	0.0	0.0	0.0
Avoca	50.0	50.0	0.0	0.0	0.0	0.0	0.0
Bedford	87.5	0.0	0.0	0.0	0.0	12.5	0.0
Belle Plaine	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Bloomfield	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Boone	76.9	3.8	11.5	0.0	0.0	7.7	0.0
Brdlwns RP	40.1	21.2	23.9	0.2	9.8	4.7	0.0
Brooklyn	83.3	0.0	16.7	0.0	0.0	0.0	0.0
Burlington	75.9	19.8	2.7	0.0	0.5	1.1	0.0
CHC	57.3	21.7	13.6	0.3	2.4	4.5	0.0
CR Urban	46.8	38.0	8.0	0.0	2.7	4.5	0.0
Carlisle	90.0	0.0	10.0	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Carroll	82.1	6.3	9.5	0.0	1.1	1.1	0.0
Cedar Falls	86.7	0.0	6.7	0.0	6.7	0.0	0.0
Centervill	87.5	0.0	8.3	0.0	0.0	4.2	0.0
Chariton	93.8	0.0	6.3	0.0	0.0	0.0	0.0
Charles City	65.5	15.5	12.1	0.0	5.2	1.7	0.0
Cherokee	83.3	0.0	13.9	0.0	0.0	2.8	0.0
Clarinda	95.5	0.0	0.0	0.0	0.0	4.5	0.0
Clarion	37.8	0.0	62.2	0.0	0.0	0.0	0.0
Clinton	77.4	8.7	10.8	0.0	0.5	2.6	0.0
Colfax	90.0	0.0	10.0	0.0	0.0	0.0	0.0
Columbus J	9.1	0.0	63.6	9.1	18.2	0.0	0.0
Corning	90.9	0.0	9.1	0.0	0.0	0.0	0.0
Corydon	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Council Blfs	77.5	4.5	14.8	0.0	1.2	1.9	0.0
Cresco	89.5	0.0	10.5	0.0	0.0	0.0	0.0
Creston	87.5	0.0	10.9	0.0	0.0	1.6	0.0
Dallas Ctr	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Dayton	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Decorah	79.3	0.0	17.2	0.0	0.0	3.4	0.0
Denison	30.6	1.8	60.4	0.0	7.2	0.0	0.0
Dewitt	72.7	9.1	9.1	0.0	0.0	9.1	0.0
Dubuque	74.7	12.5	7.1	0.3	2.7	2.7	0.0
Dyersville	80.0	20.0	0.0	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Eagle Grove	26.1	4.3	69.6	0.0	0.0	0.0	0.0
Edgerton	56.6	26.4	9.9	0.5	1.3	5.3	0.0
Eldora	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Elkader	97.4	0.0	2.6	0.0	0.0	0.0	0.0
Emmetsburg	95.8	0.0	4.2	0.0	0.0	0.0	0.0
Esthervill	66.0	2.0	30.0	0.0	0.0	2.0	0.0
Fairfiel	85.7	0.0	14.3	0.0	0.0	0.0	0.0
Forest City	85.1	6.4	8.5	0.0	0.0	0.0	0.0
Fort Dodge	75.9	8.5	13.2	0.0	0.5	1.9	0.0
Ft Madison	85.2	5.6	9.3	0.0	0.0	0.0	0.0
Garner	85.7	0.0	14.3	0.0	0.0	0.0	0.0
Glenwood	93.8	0.0	6.3	0.0	0.0	0.0	0.0
Grand Junction	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Greenfield	89.5	0.0	10.5	0.0	0.0	0.0	0.0
Grinnell	84.2	10.5	5.3	0.0	0.0	0.0	0.0
Grundy Ctr	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Guthrie Ct	92.9	0.0	7.1	0.0	0.0	0.0	0.0
Hampton	50.9	3.8	43.4	0.0	0.0	1.9	0.0
Harlan	78.0	0.0	19.5	0.0	0.0	2.4	0.0
Hawarden	61.1	0.0	38.9	0.0	0.0	0.0	0.0
Hispanic Cln	0.0	0.0	100.0	0.0	0.0	0.0	0.0
Humboldt	60.0	5.0	35.0	0.0	0.0	0.0	0.0
ILH	38.6	16.1	37.1	0.2	4.9	3.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Ida Grove	91.3	0.0	8.7	0.0	0.0	0.0	0.0
Independen	90.3	3.2	6.5	0.0	0.0	0.0	0.0
Indianola	91.1	3.6	3.6	0.0	1.8	0.0	0.0
Iowa City	34.7	43.6	16.9	0.0	2.8	1.9	0.0
Iowa Falls	82.4	0.0	17.6	0.0	0.0	0.0	0.0
Jamaica	66.7	0.0	33.3	0.0	0.0	0.0	0.0
Jefferson	66.7	0.0	33.3	0.0	0.0	0.0	0.0
KR	50.0	35.7	0.0	0.0	14.3	0.0	0.0
Keokuk	91.1	2.2	4.4	0.0	2.2	0.0	0.0
Keosauqua	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Knoxville	94.3	0.0	4.5	0.0	0.0	1.1	0.0
LCPH	66.7	22.2	5.1	1.0	2.0	3.0	0.0
Lamoni	75.0	12.5	0.0	0.0	0.0	12.5	0.0
Le Mars	73.8	7.1	16.7	0.0	2.4	0.0	0.0
Lenox	50.0	0.0	50.0	0.0	0.0	0.0	0.0
Leon	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Logan	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Manchester	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Maquoketa	82.8	3.4	6.9	0.0	0.0	6.9	0.0
Marengo	66.7	0.0	16.7	0.0	0.0	16.7	0.0
Marion	81.1	5.3	7.7	0.0	0.6	5.3	0.0
Marshalltown	41.2	4.0	45.5	0.0	8.6	0.7	0.0
Mason City	77.6	6.5	10.2	0.4	2.8	2.4	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Monticello	90.0	10.0	0.0	0.0	0.0	0.0	0.0
Mt Ayr	91.7	0.0	8.3	0.0	0.0	0.0	0.0
Mt Pleasan	91.3	0.0	6.5	0.0	2.2	0.0	0.0
Mtown Dtown	0.0	0.0	100.0	0.0	0.0	0.0	0.0
Muscatine	59.9	12.6	25.6	0.0	1.0	1.0	0.0
Nevada	66.7	0.0	33.3	0.0	0.0	0.0	0.0
New Hampto	85.0	5.0	10.0	0.0	0.0	0.0	0.0
Newton	78.2	4.6	11.5	2.3	2.3	1.1	0.0
Northwood	83.3	8.3	0.0	0.0	0.0	8.3	0.0
Norwalk	70.0	0.0	20.0	0.0	10.0	0.0	0.0
Oakridge	25.0	50.0	12.5	0.0	0.0	12.5	0.0
Oelwein	87.0	4.3	8.7	0.0	0.0	0.0	0.0
Onawa	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Orange Cty	45.5	4.5	50.0	0.0	0.0	0.0	0.0
Osage	89.3	3.6	3.6	0.0	0.0	3.6	0.0
Osceola	76.7	0.0	21.7	0.0	1.7	0.0	0.0
Oskaloosa	88.5	3.8	5.8	0.0	1.9	0.0	0.0
Ottumwa	72.7	5.6	15.4	0.0	5.8	0.3	0.3
Pella	90.5	0.0	9.5	0.0	0.0	0.0	0.0
Perry	32.8	13.1	49.2	1.6	0.0	3.3	0.0
Pocahontas	71.4	14.3	14.3	0.0	0.0	0.0	0.0
Postville	25.0	12.5	62.5	0.0	0.0	0.0	0.0
Red Oak	87.9	0.0	12.1	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Redfield	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Refugee Cln	0.0	90.0	10.0	0.0	0.0	0.0	0.0
Rock Rapid	80.8	0.0	15.4	0.0	3.8	0.0	0.0
Rock Valle	31.0	0.0	66.7	2.4	0.0	0.0	0.0
Rockwell C	100.0	0.0	0.0	0.0	0.0	0.0	0.0
SDM	42.8	15.7	28.2	0.5	8.2	4.5	0.0
Sac City	92.3	0.0	0.0	0.0	0.0	7.7	0.0
Sheldon	64.9	2.7	27.0	5.4	0.0	0.0	0.0
Shenandoah	84.4	0.0	15.6	0.0	0.0	0.0	0.0
Sibley	54.5	0.0	45.5	0.0	0.0	0.0	0.0
Sidney	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Sigourney	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Sioux Cent	34.7	0.0	65.3	0.0	0.0	0.0	0.0
Sioux City	37.0	12.3	38.8	4.3	5.4	2.1	0.0
Spencer	86.4	0.0	13.6	0.0	0.0	0.0	0.0
Spirit Lak	91.7	0.0	5.6	0.0	0.0	2.8	0.0
Storm Lake	13.7	4.8	61.3	0.0	19.4	0.8	0.0
Stuart	88.9	0.0	0.0	0.0	0.0	11.1	0.0
Tama	49.0	0.0	36.7	10.2	0.0	4.1	0.0
Tama/Settl	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Tipton	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Traer	100.0	0.0	0.0	0.0	0.0	0.0	0.0
UIHC	50.0	50.0	0.0	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Vinton	96.6	0.0	3.4	0.0	0.0	0.0	0.0
WDM	48.6	26.4	15.3	2.8	5.6	1.4	0.0
Washington	70.6	2.9	26.5	0.0	0.0	0.0	0.0
Waterloo	51.3	27.4	10.1	0.4	7.2	3.5	0.0
Waukee	72.2	5.6	22.2	0.0	0.0	0.0	0.0
Waukon	95.2	0.0	4.8	0.0	0.0	0.0	0.0
Waverly	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Wbstr Ct	56.7	3.0	37.3	0.0	3.0	0.0	0.0
West Union	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Winterset	92.3	0.0	7.7	0.0	0.0	0.0	0.0
Iowa	**59.4**	**14.1**	**19.4**	**0.5**	**4.0**	**2.5**	**0.0**
National (Prior Year)	**54.6**	**25.3**	**12.2**	**1.2**	**2.5**	**3.9**	**0.3**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Ackley	0.0	0.0	0.0	66.7	33.3	0.0	0.0
Adel	0.0	0.0	7.7	69.2	23.1	0.0	0.0
Albia	0.0	11.8	11.8	58.8	17.6	0.0	0.0
Algona	0.0	0.0	2.8	61.1	33.3	2.8	0.0
Allen WHC	0.0	2.0	8.1	59.6	29.3	1.0	0.0
Allison	0.0	0.0	0.0	63.6	36.4	0.0	0.0
Altoona	0.0	0.0	0.0	51.5	39.4	9.1	0.0
Ames	0.0	1.1	3.8	55.2	39.3	0.5	0.0
Anamosa	0.0	0.0	8.0	52.0	36.0	4.0	0.0
Ankeny	0.0	0.0	0.0	62.7	35.3	2.0	0.0
Atlantic	0.0	3.0	3.0	66.7	27.3	0.0	0.0
Audubon	0.0	0.0	0.0	58.3	41.7	0.0	0.0
Avoca	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Bedford	0.0	0.0	0.0	87.5	12.5	0.0	0.0
Belle Plaine	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Bloomfield	0.0	0.0	0.0	71.4	14.3	14.3	0.0
Boone	0.0	0.0	11.5	61.5	26.9	0.0	0.0
Brdlwns RP	0.2	2.8	5.8	56.8	31.2	3.2	0.0
Brooklyn	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Burlington	0.0	4.3	9.6	61.0	24.6	0.5	0.0
CHC	0.3	2.1	7.3	60.5	26.9	2.8	0.0
CR Urban	0.0	1.8	5.1	60.5	29.4	3.1	0.2
Carlisle	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Carroll	0.0	1.1	8.4	67.4	21.1	2.1	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cedar Falls	0.0	0.0	6.7	26.7	60.0	6.7	0.0
Centervill	0.0	0.0	8.3	75.0	12.5	4.2	0.0
Chariton	0.0	3.1	3.1	62.5	25.0	6.3	0.0
Charles City	1.7	1.7	3.4	65.5	25.9	1.7	0.0
Cherokee	0.0	0.0	8.3	75.0	16.7	0.0	0.0
Clarinda	0.0	0.0	0.0	86.4	13.6	0.0	0.0
Clarion	0.0	5.4	0.0	51.4	40.5	2.7	0.0
Clinton	0.0	2.6	12.3	62.1	20.0	3.1	0.0
Colfax	0.0	10.0	10.0	50.0	30.0	0.0	0.0
Columbus J	0.0	0.0	0.0	45.5	54.5	0.0	0.0
Corning	0.0	0.0	9.1	63.6	27.3	0.0	0.0
Corydon	0.0	0.0	0.0	70.6	29.4	0.0	0.0
Council Blfs	0.0	1.4	5.3	60.5	30.6	2.2	0.0
Cresco	0.0	5.3	5.3	47.4	42.1	0.0	0.0
Creston	0.0	3.1	15.6	50.0	29.7	1.6	0.0
Dallas Ctr	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Dayton	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Decorah	0.0	0.0	3.4	48.3	48.3	0.0	0.0
Denison	0.0	2.7	2.7	64.0	26.1	4.5	0.0
Dewitt	0.0	0.0	9.1	54.5	27.3	9.1	0.0
Dubuque	0.0	1.5	9.8	62.2	25.9	0.6	0.0
Dyersville	0.0	0.0	0.0	60.0	40.0	0.0	0.0
Eagle Grove	0.0	0.0	13.0	47.8	39.1	0.0	0.0
Edgerton	0.0	3.6	7.6	60.2	27.2	1.5	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Eldora	0.0	0.0	0.0	87.5	12.5	0.0	0.0
Elkader	0.0	0.0	5.1	69.2	23.1	2.6	0.0
Emmetsburg	0.0	0.0	8.3	66.7	25.0	0.0	0.0
Esthervill	0.0	2.0	6.0	72.0	18.0	2.0	0.0
Fairfiel	0.0	0.0	3.6	64.3	32.1	0.0	0.0
Forest City	0.0	0.0	6.4	63.8	25.5	4.3	0.0
Fort Dodge	0.0	1.9	8.5	66.5	21.2	1.9	0.0
Ft Madison	0.0	1.9	5.6	57.4	35.2	0.0	0.0
Garner	0.0	0.0	9.5	61.9	28.6	0.0	0.0
Glenwood	0.0	0.0	18.8	75.0	6.3	0.0	0.0
Grand Junction	0.0	0.0	100.0	0.0	0.0	0.0	0.0
Greenfield	0.0	0.0	10.5	57.9	31.6	0.0	0.0
Grinnell	0.0	0.0	10.5	78.9	10.5	0.0	0.0
Grundy Ctr	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Guthrie Ct	0.0	0.0	7.1	42.9	50.0	0.0	0.0
Hampton	0.0	3.8	1.9	52.8	37.7	3.8	0.0
Harlan	0.0	2.4	4.9	68.3	22.0	2.4	0.0
Hawarden	0.0	5.6	0.0	44.4	50.0	0.0	0.0
Hispanic Cln	0.0	0.0	0.0	33.3	66.7	0.0	0.0
Humboldt	0.0	5.0	0.0	55.0	40.0	0.0	0.0
ILH	0.2	2.8	7.7	57.5	28.3	3.3	0.2
Ida Grove	0.0	0.0	4.3	78.3	17.4	0.0	0.0
Independen	0.0	3.2	0.0	64.5	29.0	3.2	0.0
Indianola	0.0	3.6	10.7	55.4	28.6	1.8	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Iowa City	0.0	0.0	3.6	51.4	40.0	5.0	0.0
Iowa Falls	0.0	0.0	5.9	64.7	29.4	0.0	0.0
Jamaica	0.0	0.0	0.0	66.7	33.3	0.0	0.0
Jefferson	0.0	0.0	0.0	53.3	46.7	0.0	0.0
KR	0.0	0.0	0.0	42.9	57.1	0.0	0.0
Keokuk	0.0	2.2	13.3	66.7	17.8	0.0	0.0
Keosauqua	0.0	0.0	7.1	64.3	21.4	0.0	7.1
Knoxville	0.0	1.1	5.7	63.6	23.9	5.7	0.0
LCPH	0.0	0.0	3.0	71.7	23.2	2.0	0.0
Lamoni	0.0	0.0	0.0	87.5	12.5	0.0	0.0
Le Mars	0.0	2.4	4.8	54.8	33.3	4.8	0.0
Lenox	0.0	0.0	0.0	75.0	16.7	8.3	0.0
Leon	0.0	0.0	10.5	68.4	5.3	15.8	0.0
Logan	0.0	4.8	0.0	66.7	23.8	4.8	0.0
Manchester	0.0	0.0	0.0	90.9	9.1	0.0	0.0
Maquoketa	0.0	3.4	3.4	75.9	17.2	0.0	0.0
Marengo	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Marion	0.6	1.2	3.6	62.1	32.0	0.6	0.0
Marshalltown	0.0	2.0	7.6	59.5	28.9	2.0	0.0
Mason City	0.0	1.6	5.7	61.4	30.1	1.2	0.0
Monticello	0.0	0.0	10.0	60.0	10.0	20.0	0.0
Mt Ayr	0.0	16.7	0.0	58.3	25.0	0.0	0.0
Mt Pleasan	0.0	0.0	8.7	69.6	19.6	2.2	0.0
Mtown Dtown	0.0	0.0	0.0	0.0	100.0	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Muscatine	0.0	3.4	4.3	67.1	23.2	1.9	0.0
Nevada	0.0	0.0	0.0	16.7	83.3	0.0	0.0
New Hampto	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Newton	0.0	1.1	6.9	66.7	21.8	2.3	1.1
Northwood	0.0	0.0	0.0	83.3	16.7	0.0	0.0
Norwalk	0.0	0.0	0.0	50.0	50.0	0.0	0.0
Oakridge	0.0	0.0	0.0	75.0	25.0	0.0	0.0
Oelwein	0.0	0.0	0.0	56.5	43.5	0.0	0.0
Onawa	4.5	0.0	0.0	63.6	31.8	0.0	0.0
Orange Cty	0.0	0.0	4.5	81.8	13.6	0.0	0.0
Osage	0.0	3.6	3.6	57.1	35.7	0.0	0.0
Osceola	0.0	1.7	10.0	43.3	43.3	1.7	0.0
Oskaloosa	0.0	1.9	0.0	63.5	30.8	3.8	0.0
Ottumwa	0.0	2.7	7.7	62.3	24.4	2.9	0.0
Pella	0.0	0.0	0.0	52.4	38.1	9.5	0.0
Perry	1.6	1.6	6.6	57.4	29.5	3.3	0.0
Pocahontas	0.0	0.0	0.0	71.4	28.6	0.0	0.0
Postville	2.5	2.5	7.5	47.5	37.5	2.5	0.0
Red Oak	0.0	0.0	6.1	63.6	30.3	0.0	0.0
Redfield	0.0	0.0	40.0	40.0	20.0	0.0	0.0
Refugee Cln	0.0	0.0	0.0	30.0	60.0	10.0	0.0
Rock Rapid	0.0	0.0	0.0	53.8	38.5	7.7	0.0
Rock Valle	0.0	4.8	7.1	57.1	28.6	2.4	0.0
Rockwell C	0.0	0.0	0.0	90.9	9.1	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
SDM	0.0	1.1	6.4	57.4	32.4	2.7	0.0
Sac City	0.0	7.7	0.0	61.5	30.8	0.0	0.0
Sheldon	0.0	2.7	2.7	64.9	29.7	0.0	0.0
Shenandoah	0.0	6.3	9.4	62.5	18.8	3.1	0.0
Sibley	0.0	0.0	0.0	72.7	27.3	0.0	0.0
Sidney	0.0	0.0	0.0	55.6	44.4	0.0	0.0
Sigourney	0.0	0.0	0.0	45.5	54.5	0.0	0.0
Sioux Cent	0.0	6.1	12.2	55.1	24.5	2.0	0.0
Sioux City	0.0	1.9	6.4	59.9	29.8	1.9	0.0
Spencer	0.0	0.0	9.1	72.7	18.2	0.0	0.0
Spirit Lak	0.0	0.0	2.8	66.7	25.0	5.6	0.0
Storm Lake	0.0	3.2	4.0	58.1	28.2	6.5	0.0
Stuart	0.0	0.0	0.0	66.7	33.3	0.0	0.0
Tama	0.0	2.0	4.1	73.5	18.4	2.0	0.0
Tama/Settl	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Tipton	0.0	0.0	0.0	77.8	22.2	0.0	0.0
Traer	0.0	0.0	0.0	100.0	0.0	0.0	0.0
UIHC	0.0	0.0	25.0	25.0	50.0	0.0	0.0
Vinton	0.0	0.0	3.4	72.4	20.7	3.4	0.0
WDM	0.0	0.0	2.8	54.2	38.9	4.2	0.0
Washington	0.0	0.0	2.9	64.7	32.4	0.0	0.0
Waterloo	0.1	1.5	6.0	59.4	30.3	2.5	0.1
Waukee	0.0	5.6	5.6	55.6	30.6	2.8	0.0
Waukon	0.0	0.0	0.0	76.2	23.8	0.0	0.0

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Waverly	0.0	0.0	0.0	61.9	28.6	9.5	0.0
Wbstr Ct	0.0	1.5	7.5	61.2	29.9	0.0	0.0
West Union	0.0	0.0	0.0	40.0	53.3	6.7	0.0
Winterset	0.0	0.0	7.7	65.4	26.9	0.0	0.0
Iowa	**0.1**	**1.9**	**6.1**	**60.2**	**29.2**	**2.4**	**0.0**
National (Prior Year)	**0.1**	**2.9**	**8.8**	**61.5**	**24.9**	**1.8**	**0.1**

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)		Weight Gain (5)		Anemia (Low Hb/Hct) (6)		Gestational Diabetes During Pregnancy	Hypertension During Pregnancy
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum		
	Number % (Rank)	% (Rank)	Number % (Rank)	% (Rank)	Number % (Rank)	Number % (Rank)	Number % (Rank)	Number % (Rank)
Ackley	0.0 (1)	33.3 (5)	33.3 (116)	66.7 (136)	100.0 (141)	100.0 (141)	0.0 (1)	0.0 (1)
Adel	0.0 (1)	88.9 (135)	11.1 (19)	55.6 (117)	0.0 (1)	0.0 (1)	0.0 (1)	0.0 (1)
Albia	25.0 (148)	41.7 (9)	33.3 (116)	33.3 (31)	0.0 (1)	0.0 (1)	0.0 (1)	0.0 (1)
Algona	0.0 (1)	71.4 (102)	15.4 (30)	50.0 (99)	25.0 (117)	0.0 (1)	100.0 (49)	100.0 (67)
Allen WHC	3.3 (87)	62.0 (61)	26.7 (91)	40.7 (53)	0.0 (1)	0.0 (1)	100.0 (49)	100.0 (67)
Allison	0.0 (1)	87.5 (132)	25.0 (76)	56.3 (118)	0.0 (1)	0.0 (1)	100.0 (49)	100.0 (67)
Altoona	4.0 (98)	72.0 (107)	45.8 (143)	33.3 (31)	0.0 (1)	0.0 (1)	100.0 (49)	100.0 (67)
Ames	4.5 (106)	61.9 (59)	27.6 (96)	41.7 (58)	0.0 (1)	4.8 (97)	100.0 (49)	0.0 (1)
Anamosa	4.5 (106)	68.2 (91)	23.8 (69)	52.4 (112)	33.3 (123)	0.0 (1)	100.0 (49)	100.0 (67)
Ankeny	6.5 (128)	58.1 (36)	17.2 (34)	51.7 (111)	10.0 (103)	0.0 (1)	100.0 (49)	100.0 (67)
Atlantic	0.0 (1)	66.7 (78)	23.1 (67)	65.4 (134)	0.0 (1)	25.0 (117)	0.0 (1)	100.0 (67)
Audubon	0.0 (1)	71.4 (102)	14.3 (25)	14.3 (10)	0.0 (1)	100.0 (141)	100.0 (49)	100.0 (67)
Avoca	0.0 (1)	100.0 (138)	0.0 (1)	100.0 (145)	0.0 (1)	0.0 (1)	0.0 (1)	0.0 (1)
Bedford	16.7 (144)	33.3 (5)	33.3 (116)	16.7 (11)	0.0 (1)	0.0 (1)	0.0 (1)	0.0 (1)
Belle Plaine	0.0 (1)	85.7 (131)	28.6 (102)	42.9 (60)	0.0 (1)	0.0 (1)	100.0 (49)	100.0 (67)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Bloomfield		0.0 (1)	57.1 (33)		42.9 (139)	42.9 (60)		100.0 (141)		0.0 (1)		0.0 (1)		0.0 (1)
Boone		0.0 (1)	66.7 (78)		25.0 (76)	45.0 (79)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Brdlwns RP		5.6 (115)	53.1 (21)		23.8 (69)	43.7 (66)		3.0 (95)		0.0 (1)		100.0 (49)		100.0 (67)
Brooklyn		0.0 (1)	100.0 (138)		50.0 (144)	50.0 (99)		0.0 (1)		100.0 (141)		0.0 (1)		0.0 (1)
Burlington		5.0 (111)	59.0 (41)		15.6 (32)	50.4 (109)		5.6 (98)		7.7 (100)		100.0 (49)		100.0 (67)
CHC		6.3 (121)	54.8 (29)		23.5 (68)	47.5 (94)		38.2 (128)		50.0 (132)		100.0 (49)		100.0 (67)
CR Urban		4.1 (102)	59.8 (46)		22.4 (63)	48.7 (97)		40.2 (131)		22.6 (116)		100.0 (49)		100.0 (67)
Carlisle		0.0 (1)	66.7 (78)		66.7 (147)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (67)
Carroll		7.9 (133)	60.5 (55)		18.7 (37)	49.3 (98)		12.0 (107)		41.7 (131)		100.0 (49)		100.0 (67)
Cedar Falls		20.0 (146)	70.0 (99)		0.0 (1)	70.0 (143)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (67)
Centervill		0.0 (1)	75.0 (114)		29.2 (106)	62.5 (129)		60.0 (138)		0.0 (1)		0.0 (1)		100.0 (67)
Chariton		0.0 (1)	69.2 (97)		29.2 (106)	45.8 (86)		40.0 (129)		25.0 (117)		100.0 (49)		100.0 (67)
Charles City		4.7 (108)	60.5 (55)		30.2 (109)	39.5 (43)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Cherokee		8.3 (136)	70.8 (101)		25.0 (76)	45.8 (86)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Clarinda		0.0 (1)	81.3 (127)		25.0 (76)	43.8 (67)		0.0 (1)		40.0 (129)		0.0 (1)		100.0 (67)
Clarion		3.4 (89)	75.9 (120)		37.9 (134)	31.0 (28)		20.0 (113)		0.0 (1)		100.0 (49)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Clinton		5.9 (117)	66.0 (77)		22.8 (66)	57.0 (120)		19.2 (112)		16.7 (111)		100.0 (49)		100.0 (67)
Colfax		14.3 (142)	71.4 (102)		14.3 (25)	28.6 (21)		0.0 (1)		100.0 (141)		100.0 (49)		0.0 (1)
Columbus J		0.0 (1)	66.7 (78)		11.1 (19)	44.4 (71)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Corning		0.0 (1)	42.9 (10)		28.6 (102)	28.6 (21)		100.0 (141)		33.3 (123)		100.0 (49)		0.0 (1)
Corydon		0.0 (1)	100.0 (138)		27.3 (93)	63.6 (132)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (67)
Council Blfs		3.8 (94)	63.6 (66)		24.4 (74)	50.5 (110)		10.2 (104)		11.1 (102)		100.0 (49)		100.0 (67)
Cresco		6.3 (121)	75.0 (114)		37.5 (131)	37.5 (36)		50.0 (135)		33.3 (123)		100.0 (49)		100.0 (67)
Creston		4.9 (110)	56.1 (30)		25.6 (86)	46.2 (90)		33.3 (123)		0.0 (1)		100.0 (49)		100.0 (67)
Dallas Ctr		0.0 (1)	100.0 (138)		0.0 (1)	100.0 (145)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Dayton		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Decorah		0.0 (1)	72.7 (110)		4.5 (16)	45.5 (83)		0.0 (1)		50.0 (132)		100.0 (49)		100.0 (67)
Denison		3.3 (87)	64.1 (70)		33.0 (115)	29.5 (26)		3.4 (97)		0.0 (1)		100.0 (49)		100.0 (67)
Dewitt		0.0 (1)	77.8 (124)		33.3 (116)	44.4 (71)		100.0 (141)		0.0 (1)		0.0 (1)		100.0 (67)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Dubuque		2.6 (81)	63.2 (65)		24.0 (71)	47.2 (93)		50.0 (135)		36.0 (128)		100.0 (49)		100.0 (67)
Dyersville		0.0 (1)	100.0 (138)		33.3 (116)	66.7 (136)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Eagle Grove		0.0 (1)	53.3 (22)		20.0 (42)	40.0 (46)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (67)
Edgerton		3.7 (92)	61.7 (57)		21.4 (57)	52.7 (113)		23.3 (116)		17.4 (113)		100.0 (49)		100.0 (67)
Eldora		0.0 (1)	40.0 (8)		0.0 (1)	40.0 (46)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Elkader		0.0 (1)	69.0 (96)		19.2 (40)	53.8 (114)		14.3 (109)		25.0 (117)		100.0 (49)		100.0 (67)
Emmetsburg		0.0 (1)	50.0 (13)		20.0 (42)	60.0 (124)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Esthervill		2.9 (83)	60.0 (47)		30.3 (110)	39.4 (42)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Fairfiel		0.0 (1)	63.6 (66)		14.3 (25)	57.1 (121)		0.0 (1)		40.0 (129)		0.0 (1)		0.0 (1)
Forest City		8.1 (135)	59.5 (44)		19.4 (41)	44.4 (71)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Fort Dodge		0.6 (74)	68.5 (93)		19.1 (39)	50.0 (99)		2.6 (94)		0.0 (1)		100.0 (49)		100.0 (67)
Ft Madison		0.0 (1)	76.3 (123)		14.3 (25)	54.3 (115)		8.3 (100)		0.0 (1)		100.0 (49)		100.0 (67)
Garner		6.3 (121)	68.8 (94)		25.0 (76)	50.0 (99)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Glenwood		7.7 (130)	53.8 (24)		30.8 (112)	23.1 (13)		0.0 (1)		25.0 (117)		0.0 (1)		0.0 (1)
Grand Junction		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Greenfield		6.3 (121)	81.3 (127)		20.0 (42)	40.0 (46)		100.0 (141)		0.0 (1)		100.0 (49)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Grinnell		5.9 (117)	52.9 (20)		13.3 (23)	66.7 (136)		0.0 (1)		66.7 (140)		0.0 (1)		0.0 (1)
Grundy Ctr		0.0 (1)	100.0 (138)		25.0 (76)	50.0 (99)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Guthrie Ct		0.0 (1)	87.5 (132)		25.0 (76)	62.5 (129)		100.0 (141)		0.0 (1)		100.0 (49)		0.0 (1)
Hampton		2.4 (79)	75.6 (119)		34.2 (124)	39.5 (43)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Harlan		6.9 (129)	72.4 (109)		17.2 (34)	65.5 (135)		0.0 (1)		12.5 (106)		100.0 (49)		100.0 (67)
Hawarden		6.3 (121)	75.0 (114)		26.7 (91)	46.7 (91)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Hispanic Cln		0.0 (1)	66.7 (78)		50.0 (144)	50.0 (99)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Humboldt		0.0 (1)	66.7 (78)		44.4 (142)	27.8 (20)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
ILH		4.8 (109)	61.9 (59)		27.8 (99)	38.6 (40)		1.4 (92)		3.8 (96)		100.0 (49)		100.0 (67)
Ida Grove		5.0 (111)	65.0 (72)		35.0 (125)	45.0 (79)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Independen		0.0 (1)	80.0 (125)		36.8 (129)	26.3 (17)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Indianola		7.9 (133)	50.0 (13)		31.6 (114)	39.5 (43)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Iowa City		3.4 (89)	61.7 (57)		20.8 (53)	43.5 (64)		18.5 (111)		9.4 (101)		100.0 (49)		100.0 (67)
Iowa Falls		7.7 (130)	69.2 (97)		15.4 (30)	38.5 (39)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Jamaica		0.0 (1)	100.0 (138)		0.0 (1)	100.0 (145)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Jefferson		7.7 (130)	84.6 (130)		30.8 (112)	61.5 (128)		40.0 (129)		0.0 (1)		0.0 (1)		100.0 (67)
KR		0.0 (1)	63.6 (66)		27.3 (93)	36.4 (34)		0.0 (1)		33.3 (123)		100.0 (49)		100.0 (67)
Keokuk		12.5 (141)	59.4 (43)		27.6 (96)	31.0 (28)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Keosauqua		0.0 (1)	66.7 (78)		41.7 (137)	41.7 (58)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Knoxville		1.6 (75)	65.6 (74)		20.3 (51)	40.7 (53)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
LCPH		0.0 (1)	58.8 (40)		22.4 (63)	55.2 (116)		58.3 (137)		33.3 (123)		100.0 (49)		100.0 (67)
Lamoni		0.0 (1)	75.0 (114)		25.0 (76)	50.0 (99)		0.0 (1)		50.0 (132)		100.0 (49)		0.0 (1)
Le Mars		3.1 (84)	71.9 (106)		16.1 (33)	45.2 (81)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Lenox		0.0 (1)	62.5 (62)		37.5 (131)	37.5 (36)		0.0 (1)		100.0 (141)		100.0 (49)		0.0 (1)
Leon		0.0 (1)	66.7 (78)		63.6 (146)	27.3 (19)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Logan		0.0 (1)	60.0 (47)		14.3 (25)	50.0 (99)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Manchester		0.0 (1)	88.9 (135)		22.2 (59)	44.4 (71)		0.0 (1)		100.0 (141)		0.0 (1)		0.0 (1)
Maquoketa		8.7 (138)	65.2 (73)		13.0 (22)	56.5 (119)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Marengo		20.0 (146)	80.0 (125)		40.0 (135)	60.0 (124)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Marion		2.3 (78)	65.9 (76)		20.6 (52)	46.0 (89)		37.0 (127)		14.3 (109)		100.0 (49)		100.0 (67)
Marshalltown		4.3 (103)	59.1 (42)		27.6 (96)	36.8 (35)		12.5 (108)		18.2 (114)		100.0 (49)		100.0 (67)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Mason City		3.1 (84)	63.0 (64)		20.0 (42)	44.7 (76)		0.0 (1)		6.3 (98)		100.0 (49)		100.0 (67)
Monticello		0.0 (1)	44.4 (12)		22.2 (59)	66.7 (136)		66.7 (140)		0.0 (1)		100.0 (49)		100.0 (67)
Mt Ayr		0.0 (1)	60.0 (47)		30.0 (108)	40.0 (46)		0.0 (1)		33.3 (123)		100.0 (49)		0.0 (1)
Mt Pleasan		2.8 (82)	66.7 (78)		20.0 (42)	42.9 (60)		9.1 (102)		0.0 (1)		100.0 (49)		100.0 (67)
Mtown Dtown		0.0 (1)	0.0 (1)		100.0 (148)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Muscatine		1.9 (76)	57.9 (34)		25.6 (86)	44.2 (70)		3.1 (96)		14.3 (109)		100.0 (49)		100.0 (67)
Nevada		0.0 (1)	100.0 (138)		0.0 (1)	66.7 (136)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
New Hampto		5.6 (115)	72.2 (108)		22.2 (59)	38.9 (41)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Newton		6.3 (121)	58.7 (39)		22.4 (63)	44.8 (78)		28.6 (122)		0.0 (1)		100.0 (49)		100.0 (67)
Northwood		0.0 (1)	87.5 (132)		0.0 (1)	57.1 (121)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Norwalk		0.0 (1)	60.0 (47)		20.0 (42)	40.0 (46)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Oakridge		0.0 (1)	50.0 (13)		40.0 (135)	60.0 (124)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Oelwein		0.0 (1)	81.8 (129)		19.0 (38)	47.6 (95)		44.4 (133)		0.0 (1)		100.0 (49)		100.0 (67)
Onawa		6.3 (121)	56.3 (31)		13.3 (23)	40.0 (46)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Orange Cty		0.0 (1)	60.0 (47)		20.0 (42)	20.0 (12)		33.3 (123)		0.0 (1)		100.0 (49)		0.0 (1)
Osage		0.0 (1)	68.2 (91)		18.2 (36)	59.1 (123)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Osceola		4.3 (103)	63.8 (69)		33.3 (116)	37.8 (38)		42.9 (132)		50.0 (132)		100.0 (49)		100.0 (67)
Oskaloosa		8.3 (136)	58.3 (37)		21.9 (58)	43.8 (67)		20.0 (113)		50.0 (132)		0.0 (1)		100.0 (67)
Ottumwa		4.3 (103)	56.3 (31)		28.8 (105)	41.0 (56)		26.4 (121)		12.5 (106)		100.0 (49)		100.0 (67)
Pella		0.0 (1)	53.3 (22)		20.0 (42)	46.7 (91)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Perry		6.0 (119)	54.0 (27)		28.6 (102)	32.7 (30)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Pocahontas		16.7 (144)	50.0 (13)		0.0 (1)	0.0 (1)		100.0 (141)		0.0 (1)		0.0 (1)		0.0 (1)
Postville		0.0 (1)	50.0 (13)		41.9 (138)	29.0 (24)		44.4 (133)		50.0 (132)		100.0 (49)		0.0 (1)
Red Oak		4.0 (98)	76.0 (121)		24.0 (71)	40.0 (46)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Redfield		0.0 (1)	75.0 (114)		25.0 (76)	25.0 (14)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Refugee Cln		0.0 (1)	62.5 (62)		12.5 (21)	12.5 (9)		100.0 (141)		100.0 (141)		100.0 (49)		0.0 (1)
Rock Rapid		0.0 (1)	68.8 (94)		20.0 (42)	60.0 (124)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Rock Valle		3.1 (84)	50.0 (13)		35.5 (127)	25.8 (16)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Rockwell C		0.0 (1)	33.3 (5)		37.5 (131)	50.0 (99)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (67)
SDM		5.3 (114)	59.5 (44)		24.7 (75)	44.5 (75)		1.5 (93)		0.0 (1)		100.0 (49)		100.0 (67)
Sac City		9.1 (139)	72.7 (110)		36.4 (128)	45.5 (83)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Sheldon		3.6 (91)	67.9 (89)		28.0 (101)	48.0 (96)		0.0 (1)		11.1 (102)		100.0 (49)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Shenandoah		3.8 (94)	73.1 (112)		20.8 (53)	50.0 (99)		20.0 (113)		25.0 (117)		100.0 (49)		100.0 (67)
Sibley		0.0 (1)	100.0 (138)		0.0 (1)	25.0 (14)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Sidney		0.0 (1)	60.0 (47)		25.0 (76)	75.0 (144)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Sigourney		0.0 (1)	54.5 (28)		27.3 (93)	9.1 (8)		25.0 (117)		0.0 (1)		0.0 (1)		0.0 (1)
Sioux Cent		2.5 (80)	60.0 (47)		33.3 (116)	30.8 (27)		11.1 (106)		0.0 (1)		100.0 (49)		0.0 (1)
Sioux City		4.0 (98)	60.4 (54)		24.1 (73)	40.9 (55)		10.4 (105)		11.4 (105)		100.0 (49)		100.0 (67)
Spencer		2.0 (77)	68.0 (90)		30.4 (111)	41.3 (57)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Spirit Lak		4.0 (98)	76.0 (121)		20.8 (53)	45.8 (86)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Storm Lake		3.9 (97)	65.8 (75)		27.9 (100)	44.1 (69)		0.0 (1)		16.7 (111)		100.0 (49)		100.0 (67)
Stuart		0.0 (1)	71.4 (102)		42.9 (139)	42.9 (60)		0.0 (1)		100.0 (141)		0.0 (1)		0.0 (1)
Tama		5.0 (111)	70.0 (99)		21.1 (56)	44.7 (76)		25.0 (117)		25.0 (117)		100.0 (49)		100.0 (67)
Tama/Settl		0.0 (1)	100.0 (138)		0.0 (1)	100.0 (145)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Tipton		14.3 (142)	42.9 (10)		42.9 (139)	28.6 (21)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (67)
Traer		0.0 (1)	100.0 (138)		33.3 (116)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (67)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
UIHC		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		100.0 (49)		0.0 (1)
Vinton		0.0 (1)	64.3 (71)		7.1 (17)	67.9 (141)		33.3 (123)		0.0 (1)		0.0 (1)		100.0 (67)
WDM		6.1 (120)	67.3 (88)		26.1 (88)	43.5 (64)		8.3 (100)		11.1 (102)		100.0 (49)		100.0 (67)
Washington		11.5 (140)	53.8 (24)		26.1 (88)	34.8 (33)		14.3 (109)		12.5 (106)		100.0 (49)		100.0 (67)
Waterloo		3.7 (92)	58.0 (35)		22.2 (59)	45.5 (83)		7.8 (99)		7.0 (99)		100.0 (49)		100.0 (67)
Waukee		3.8 (94)	53.8 (24)		4.2 (15)	62.5 (129)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Waukon		0.0 (1)	66.7 (78)		35.3 (126)	29.4 (25)		60.0 (138)		0.0 (1)		0.0 (1)		100.0 (67)
Waverly		0.0 (1)	89.5 (137)		10.5 (18)	68.4 (142)		0.0 (1)		0.0 (1)		100.0 (49)		100.0 (67)
Wbstr Ct		0.0 (1)	52.8 (19)		26.4 (90)	45.3 (82)		0.0 (1)		50.0 (132)		100.0 (49)		100.0 (67)
West Union		0.0 (1)	58.3 (37)		0.0 (1)	63.6 (132)		0.0 (1)		50.0 (132)		100.0 (49)		0.0 (1)
Winterset		0.0 (1)	73.7 (113)		36.8 (129)	26.3 (17)		25.0 (117)		20.0 (115)		100.0 (49)		100.0 (67)
Iowa		**4.0**	**61.2**		**24.1**	**44.7**		**13.9**		**13.1**		**100.0**		**100.0**
National (Prior Year)	**128,555**	**4.0**	**61.6**	**89,471**	**22.3**	**49.7**	**34,531**	**19.0**	**48,409**	**17.0**	**94,843**	**9.2**	**94,594**	**9.6**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley		33.3 (54)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Adel		12.5 (131)	0.0 (1)		0.0 (34)		20.0 (70)		100.0 (1)
Albia		7.7 (137)	23.1 (133)		0.0 (34)		28.6 (44)		85.7 (119)
Algona		65.2 (6)	8.7 (73)		0.0 (34)		6.3 (117)		95.2 (63)
Allen WHC		21.9 (107)	28.1 (139)		0.0 (34)		27.8 (48)		91.9 (88)
Allison		33.3 (54)	0.0 (1)		0.0 (34)		33.3 (34)		93.3 (78)
Altoona		31.8 (65)	13.6 (101)		0.0 (34)		14.3 (97)		94.1 (68)
Ames		42.9 (19)	6.5 (59)		2.0 (19)		20.6 (69)		94.1 (68)
Anamosa		35.0 (50)	5.0 (49)		0.0 (34)		33.3 (34)		93.3 (78)
Ankeny		28.6 (79)	10.7 (85)		0.0 (34)		18.2 (82)		93.3 (78)
Atlantic		25.0 (94)	8.3 (67)		0.0 (34)		12.5 (103)		100.0 (1)
Audubon		37.5 (40)	12.5 (95)		0.0 (34)		50.0 (10)		100.0 (1)
Avoca		0.0 (138)	50.0 (147)		0.0 (34)		0.0 (118)		100.0 (1)
Bedford		0.0 (138)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Belle Plaine		12.5 (131)	0.0 (1)		0.0 (34)		0.0 (118)		83.3 (123)
Bloomfield		40.0 (27)	20.0 (127)		0.0 (34)		0.0 (118)		0.0 (141)
Boone		15.4 (128)	7.7 (65)		5.3 (4)		12.5 (103)		100.0 (1)
Brdlwns RP		26.7 (89)	11.1 (87)		0.5 (30)		11.5 (109)		89.8 (102)
Brooklyn		33.3 (54)	33.3 (143)		0.0 (34)		0.0 (118)		100.0 (1)
Burlington		33.8 (53)	2.2 (42)		1.7 (21)		17.6 (86)		96.2 (57)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
CHC		26.5 (90)	18.9 (126)		1.1 (24)		11.1 (110)		90.9 (93)
CR Urban		24.0 (101)	9.2 (76)		0.3 (33)		16.2 (90)		87.7 (111)
Carlisle		50.0 (10)	0.0 (1)		0.0 (34)		50.0 (10)		100.0 (1)
Carroll		27.1 (88)	11.4 (90)		1.4 (22)		27.3 (51)		95.7 (58)
Cedar Falls		33.3 (54)	8.3 (67)		0.0 (34)		71.4 (5)		90.9 (93)
Centervill		30.0 (71)	15.0 (112)		0.0 (34)		100.0 (1)		100.0 (1)
Chariton		40.0 (27)	4.0 (45)		0.0 (34)		22.2 (63)		77.8 (132)
Charles City		39.6 (33)	8.3 (67)		0.0 (34)		16.7 (87)		93.8 (74)
Cherokee		38.5 (37)	3.8 (44)		4.0 (9)		14.3 (97)		100.0 (1)
Clarinda		36.8 (44)	0.0 (1)		0.0 (34)		42.9 (23)		100.0 (1)
Clarion		30.0 (71)	20.0 (127)		0.0 (34)		14.8 (96)		86.7 (117)
Clinton		42.0 (23)	5.3 (50)		0.8 (27)		41.7 (25)		86.8 (115)
Colfax		16.7 (124)	16.7 (116)		0.0 (34)		40.0 (28)		100.0 (1)
Columbus J		22.2 (105)	11.1 (87)		0.0 (34)		0.0 (118)		62.5 (139)
Corning		28.6 (79)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Corydon		33.3 (54)	8.3 (67)		0.0 (34)		22.2 (63)		77.8 (132)
Council Blfs		41.3 (24)	5.9 (54)		1.2 (23)		23.6 (60)		86.5 (118)
Cresco		42.1 (22)	5.3 (50)		0.0 (34)		22.2 (63)		88.9 (104)
Creston		51.2 (9)	17.1 (122)		0.0 (34)		40.9 (27)		95.5 (60)
Dallas Ctr		0.0 (138)	0.0 (1)		0.0 (34)		0.0 (118)		0.0 (141)
Dayton		0.0 (138)	0.0 (1)		0.0 (34)		0.0 (118)		0.0 (141)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Decorah		20.8 (113)	4.2 (47)		0.0 (34)		19.0 (79)		90.9 (93)
Denison		17.1 (123)	7.9 (66)		0.0 (34)		22.7 (62)		96.3 (55)
Dewitt		37.5 (40)	0.0 (1)		0.0 (34)		50.0 (10)		83.3 (123)
Dubuque		38.8 (36)	12.5 (95)		5.5 (3)		21.3 (67)		90.3 (98)
Dyersville		33.3 (54)	0.0 (1)		0.0 (34)		50.0 (10)		100.0 (1)
Eagle Grove		27.8 (86)	27.8 (138)		0.0 (34)		11.8 (108)		88.9 (104)
Edgerton		31.0 (69)	11.8 (93)		1.9 (20)		0.0 (118)		75.0 (135)
Eldora		80.0 (3)	0.0 (1)		0.0 (34)		50.0 (10)		80.0 (129)
Elkader		25.8 (92)	6.5 (59)		3.8 (10)		33.3 (34)		93.9 (72)
Emmetsburg		43.8 (18)	18.8 (125)		0.0 (34)		66.7 (6)		93.3 (78)
Esthervill		22.9 (104)	11.4 (90)		0.0 (34)		41.7 (25)		91.7 (89)
Fairfiel		35.0 (50)	30.0 (141)		4.5 (5)		100.0 (1)		100.0 (1)
Forest City		38.9 (34)	8.3 (67)		0.0 (34)		18.2 (82)		83.8 (122)
Fort Dodge		23.9 (103)	17.0 (121)		3.2 (13)		17.8 (85)		93.4 (76)
Ft Madison		30.0 (71)	2.5 (43)		0.0 (34)		23.5 (61)		95.1 (64)
Garner		46.7 (17)	20.0 (127)		0.0 (34)		40.0 (28)		85.7 (119)
Glenwood		10.0 (136)	10.0 (81)		0.0 (34)		0.0 (118)		66.7 (137)
Grand Junction		100.0 (1)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Greenfield		21.4 (109)	7.1 (62)		0.0 (34)		22.2 (63)		87.5 (113)
Grinnell		14.3 (129)	7.1 (62)		0.0 (34)		0.0 (118)		83.3 (123)
Grundy Ctr		0.0 (138)	16.7 (116)		0.0 (34)		50.0 (10)		100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Guthrie Ct		28.6 (79)	14.3 (105)		0.0 (34)		0.0 (118)		0.0 (141)
Hampton		30.6 (70)	30.6 (142)		3.3 (12)		30.3 (41)		86.8 (115)
Harlan		25.0 (94)	0.0 (1)		0.0 (34)		10.0 (113)		100.0 (1)
Hawarden		37.5 (40)	12.5 (95)		0.0 (34)		30.0 (43)		91.7 (89)
Hispanic Cln		16.7 (124)	0.0 (1)		0.0 (34)		100.0 (1)		100.0 (1)
Humboldt		33.3 (54)	26.7 (135)		0.0 (34)		12.5 (103)		93.8 (74)
ILH		32.0 (64)	11.6 (92)		0.7 (29)		20.9 (68)		87.7 (111)
Ida Grove		38.9 (34)	0.0 (1)		0.0 (34)		20.0 (70)		88.2 (109)
Independen		40.0 (27)	20.0 (127)		0.0 (34)		28.6 (44)		100.0 (1)
Indianola		19.4 (119)	22.2 (132)		0.0 (34)		15.6 (92)		94.1 (68)
Iowa City		22.0 (106)	18.5 (124)		1.0 (26)		19.7 (76)		88.1 (110)
Iowa Falls		36.4 (45)	0.0 (1)		0.0 (34)		11.1 (110)		88.9 (104)
Jamaica		50.0 (10)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Jefferson		33.3 (54)	16.7 (116)		0.0 (34)		20.0 (70)		100.0 (1)
KR		16.7 (124)	33.3 (143)		0.0 (34)		40.0 (28)		100.0 (1)
Keokuk		20.7 (114)	0.0 (1)		7.7 (2)		27.3 (51)		93.1 (84)
Keosauqua		25.0 (94)	0.0 (1)		0.0 (34)		50.0 (10)		100.0 (1)
Knoxville		35.7 (49)	10.7 (85)		0.0 (34)		23.7 (59)		100.0 (1)
LCPH		37.7 (39)	6.6 (61)		0.0 (34)		13.3 (100)		94.0 (71)
Lamoni		28.6 (79)	28.6 (140)		0.0 (34)		0.0 (118)		100.0 (1)
Le Mars		16.1 (127)	9.7 (79)		0.0 (34)		46.2 (20)		92.9 (85)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Lenox		55.6 (8)	11.1 (87)		0.0 (34)		50.0 (10)		66.7 (137)
Leon		36.4 (45)	27.3 (137)		0.0 (34)		28.6 (44)		100.0 (1)
Logan		28.6 (79)	0.0 (1)		0.0 (34)		66.7 (6)		100.0 (1)
Manchester		40.0 (27)	10.0 (81)		0.0 (34)		11.1 (110)		100.0 (1)
Maquoketa		36.4 (45)	9.1 (75)		0.0 (34)		20.0 (70)		89.5 (103)
Marengo		0.0 (138)	0.0 (1)		0.0 (34)		0.0 (118)		0.0 (141)
Marion		29.0 (78)	13.8 (103)		0.8 (27)		19.6 (77)		93.3 (78)
Marshalltown		35.0 (50)	5.9 (54)		1.1 (24)		16.1 (91)		92.0 (87)
Mason City		40.7 (25)	13.7 (102)		0.0 (34)		24.7 (58)		88.3 (108)
Monticello		42.9 (19)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Mt Ayr		50.0 (10)	0.0 (1)		0.0 (34)		25.0 (54)		100.0 (1)
Mt Pleasan		20.6 (115)	8.8 (74)		0.0 (34)		40.0 (28)		90.6 (97)
Mtown Dtown		0.0 (138)	0.0 (1)		0.0 (34)		0.0 (118)		0.0 (141)
Muscatine		25.5 (93)	9.3 (77)		0.0 (34)		18.3 (81)		93.4 (76)
Nevada		25.0 (94)	25.0 (134)		0.0 (34)		33.3 (34)		100.0 (1)
New Hampto		25.0 (94)	16.7 (116)		0.0 (34)		57.1 (8)		100.0 (1)
Newton		24.0 (101)	14.0 (104)		0.0 (34)		11.9 (107)		95.3 (62)
Northwood		66.7 (4)	0.0 (1)		0.0 (34)		12.5 (103)		88.9 (104)
Norwalk		50.0 (10)	0.0 (1)		0.0 (34)		20.0 (70)		80.0 (129)
Oakridge		40.0 (27)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Oelwein		26.3 (91)	10.5 (84)		0.0 (34)		27.3 (51)		95.5 (60)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Onawa		14.3 (129)	14.3 (105)		0.0 (34)		0.0 (118)		100.0 (1)
Orange Cty		28.6 (79)	14.3 (105)		0.0 (34)		40.0 (28)		80.0 (129)
Osage		21.7 (108)	21.7 (131)		0.0 (34)		43.8 (22)		87.5 (113)
Osceola		27.7 (87)	8.5 (72)		2.4 (16)		31.3 (40)		93.9 (72)
Oskaloosa		40.6 (26)	9.4 (78)		3.2 (13)		0.0 (118)		0.0 (141)
Ottumwa		29.7 (74)	27.2 (136)		2.4 (16)		28.6 (44)		42.9 (140)
Pella		21.4 (109)	14.3 (105)		0.0 (34)		18.2 (82)		100.0 (1)
Perry		18.8 (122)	16.7 (116)		2.2 (18)		10.0 (113)		93.3 (78)
Pocahontas		66.7 (4)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Postville		19.4 (119)	12.9 (100)		0.0 (34)		19.4 (78)		90.3 (98)
Red Oak		38.1 (38)	4.8 (48)		4.2 (7)		16.7 (87)		83.3 (123)
Redfield		50.0 (10)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Refugee Cln		0.0 (138)	12.5 (95)		0.0 (34)		0.0 (118)		71.4 (136)
Rock Rapid		29.4 (75)	5.9 (54)		0.0 (34)		15.4 (94)		100.0 (1)
Rock Valle		37.5 (40)	15.6 (114)		0.0 (34)		13.3 (100)		81.3 (128)
Rockwell C		33.3 (54)	0.0 (1)		0.0 (34)		20.0 (70)		100.0 (1)
SDM		24.8 (99)	9.7 (79)		0.0 (34)		15.6 (92)		90.3 (98)
Sac City		42.9 (19)	14.3 (105)		0.0 (34)		25.0 (54)		100.0 (1)
Sheldon		20.0 (116)	4.0 (45)		0.0 (34)		44.4 (21)		96.3 (55)
Shenandoah		48.0 (15)	0.0 (1)		4.5 (5)		16.7 (87)		94.4 (65)
Sibley		20.0 (116)	0.0 (1)		25.0 (1)		50.0 (10)		83.3 (123)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Sidney		0.0 (138)	33.3 (143)		0.0 (34)		0.0 (118)		100.0 (1)
Sigourney		40.0 (27)	0.0 (1)		0.0 (34)		100.0 (1)		100.0 (1)
Sioux Cent		19.0 (121)	7.1 (62)		0.0 (34)		9.5 (115)		95.7 (58)
Sioux City		28.0 (85)	5.7 (53)		0.5 (30)		27.7 (50)		90.7 (96)
Spencer		11.9 (134)	11.9 (94)		2.8 (15)		51.1 (9)		91.5 (92)
Spirit Lak		36.0 (48)	16.0 (115)		4.2 (7)		48.0 (19)		100.0 (1)
Storm Lake		21.1 (112)	5.3 (50)		0.0 (34)		34.6 (33)		100.0 (1)
Stuart		12.5 (131)	12.5 (95)		0.0 (34)		25.0 (54)		100.0 (1)
Tama		31.4 (66)	0.0 (1)		0.0 (34)		13.8 (99)		100.0 (1)
Tama/Settl		0.0 (138)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
Tipton		20.0 (116)	60.0 (148)		0.0 (34)		0.0 (118)		100.0 (1)
Traer		100.0 (1)	0.0 (1)		0.0 (34)		0.0 (118)		100.0 (1)
UIHC		0.0 (138)	0.0 (1)		0.0 (34)		0.0 (118)		0.0 (141)
Vinton		47.6 (16)	14.3 (105)		0.0 (34)		33.3 (34)		100.0 (1)
WDM		33.3 (54)	10.3 (83)		0.0 (34)		7.7 (116)		94.4 (65)
Washington		10.7 (135)	14.3 (105)		0.0 (34)		15.0 (95)		90.0 (101)
Waterloo		29.2 (76)	17.1 (122)		0.5 (30)		30.3 (41)		92.7 (86)
Waukee		29.2 (76)	0.0 (1)		3.6 (11)		18.8 (80)		100.0 (1)
Waukon		31.3 (67)	6.3 (57)		0.0 (34)		27.8 (48)		94.4 (65)
Waverly		61.5 (7)	15.4 (113)		0.0 (34)		33.3 (34)		100.0 (1)
Wbstr Ct		24.1 (100)	35.2 (146)		0.0 (34)		13.0 (102)		76.5 (134)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Comparison of Maternal Behavioral Indicators (2,3)									
Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
West Union		21.4 (109)	0.0 (1)		0.0 (34)		42.9 (23)		85.7 (119)
Winterset		31.3 (67)	6.3 (57)		0.0 (34)		25.0 (54)		91.7 (89)
Iowa		**29.9**	**11.6**		**1.0**		**22.2**		**5109.0**
National (Prior Year)	**100,055**	**77.9**	**5.0**	**134,286**	**21.1**	**82,984**	**20.5**	**79,295**	**89.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley		33.3 (85)		33.3 (128)		0.0 (104)		0.0 (1)		0.0 (1)
Adel		25.0 (54)		33.3 (128)		0.0 (104)		0.0 (1)		0.0 (1)
Albia		30.8 (77)		30.0 (125)		25.0 (70)		7.1 (82)		0.0 (1)
Algona		37.5 (101)		25.0 (114)		22.2 (80)		4.3 (55)		0.0 (1)
Allen WHC		18.4 (36)		10.7 (53)		35.7 (45)		6.8 (78)		1.5 (98)
Allison		41.2 (114)		23.8 (110)		42.9 (38)		11.8 (112)		0.0 (1)
Altoona		27.8 (68)		5.3 (31)		100.0 (1)		5.0 (59)		0.0 (1)
Ames		21.0 (43)		7.8 (41)		19.0 (88)		8.6 (98)		3.1 (111)
Anamosa		33.3 (85)		23.8 (110)		25.0 (70)		5.3 (61)		0.0 (1)
Ankeny		4.3 (13)		6.7 (35)		0.0 (104)		0.0 (1)		2.9 (110)
Atlantic		46.2 (126)		26.1 (121)		45.5 (35)		7.7 (89)		0.0 (1)
Audubon		12.5 (26)		0.0 (1)		100.0 (1)		12.5 (114)		9.1 (131)
Avoca		50.0 (129)		0.0 (1)		100.0 (1)		50.0 (147)		0.0 (1)
Bedford		50.0 (129)		0.0 (1)		66.7 (9)		16.7 (131)		0.0 (1)
Belle Plaine		50.0 (129)		40.0 (135)		33.3 (47)		12.5 (114)		0.0 (1)
Bloomfield		0.0 (1)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Boone		21.4 (44)		12.5 (60)		25.0 (70)		7.1 (82)		0.0 (1)
Brdlwns RP		16.6 (32)		13.4 (65)		22.0 (82)		3.9 (53)		1.6 (99)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Brooklyn		25.0 (54)		25.0 (114)		0.0 (104)		20.0 (137)		0.0 (1)
Burlington		46.4 (127)		21.0 (98)		47.0 (33)		13.6 (123)		2.3 (103)
CHC		32.0 (83)		16.2 (79)		22.2 (80)		13.6 (123)		4.4 (117)
CR Urban		28.3 (69)		15.8 (75)		25.8 (68)		6.9 (80)		0.7 (87)
Carlisle		100.0 (147)		100.0 (147)		0.0 (104)		0.0 (1)		50.0 (147)
Carroll		37.5 (101)		18.6 (89)		50.0 (16)		6.9 (80)		4.7 (119)
Cedar Falls		16.7 (33)		0.0 (1)		0.0 (104)		7.7 (89)		0.0 (1)
Centervill		38.1 (107)		16.7 (80)		0.0 (104)		9.5 (101)		0.0 (1)
Chariton		24.1 (53)		7.1 (37)		50.0 (16)		7.1 (82)		0.0 (1)
Charles City		27.5 (66)		12.7 (63)		46.2 (34)		6.0 (72)		0.0 (1)
Cherokee		25.0 (54)		10.7 (53)		50.0 (16)		3.6 (49)		3.6 (114)
Clarinda		47.4 (128)		33.3 (128)		37.5 (43)		10.5 (107)		0.0 (1)
Clarion		3.2 (12)		2.9 (25)		100.0 (1)		9.7 (102)		0.0 (1)
Clinton		42.8 (119)		23.3 (106)		35.5 (46)		8.3 (95)		0.6 (86)
Colfax		14.3 (28)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Columbus J		0.0 (1)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Corning		14.3 (28)		20.0 (93)		0.0 (104)		28.6 (144)		11.1 (133)
Corydon		42.9 (120)		22.2 (101)		33.3 (47)		12.5 (114)		0.0 (1)
Council Blfs		38.6 (109)		23.5 (108)		24.3 (79)		7.1 (82)		0.3 (85)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Cresco		31.6 (82)		25.0 (114)		16.7 (90)		21.1 (138)		6.3 (126)
Creston		31.1 (80)		15.9 (77)		30.8 (59)		4.4 (56)		2.3 (103)
Dallas Ctr		0.0 (1)		100.0 (147)		0.0 (104)		0.0 (1)		0.0 (1)
Dayton		0.0 (1)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Decorah		30.4 (74)		9.1 (44)		33.3 (47)		17.4 (135)		4.8 (120)
Denison		11.1 (22)		7.5 (39)		20.0 (83)		1.1 (43)		0.0 (1)
Dewitt		33.3 (85)		20.0 (93)		0.0 (104)		22.2 (139)		0.0 (1)
Dubuque		16.7 (33)		8.0 (42)		14.6 (95)		0.4 (42)		0.0 (1)
Dyersville		25.0 (54)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Eagle Grove		25.0 (54)		5.0 (29)		60.0 (12)		5.0 (59)		5.0 (122)
Edgerton		27.6 (67)		16.9 (84)		26.4 (67)		12.2 (113)		1.3 (95)
Eldora		40.0 (110)		33.3 (128)		50.0 (16)		60.0 (148)		16.7 (142)
Elkader		27.3 (64)		15.2 (73)		44.4 (37)		12.5 (114)		9.1 (131)
Emmetsburg		43.8 (122)		16.7 (80)		57.1 (14)		12.5 (114)		16.7 (142)
Esthervill		22.2 (47)		12.2 (59)		37.5 (43)		16.7 (131)		16.3 (141)
Fairfiel		38.1 (107)		15.4 (74)		0.0 (104)		4.8 (58)		0.0 (1)
Forest City		30.6 (76)		8.9 (43)		50.0 (16)		5.6 (63)		0.0 (1)
Fort Dodge		41.9 (117)		23.7 (109)		31.9 (58)		14.6 (128)		2.6 (109)
Ft Madison		45.0 (124)		19.6 (91)		50.0 (16)		12.5 (114)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Garner		40.0 (110)		10.0 (49)		66.7 (9)		0.0 (1)		0.0 (1)
Glenwood		70.0 (145)		20.0 (93)		28.6 (61)		0.0 (1)		0.0 (1)
Grand Junction		100.0 (147)		0.0 (1)		100.0 (1)		0.0 (1)		0.0 (1)
Greenfield		12.5 (26)		7.7 (40)		50.0 (16)		0.0 (1)		7.7 (128)
Grinnell		25.0 (54)		25.0 (114)		33.3 (47)		12.5 (114)		13.3 (138)
Grundy Ctr		14.3 (28)		33.3 (128)		0.0 (104)		0.0 (1)		0.0 (1)
Guthrie Ct		71.4 (146)		50.0 (140)		0.0 (104)		14.3 (125)		0.0 (1)
Hampton		9.5 (20)		10.2 (51)		20.0 (83)		2.4 (46)		0.0 (1)
Harlan		42.3 (118)		27.3 (123)		16.7 (90)		3.8 (51)		0.0 (1)
Hawarden		35.3 (94)		14.3 (66)		40.0 (41)		0.0 (1)		0.0 (1)
Hispanic Cln		0.0 (1)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Humboldt		16.7 (33)		5.0 (29)		33.3 (47)		5.6 (63)		0.0 (1)
ILH		30.4 (74)		19.7 (92)		24.6 (78)		4.7 (57)		1.0 (92)
Ida Grove		41.2 (114)		15.8 (75)		14.3 (96)		0.0 (1)		5.0 (122)
Independen		25.0 (54)		22.7 (104)		20.0 (83)		10.0 (103)		4.5 (118)
Indianola		29.0 (70)		20.7 (97)		16.7 (90)		3.0 (47)		3.4 (113)
Iowa City		15.0 (31)		4.6 (28)		41.7 (40)		3.6 (49)		0.9 (89)
Iowa Falls		25.0 (54)		0.0 (1)		0.0 (104)		8.3 (95)		0.0 (1)
Jamaica		50.0 (129)		50.0 (140)		50.0 (16)		0.0 (1)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Jefferson		30.8 (77)		7.1 (37)		25.0 (70)		0.0 (1)		0.0 (1)
KR		22.2 (47)		10.0 (49)		50.0 (16)		11.1 (108)		0.0 (1)
Keokuk		43.3 (121)		23.8 (110)		28.6 (61)		23.3 (141)		2.5 (108)
Keosauqua		22.2 (47)		44.4 (139)		0.0 (104)		0.0 (1)		11.1 (133)
Knoxville		36.0 (97)		24.0 (113)		17.6 (89)		11.5 (111)		2.0 (102)
LCPH		31.3 (81)		10.3 (52)		33.3 (47)		1.5 (44)		0.0 (1)
Lamoni		50.0 (129)		16.7 (80)		33.3 (47)		12.5 (114)		0.0 (1)
Le Mars		21.9 (46)		6.1 (33)		42.9 (38)		3.1 (48)		0.0 (1)
Lenox		10.0 (21)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Leon		30.8 (77)		17.6 (86)		25.0 (70)		7.7 (89)		5.9 (124)
Logan		35.7 (95)		5.6 (32)		50.0 (16)		7.1 (82)		0.0 (1)
Manchester		20.0 (39)		11.1 (55)		0.0 (104)		0.0 (1)		0.0 (1)
Maquoketa		54.2 (138)		31.8 (127)		16.7 (90)		13.0 (122)		8.7 (130)
Marengo		66.7 (142)		50.0 (140)		0.0 (104)		0.0 (1)		0.0 (1)
Marion		33.3 (85)		19.3 (90)		29.2 (60)		8.6 (98)		0.7 (87)
Marshalltown		23.1 (51)		14.8 (68)		13.9 (97)		6.7 (76)		2.4 (107)
Mason City		32.5 (84)		15.0 (71)		27.7 (65)		6.8 (78)		1.4 (96)
Monticello		57.1 (139)		50.0 (140)		0.0 (104)		28.6 (144)		0.0 (1)
Mt Ayr		50.0 (129)		14.3 (66)		0.0 (104)		10.0 (103)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indic tors (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Mt Pleasan		37.1 (100)		23.3 (106)		33.3 (47)		5.7 (67)		2.3 (103)
Mtown Dtown		0.0 (1)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Muscatine		29.3 (71)		17.5 (85)		33.3 (47)		11.4 (109)		1.0 (92)
Nevada		33.3 (85)		20.0 (93)		0.0 (104)		0.0 (1)		0.0 (1)
New Hampto		11.1 (22)		6.7 (35)		50.0 (16)		0.0 (1)		0.0 (1)
Newton		41.3 (116)		30.7 (126)		20.0 (83)		5.8 (69)		1.4 (96)
Northwood		11.1 (22)		9.1 (44)		100.0 (1)		0.0 (1)		0.0 (1)
Norwalk		40.0 (110)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Oakridge		20.0 (39)		40.0 (135)		0.0 (104)		0.0 (1)		0.0 (1)
Oelwein		50.0 (129)		55.6 (146)		10.0 (100)		23.8 (142)		38.9 (146)
Onawa		18.8 (37)		25.0 (114)		0.0 (104)		6.3 (73)		0.0 (1)
Orange Cty		6.7 (16)		0.0 (1)		0.0 (104)		6.7 (76)		5.9 (124)
Osage		20.8 (42)		14.8 (68)		20.0 (83)		0.0 (1)		3.8 (116)
Osceola		36.7 (99)		26.1 (121)		6.3 (103)		8.0 (93)		0.0 (1)
Oskaloosa		25.7 (62)		25.0 (114)		12.5 (98)		5.7 (67)		0.0 (1)
Ottumwa		26.0 (63)		18.4 (88)		8.8 (101)		10.1 (105)		1.1 (94)
Pella		9.1 (19)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Perry		21.4 (44)		12.5 (60)		0.0 (104)		2.2 (45)		0.0 (1)
Pocahontas		50.0 (129)		42.9 (138)		0.0 (104)		16.7 (131)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)										
Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Postville		5.7 (14)		3.0 (26)		50.0 (16)		5.9 (70)		3.1 (111)
Red Oak		63.6 (141)		39.1 (134)		8.3 (102)		9.1 (100)		0.0 (1)
Redfield		0.0 (1)		50.0 (140)		0.0 (104)		0.0 (1)		50.0 (147)
Refugee Cln		0.0 (1)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Rock Rapid		11.1 (22)		4.5 (27)		50.0 (16)		5.6 (63)		0.0 (1)
Rock Valle		3.1 (11)		0.0 (1)		100.0 (1)		0.0 (1)		0.0 (1)
Rockwell C		66.7 (142)		40.0 (135)		16.7 (90)		22.2 (139)		0.0 (1)
SDM		23.2 (52)		9.2 (47)		27.9 (64)		6.5 (75)		0.9 (89)
Sac City		57.1 (139)		33.3 (128)		40.0 (41)		14.3 (125)		8.3 (129)
Sheldon		29.6 (72)		9.1 (44)		55.6 (15)		7.1 (82)		12.5 (136)
Shenandoah		44.4 (123)		22.2 (101)		27.3 (66)		7.4 (88)		3.7 (115)
Sibley		33.3 (85)		11.1 (55)		0.0 (104)		33.3 (146)		11.1 (133)
Sidney		50.0 (129)		16.7 (80)		50.0 (16)		16.7 (131)		0.0 (1)
Sigourney		45.5 (125)		50.0 (140)		0.0 (104)		18.2 (136)		0.0 (1)
Sioux Cent		6.8 (17)		6.3 (34)		33.3 (47)		0.0 (1)		0.0 (1)
Sioux City		18.8 (37)		9.7 (48)		33.3 (47)		4.2 (54)		0.9 (89)
Spencer		36.2 (98)		22.0 (100)		25.0 (70)		14.9 (129)		15.3 (139)
Spirit Lak		34.6 (92)		22.6 (103)		62.5 (11)		15.4 (130)		16.7 (142)
Storm Lake		7.3 (18)		1.8 (24)		60.0 (12)		8.5 (97)		4.9 (121)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Stuart		37.5 (101)		28.6 (124)		0.0 (104)		25.0 (143)		0.0 (1)
Tama		37.5 (101)		17.9 (87)		12.5 (98)		7.7 (89)		2.3 (103)
Tama/Settl		0.0 (1)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Tipton		20.0 (39)		25.0 (114)		0.0 (104)		0.0 (1)		12.5 (136)
Traer		66.7 (142)		0.0 (1)		100.0 (1)		0.0 (1)		0.0 (1)
UIHC		0.0 (1)		0.0 (1)		0.0 (104)		0.0 (1)		0.0 (1)
Vinton		40.0 (110)		15.0 (71)		50.0 (16)		8.0 (93)		0.0 (1)
WDM		27.3 (64)		14.9 (70)		28.6 (61)		0.0 (1)		0.0 (1)
Washington		37.9 (106)		21.4 (99)		25.0 (70)		10.3 (106)		0.0 (1)
Waterloo		30.1 (73)		16.1 (78)		25.6 (69)		11.4 (109)		1.6 (99)
Waukee		6.3 (15)		0.0 (1)		50.0 (16)		3.8 (51)		0.0 (1)
Waukon		22.2 (47)		11.8 (57)		25.0 (70)		5.6 (63)		35.3 (145)
Waverly		33.3 (85)		13.3 (64)		0.0 (104)		5.9 (70)		6.3 (126)
Wbstr Ct		35.1 (93)		12.5 (60)		45.0 (36)		5.3 (61)		1.6 (99)
West Union		35.7 (95)		23.1 (105)		0.0 (104)		14.3 (125)		15.4 (140)
Winterset		37.5 (101)		11.8 (57)		50.0 (16)		6.3 (73)		0.0 (1)
Iowa		**27.2**		**15.4**		**27.6**		**7.3**		**2.0**
National (Prior Year)	**126,132**	**26.9**	**133,382**	**14.1**	**21,263**	**35.7**	**97,728**	**11.9**	**118,446**	**6.6**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley		0.0 (1)	66.7 (148)		0.0 (1)		0.0 (1)		66.7 (91)
Adel		0.0 (1)	15.4 (132)		0.0 (1)		0.0 (1)		53.8 (135)
Albia		5.9 (65)	11.8 (109)		7.1 (52)		0.0 (1)		52.9 (137)
Algona		11.1 (116)	8.3 (83)		10.3 (81)		3.8 (104)		80.6 (31)
Allen WHC		7.8 (89)	6.7 (62)		6.9 (50)		6.2 (121)		70.0 (80)
Allison		19.0 (143)	9.5 (98)		23.5 (141)		7.7 (130)		76.2 (49)
Altoona		0.0 (1)	3.2 (37)		3.7 (34)		0.0 (1)		77.4 (47)
Ames		8.4 (95)	9.0 (91)		8.7 (68)		5.2 (115)		79.6 (38)
Anamosa		12.5 (126)	8.3 (83)		12.5 (106)		4.8 (111)		62.5 (112)
Ankeny		8.2 (92)	12.2 (114)		7.3 (54)		0.0 (1)		83.7 (21)
Atlantic		0.0 (1)	6.3 (59)		3.4 (33)		0.0 (1)		59.4 (127)
Audubon		9.1 (98)	27.3 (146)		10.0 (75)		0.0 (1)		63.6 (109)
Avoca		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		50.0 (139)
Bedford		0.0 (1)	12.5 (115)		0.0 (1)		0.0 (1)		62.5 (112)
Belle Plaine		0.0 (1)	12.5 (115)		0.0 (1)		0.0 (1)		75.0 (54)
Bloomfield		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		60.0 (120)
Boone		16.0 (134)	0.0 (1)		18.2 (135)		11.1 (138)		72.0 (73)
Brdlwns RP		8.4 (95)	6.6 (61)		10.7 (90)		3.4 (100)		78.3 (44)
Brooklyn		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		60.0 (120)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Burlington		6.6 (74)	8.3 (83)		6.8 (49)		4.3 (107)		49.7 (144)
CHC		11.9 (124)	8.1 (80)		11.3 (98)		6.6 (124)		67.5 (90)
CR Urban		8.7 (97)	9.1 (92)		7.8 (59)		4.4 (108)		73.9 (66)
Carlisle		0.0 (1)	0.0 (1)		50.0 (148)		0.0 (1)		77.8 (46)
Carroll		4.3 (50)	8.5 (88)		7.4 (55)		1.1 (84)		74.5 (60)
Cedar Falls		6.7 (76)	13.3 (123)		7.1 (52)		0.0 (1)		80.0 (34)
Centervill		4.2 (47)	12.5 (115)		8.3 (64)		0.0 (1)		54.2 (134)
Chariton		6.7 (76)	16.7 (133)		17.9 (134)		0.0 (1)		73.3 (68)
Charles City		11.1 (116)	11.1 (105)		10.6 (89)		9.5 (137)		75.9 (52)
Cherokee		11.4 (122)	5.7 (55)		12.5 (106)		7.1 (128)		80.0 (34)
Clarinda		4.8 (56)	4.8 (45)		10.5 (88)		0.0 (1)		75.0 (54)
Clarion		8.1 (91)	5.4 (52)		16.1 (125)		0.0 (1)		67.6 (89)
Clinton		6.3 (69)	4.8 (45)		13.9 (115)		2.2 (89)		68.8 (85)
Colfax		10.0 (105)	10.0 (101)		22.2 (140)		0.0 (1)		70.0 (80)
Columbus J		9.1 (98)	0.0 (1)		11.1 (92)		12.5 (141)		90.9 (10)
Corning		0.0 (1)	10.0 (101)		0.0 (1)		0.0 (1)		60.0 (120)
Corydon		6.3 (69)	25.0 (144)		16.7 (126)		0.0 (1)		87.5 (14)
Council Blfs		11.0 (115)	8.2 (81)		12.3 (105)		6.2 (121)		71.8 (75)
Cresco		11.1 (116)	22.2 (143)		16.7 (126)		0.0 (1)		83.3 (22)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Creston		3.3 (45)	11.5 (108)		4.2 (35)		2.2 (89)		70.5 (78)
Dallas Ctr		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		100.0 (1)
Dayton		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (147)
Decorah		10.7 (114)	17.9 (139)		8.0 (62)		4.5 (109)		93.1 (8)
Denison		9.3 (103)	2.8 (35)		7.8 (59)		4.2 (106)		65.7 (101)
Dewitt		9.1 (98)	0.0 (1)		11.1 (92)		0.0 (1)		63.6 (109)
Dubuque		6.3 (69)	8.6 (89)		10.4 (84)		2.2 (89)		55.8 (131)
Dyersville		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		50.0 (139)
Eagle Grove		4.3 (50)	4.3 (40)		14.3 (118)		0.0 (1)		78.3 (44)
Edgerton		8.0 (90)	9.4 (97)		9.4 (70)		1.9 (88)		68.4 (87)
Eldora		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		75.0 (54)
Elkader		5.7 (63)	14.3 (125)		10.0 (75)		0.0 (1)		85.7 (18)
Emmetsburg		13.0 (129)	13.0 (122)		0.0 (1)		12.5 (141)		60.9 (119)
Esthervill		6.3 (69)	12.5 (115)		10.3 (81)		0.0 (1)		62.5 (112)
Fairfiel		11.1 (116)	11.1 (105)		8.3 (64)		9.1 (135)		70.4 (79)
Forest City		4.4 (52)	17.8 (138)		10.3 (81)		2.9 (95)		66.7 (91)
Fort Dodge		7.7 (84)	8.2 (81)		11.3 (98)		0.7 (83)		66.5 (100)
Ft Madison		7.7 (84)	9.6 (99)		2.4 (32)		2.5 (93)		55.8 (131)
Garner		15.0 (132)	10.0 (101)		6.7 (46)		0.0 (1)		70.0 (80)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Glenwood		18.8 (142)	0.0 (1)		14.3 (118)		8.3 (132)		62.5 (112)
Grand Junction		100.0 (147)	0.0 (1)		0.0 (1)		100.0 (148)		100.0 (1)
Greenfield		0.0 (1)	12.5 (115)		7.7 (57)		0.0 (1)		81.3 (29)
Grinnell		5.9 (65)	11.8 (109)		6.7 (46)		0.0 (1)		64.7 (106)
Grundy Ctr		12.5 (126)	12.5 (115)		14.3 (118)		0.0 (1)		50.0 (139)
Guthrie Ct		7.1 (80)	0.0 (1)		7.7 (57)		8.3 (132)		78.6 (40)
Hampton		4.2 (47)	16.7 (133)		4.7 (38)		0.0 (1)		65.3 (104)
Harlan		2.4 (41)	7.3 (71)		13.9 (115)		0.0 (1)		80.5 (33)
Hawarden		11.8 (123)	17.6 (137)		11.8 (100)		6.7 (125)		82.4 (26)
Hispanic Cln		16.7 (135)	16.7 (133)		0.0 (1)		16.7 (145)		100.0 (1)
Humboldt		0.0 (1)	5.6 (53)		11.8 (100)		0.0 (1)		88.9 (12)
ILH		8.2 (92)	6.0 (56)		9.8 (74)		2.9 (95)		71.9 (74)
Ida Grove		0.0 (1)	4.5 (42)		4.5 (36)		0.0 (1)		86.4 (17)
Independen		10.3 (108)	17.2 (136)		12.5 (106)		4.8 (111)		62.1 (117)
Indianola		18.2 (140)	14.5 (128)		12.5 (106)		11.4 (140)		72.7 (70)
Iowa City		11.1 (116)	6.8 (64)		10.4 (84)		5.1 (114)		82.7 (25)
Iowa Falls		5.9 (65)	0.0 (1)		0.0 (1)		6.7 (125)		76.5 (48)
Jamaica		33.3 (146)	0.0 (1)		33.3 (145)		0.0 (1)		66.7 (91)
Jefferson		7.1 (80)	7.1 (69)		21.4 (139)		0.0 (1)		78.6 (40)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
KR		14.3 (130)	14.3 (125)		25.0 (142)		11.1 (138)		78.6 (40)
Keokuk		7.3 (82)	4.9 (47)		17.6 (133)		3.6 (102)		56.1 (130)
Keosauqua		7.7 (84)	7.7 (77)		8.3 (64)		0.0 (1)		53.8 (135)
Knoxville		5.8 (64)	7.0 (67)		9.4 (70)		1.7 (87)		75.6 (53)
LCPH		10.5 (111)	7.4 (75)		15.2 (123)		1.5 (85)		65.6 (102)
Lamoni		0.0 (1)	25.0 (144)		12.5 (106)		0.0 (1)		75.0 (54)
Le Mars		4.9 (57)	7.3 (71)		11.1 (92)		0.0 (1)		92.7 (9)
Lenox		8.3 (94)	8.3 (83)		0.0 (1)		9.1 (135)		58.3 (129)
Leon		11.1 (116)	5.6 (53)		20.0 (136)		0.0 (1)		66.7 (91)
Logan		20.0 (144)	20.0 (142)		5.0 (39)		15.8 (144)		95.0 (7)
Manchester		0.0 (1)	9.1 (92)		10.0 (75)		0.0 (1)		36.4 (146)
Maquoketa		6.9 (79)	6.9 (65)		16.0 (124)		0.0 (1)		62.1 (117)
Marengo		16.7 (135)	33.3 (147)		0.0 (1)		16.7 (145)		50.0 (139)
Marion		4.4 (52)	6.9 (65)		10.0 (75)		1.5 (85)		68.6 (86)
Marshalltown		6.6 (74)	7.3 (71)		9.0 (69)		5.0 (113)		74.5 (60)
Mason City		10.3 (108)	7.3 (71)		14.4 (122)		4.6 (110)		64.4 (107)
Monticello		0.0 (1)	11.1 (105)		11.1 (92)		0.0 (1)		66.7 (91)
Mt Ayr		0.0 (1)	0.0 (1)		10.0 (75)		0.0 (1)		66.7 (91)
Mt Pleasan		9.1 (98)	9.1 (92)		5.7 (42)		3.0 (98)		63.6 (109)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Mtown Dtown		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		100.0 (1)
Muscatine		10.6 (113)	8.0 (79)		12.7 (114)		4.1 (105)		76.0 (50)
Nevada		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		80.0 (34)
New Hampto		5.0 (59)	15.0 (129)		5.3 (40)		0.0 (1)		80.0 (34)
Newton		10.0 (105)	5.0 (49)		10.4 (84)		6.8 (127)		66.7 (91)
Northwood		0.0 (1)	0.0 (1)		11.1 (92)		0.0 (1)		81.8 (27)
Norwalk		30.0 (145)	0.0 (1)		33.3 (145)		0.0 (1)		60.0 (120)
Oakridge		0.0 (1)	0.0 (1)		16.7 (126)		0.0 (1)		42.9 (145)
Oelwein		4.5 (54)	4.5 (42)		9.5 (72)		5.3 (116)		59.1 (128)
Onawa		10.5 (111)	0.0 (1)		5.9 (43)		6.3 (123)		68.4 (87)
Orange Cty		4.5 (54)	9.1 (92)		0.0 (1)		5.3 (116)		81.8 (27)
Osage		3.7 (46)	7.4 (75)		4.5 (36)		0.0 (1)		74.1 (64)
Osceola		9.1 (98)	9.1 (92)		14.3 (118)		0.0 (1)		72.7 (70)
Oskaloosa		2.2 (40)	4.3 (40)		8.6 (67)		0.0 (1)		52.2 (138)
Ottumwa		5.2 (60)	6.7 (62)		7.8 (59)		2.4 (92)		69.7 (83)
Pella		0.0 (1)	19.0 (141)		0.0 (1)		0.0 (1)		81.0 (30)
Perry		10.0 (105)	5.0 (49)		12.5 (106)		6.1 (120)		73.3 (68)
Pocahontas		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		71.4 (76)
Postville		7.7 (84)	7.7 (77)		8.1 (63)		2.9 (95)		84.6 (20)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Red Oak		3.0 (43)	12.1 (113)		6.7 (46)		0.0 (1)		72.7 (70)
Redfield		0.0 (1)	0.0 (1)		20.0 (136)		0.0 (1)		60.0 (120)
Refugee Cln		14.3 (130)	14.3 (125)		12.5 (106)		14.3 (143)		100.0 (1)
Rock Rapid		0.0 (1)	12.0 (112)		12.0 (103)		0.0 (1)		76.0 (50)
Rock Valle		4.9 (57)	9.8 (100)		16.7 (126)		0.0 (1)		87.8 (13)
Rockwell C		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		60.0 (120)
SDM		6.1 (68)	8.4 (87)		11.0 (91)		3.2 (99)		74.0 (65)
Sac City		16.7 (135)	0.0 (1)		33.3 (145)		0.0 (1)		75.0 (54)
Sheldon		17.1 (139)	2.9 (36)		17.2 (132)		0.0 (1)		85.7 (18)
Shenandoah		16.7 (135)	0.0 (1)		26.9 (143)		5.3 (116)		83.3 (22)
Sibley		0.0 (1)	10.0 (101)		28.6 (144)		0.0 (1)		90.0 (11)
Sidney		0.0 (1)	12.5 (115)		0.0 (1)		0.0 (1)		87.5 (14)
Sigourney		18.2 (140)	18.2 (140)		0.0 (1)		18.2 (147)		54.5 (133)
Sioux Cent		4.2 (47)	6.3 (59)		6.5 (44)		0.0 (1)		87.5 (14)
Sioux City		7.3 (82)	4.9 (47)		11.2 (97)		3.6 (102)		70.6 (77)
Spencer		5.3 (61)	8.8 (90)		6.5 (44)		0.0 (1)		66.7 (91)
Spirit Lak		3.0 (43)	6.1 (57)		12.0 (103)		0.0 (1)		78.8 (39)
Storm Lake		15.0 (132)	3.3 (38)		16.7 (126)		7.5 (129)		74.2 (63)
Stuart		12.5 (126)	0.0 (1)		12.5 (106)		0.0 (1)		50.0 (139)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Tama		6.5 (73)	15.2 (131)		9.5 (72)		2.6 (94)		65.2 (105)
Tama/Settl		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (147)
Tipton		0.0 (1)	0.0 (1)		16.7 (126)		0.0 (1)		62.5 (112)
Traer		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		66.7 (91)
UIHC		100.0 (147)	0.0 (1)		0.0 (1)		0.0 (1)		100.0 (1)
Vinton		0.0 (1)	13.8 (124)		6.9 (50)		0.0 (1)		69.0 (84)
WDM		10.4 (110)	4.5 (42)		14.0 (117)		8.2 (131)		80.6 (31)
Washington		6.7 (76)	0.0 (1)		7.4 (55)		0.0 (1)		83.3 (22)
Waterloo		9.3 (103)	7.0 (67)		10.4 (84)		3.5 (101)		74.3 (62)
Waukee		2.9 (42)	11.8 (109)		0.0 (1)		0.0 (1)		73.5 (67)
Waukon		0.0 (1)	15.0 (129)		11.8 (100)		0.0 (1)		75.0 (54)
Waverly		5.3 (61)	5.3 (51)		5.3 (40)		0.0 (1)		60.0 (120)
Wbstr Ct		12.3 (125)	6.2 (58)		20.0 (136)		8.3 (132)		78.5 (43)
West Union		0.0 (1)	7.1 (69)		0.0 (1)		0.0 (1)		64.3 (108)
Winterset		7.7 (84)	3.8 (39)		10.0 (75)		5.6 (119)		65.4 (103)
Iowa		**8.1**	**7.6**		**10.4**		**3.5**		**72.0**
National (Prior Year)	**99,198**	**9.7**	**6.5**	**100,365**	**12.4**	**84,851**	**4.1**	**99,674**	**52.7**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Ackley	54.5	0.0	45.5	0.0	0.0	0.0	0.0
Adel	84.8	3.0	12.1	0.0	0.0	0.0	0.0
Albia	94.2	1.9	1.9	1.9	0.0	0.0	0.0
Algona	79.7	3.4	14.4	0.0	0.0	2.5	0.0
Allen WHC	49.0	32.3	8.0	0.0	7.3	3.5	0.0
Allison	91.0	1.5	6.0	0.0	0.0	1.5	0.0
Altoona	76.9	13.0	6.5	0.0	2.8	0.9	0.0
Ames	70.5	9.7	9.9	0.3	7.8	1.8	0.0
Anamosa	95.1	1.2	3.7	0.0	0.0	0.0	0.0
Ankeny	75.2	9.8	8.5	0.0	3.3	3.3	0.0
Atlantic	90.8	0.9	8.3	0.0	0.0	0.0	0.0
Audubon	92.1	0.0	7.9	0.0	0.0	0.0	0.0
Avoca	83.3	11.1	5.6	0.0	0.0	0.0	0.0
Bedford	91.7	0.0	0.0	0.0	0.0	8.3	0.0
Belle Plaine	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Bloomfield	89.5	5.3	5.3	0.0	0.0	0.0	0.0
Boone	81.5	2.5	11.1	0.0	0.0	4.9	0.0
Brdlwns RP	39.2	24.3	21.4	0.3	9.9	4.8	0.0
Brooklyn	66.7	0.0	33.3	0.0	0.0	0.0	0.0
Burlington	75.5	18.6	4.3	0.0	0.2	1.4	0.0
CHC	57.6	21.2	14.4	0.3	2.1	4.5	0.0
CR Urban	48.2	37.4	7.1	0.1	1.6	5.7	0.0
Carlisle	89.2	2.7	5.4	0.0	2.7	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Carroll	75.9	6.8	16.0	0.0	0.8	0.4	0.0
Cedar Falls	81.1	0.0	10.8	0.0	8.1	0.0	0.0
Centervill	91.1	1.3	6.3	0.0	0.0	1.3	0.0
Chariton	94.6	0.0	5.4	0.0	0.0	0.0	0.0
Charles City	73.5	10.8	10.3	0.0	4.3	1.1	0.0
Cherokee	86.4	0.9	11.8	0.0	0.0	0.9	0.0
Clarinda	93.5	0.0	1.6	0.0	0.0	4.8	0.0
Clarion	39.2	0.0	60.8	0.0	0.0	0.0	0.0
Clinton	78.7	9.3	9.3	0.0	0.4	2.4	0.0
Colfax	93.1	0.0	6.9	0.0	0.0	0.0	0.0
Columbus J	9.5	0.0	47.6	4.8	38.1	0.0	0.0
Corning	93.9	0.0	6.1	0.0	0.0	0.0	0.0
Corydon	98.0	0.0	2.0	0.0	0.0	0.0	0.0
Council Blfs	78.1	3.4	15.4	0.4	0.7	1.9	0.0
Cresco	91.5	0.0	7.0	0.0	1.4	0.0	0.0
Creston	86.0	1.3	11.8	0.0	0.0	0.9	0.0
Dallas Ctr	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Dayton	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Decorah	81.6	1.3	15.8	0.0	0.0	1.3	0.0
Denison	31.6	3.6	58.1	0.3	6.3	0.0	0.0
Dewitt	76.9	3.8	11.5	0.0	0.0	7.7	0.0
Dubuque	72.5	14.0	6.9	0.5	3.9	2.3	0.0
Dyersville	81.3	12.5	6.3	0.0	0.0	0.0	0.0
Eagle Grove	32.4	1.4	66.2	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Edgerton	56.6	25.9	9.4	0.7	1.7	5.6	0.0
Eldora	87.0	0.0	13.0	0.0	0.0	0.0	0.0
Elkader	94.1	0.0	4.9	0.0	1.0	0.0	0.0
Emmetsburg	92.5	3.0	4.5	0.0	0.0	0.0	0.0
Esthervill	64.2	2.2	32.1	0.0	0.0	1.5	0.0
Fairfiel	88.7	0.0	7.0	0.0	4.2	0.0	0.0
Forest City	83.4	6.2	10.3	0.0	0.0	0.0	0.0
Fort Dodge	76.3	9.4	11.4	0.5	0.3	2.2	0.0
Ft Madison	88.8	2.9	7.6	0.0	0.0	0.6	0.0
Garner	79.0	1.2	17.3	1.2	0.0	1.2	0.0
Glenwood	88.5	0.0	11.5	0.0	0.0	0.0	0.0
Grand Junction	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Greenfield	87.3	0.0	10.9	0.0	1.8	0.0	0.0
Grinnell	87.2	4.7	7.0	0.0	0.0	1.2	0.0
Grundy Ctr	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Guthrie Ct	88.7	1.9	7.5	0.0	0.0	1.9	0.0
Hampton	52.2	2.2	44.9	0.0	0.0	0.7	0.0
Harlan	83.0	0.9	15.1	0.0	0.0	0.9	0.0
Hawarden	59.3	0.0	39.0	1.7	0.0	0.0	0.0
Hispanic Cln	9.1	0.0	90.9	0.0	0.0	0.0	0.0
Humboldt	63.2	2.9	32.4	0.0	0.0	1.5	0.0
ILH	38.7	16.4	36.3	0.1	5.5	3.0	0.0
Ida Grove	83.3	1.3	14.1	0.0	0.0	1.3	0.0
Independen	94.9	2.0	3.1	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Indianola	90.1	3.5	3.5	0.0	1.2	1.7	0.0
Iowa City	35.1	42.3	17.7	0.3	2.2	2.3	0.0
Iowa Falls	82.2	2.2	13.3	0.0	0.0	2.2	0.0
Jamaica	75.0	0.0	25.0	0.0	0.0	0.0	0.0
Jefferson	83.9	0.0	14.3	0.0	1.8	0.0	0.0
KR	55.0	22.5	2.5	0.0	17.5	2.5	0.0
Keokuk	89.7	3.8	4.5	0.0	0.6	1.3	0.0
Keosauqua	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Knoxville	92.7	2.0	4.1	0.0	0.0	1.2	0.0
LCPH	66.7	22.2	5.1	1.0	2.0	3.0	0.0
Lamoni	66.7	11.1	11.1	0.0	0.0	11.1	0.0
Le Mars	66.7	5.1	23.2	0.7	3.6	0.7	0.0
Lenox	63.6	0.0	33.3	0.0	3.0	0.0	0.0
Leon	98.0	2.0	0.0	0.0	0.0	0.0	0.0
Logan	93.0	0.0	5.8	1.2	0.0	0.0	0.0
Manchester	95.6	0.0	4.4	0.0	0.0	0.0	0.0
Manning	50.0	0.0	0.0	0.0	50.0	0.0	0.0
Maquoketa	82.4	5.5	3.3	0.0	4.4	4.4	0.0
Marengo	88.0	0.0	4.0	0.0	0.0	8.0	0.0
Marion	82.7	5.1	7.7	0.0	0.7	3.7	0.0
Marshalltown	41.8	3.3	45.0	0.2	8.6	1.0	0.0
Mason City	76.8	5.6	11.0	0.3	3.3	3.0	0.0
Monticello	92.9	3.6	3.6	0.0	0.0	0.0	0.0
Mt Ayr	90.3	0.0	9.7	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Mt Pleasan	81.9	2.4	14.5	0.0	1.2	0.0	0.0
Mtown Dtown	0.0	0.0	66.7	0.0	33.3	0.0	0.0
Muscatine	58.3	8.5	30.6	0.0	2.0	0.6	0.0
Nevada	53.3	0.0	46.7	0.0	0.0	0.0	0.0
New Hampto	75.0	1.8	21.4	0.0	0.0	1.8	0.0
Newton	80.1	4.9	10.9	1.5	2.3	0.4	0.0
North Towne	62.4	19.9	8.2	0.0	5.4	4.1	0.0
Northwood	83.3	3.3	6.7	0.0	3.3	3.3	0.0
Norwalk	81.4	0.0	14.0	0.0	4.7	0.0	0.0
Oakridge	25.0	50.0	12.5	0.0	0.0	12.5	0.0
Oelwein	86.0	4.7	9.3	0.0	0.0	0.0	0.0
Onawa	97.3	0.0	0.0	1.4	0.0	1.4	0.0
Orange Cty	43.4	2.6	52.6	0.0	1.3	0.0	0.0
Osage	90.7	1.2	5.8	0.0	0.0	2.3	0.0
Osceola	71.8	0.6	26.3	0.0	1.3	0.0	0.0
Oskaloosa	89.3	5.3	4.7	0.0	0.7	0.0	0.0
Ottumwa	72.1	7.1	14.8	0.0	5.0	0.9	0.2
Pella	88.9	0.0	6.7	0.0	4.4	0.0	0.0
Perry	33.9	9.1	51.5	0.6	1.2	3.6	0.0
Pocahontas	87.0	8.7	4.3	0.0	0.0	0.0	0.0
Pomeroy	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Postville	32.0	9.0	59.0	0.0	0.0	0.0	0.0
Red Oak	92.1	0.0	7.9	0.0	0.0	0.0	0.0
Redfield	100.0	0.0	0.0	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Refugee Cln	3.8	92.3	3.8	0.0	0.0	0.0	0.0
Rock Rapid	83.0	0.0	13.2	0.0	3.8	0.0	0.0
Rock Valle	37.6	0.0	58.7	2.8	0.0	0.9	0.0
Rockwell C	95.3	0.0	4.7	0.0	0.0	0.0	0.0
SDM	45.0	14.3	27.8	0.4	8.2	4.3	0.0
Sac City	97.3	0.0	0.0	0.0	0.0	2.7	0.0
Sheldon	64.5	3.6	27.3	2.7	1.8	0.0	0.0
Shenandoah	87.3	0.0	12.7	0.0	0.0	0.0	0.0
Sibley	52.5	5.0	42.5	0.0	0.0	0.0	0.0
Sidney	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Sigourney	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Sioux Cent	29.3	0.0	69.4	0.0	0.0	1.4	0.0
Sioux City	41.1	11.3	37.4	3.8	3.9	2.5	0.0
Spencer	82.9	0.5	15.0	0.0	1.6	0.0	0.0
Spirit Lak	87.7	1.4	8.2	0.0	0.0	2.7	0.0
Storm Lake	21.4	4.8	55.5	0.3	17.4	0.5	0.0
Stuart	88.5	3.8	0.0	0.0	0.0	7.7	0.0
Tama	44.2	0.7	42.0	9.4	0.0	3.6	0.0
Tama/Settl	33.3	0.0	0.0	66.7	0.0	0.0	0.0
Tipton	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Traer	100.0	0.0	0.0	0.0	0.0	0.0	0.0
UIHC	63.3	23.3	10.0	0.0	3.3	0.0	0.0
Vinton	96.0	0.0	4.0	0.0	0.0	0.0	0.0
WDM	49.8	23.4	20.0	1.0	3.9	2.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 3A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Combined 3 Years

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wapello	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Washington	65.7	1.9	30.6	0.0	0.0	1.9	0.0
Waterloo	54.9	25.6	9.6	0.5	5.9	3.5	0.0
Waukee	74.1	4.6	14.8	0.0	2.8	3.7	0.0
Waukon	92.8	0.0	5.8	0.0	1.4	0.0	0.0
Waverly	96.8	3.2	0.0	0.0	0.0	0.0	0.0
Wbstr Ct	59.7	2.6	32.5	0.0	5.2	0.0	0.0
West Union	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Williamsburg	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Winterset	90.3	1.6	8.1	0.0	0.0	0.0	0.0
Iowa	**60.1**	**14.0**	**18.9**	**0.5**	**3.9**	**2.6**	**0.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Ackley	0.0	0.0	0.0	63.6	36.4	0.0	0.0
Adel	0.0	3.0	6.1	72.7	18.2	0.0	0.0
Albia	0.0	5.8	7.7	63.5	23.1	0.0	0.0
Algona	0.0	0.8	4.2	56.8	37.3	0.8	0.0
Allen WHC	0.3	2.8	6.9	56.9	31.3	1.7	0.0
Allison	0.0	0.0	0.0	61.2	38.8	0.0	0.0
Altoona	0.0	2.8	2.8	51.9	38.9	3.7	0.0
Ames	0.0	1.3	4.9	58.8	34.2	0.8	0.0
Anamosa	0.0	2.4	4.9	61.0	29.3	2.4	0.0
Ankeny	0.0	2.0	2.0	66.0	26.8	3.3	0.0
Atlantic	0.0	1.8	4.6	63.3	28.4	0.9	0.9
Audubon	0.0	5.3	10.5	57.9	26.3	0.0	0.0
Avoca	0.0	0.0	0.0	66.7	33.3	0.0	0.0
Bedford	0.0	0.0	4.2	79.2	16.7	0.0	0.0
Belle Plaine	0.0	0.0	0.0	61.9	38.1	0.0	0.0
Bloomfield	0.0	0.0	0.0	73.7	15.8	10.5	0.0
Boone	0.0	0.0	7.4	64.2	27.2	1.2	0.0
Brdlwns RP	0.1	2.3	6.0	57.3	31.3	3.0	0.0
Brooklyn	0.0	0.0	0.0	88.9	11.1	0.0	0.0
Burlington	0.0	4.0	8.3	60.7	26.5	0.5	0.0
CHC	0.3	2.3	7.2	62.8	25.3	2.2	0.0
CR Urban	0.0	1.6	6.1	60.3	29.2	2.7	0.1
Carlisle	0.0	0.0	5.4	59.5	32.4	2.7	0.0
Carroll	0.0	0.8	5.9	64.6	26.6	2.1	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Cedar Falls	0.0	0.0	2.7	45.9	48.6	2.7	0.0
Centervill	0.0	0.0	7.6	65.8	24.1	2.5	0.0
Chariton	0.0	1.4	8.1	63.5	23.0	4.1	0.0
Charles City	0.5	1.1	3.2	67.0	27.6	0.5	0.0
Cherokee	0.0	0.9	11.8	70.9	16.4	0.0	0.0
Clarinda	0.0	0.0	6.5	77.4	12.9	3.2	0.0
Clarion	0.0	5.6	4.8	52.8	35.2	1.6	0.0
Clinton	0.0	2.5	9.3	64.4	21.6	2.2	0.0
Colfax	0.0	6.9	17.2	51.7	24.1	0.0	0.0
Columbus J	0.0	0.0	0.0	45.2	47.6	7.1	0.0
Corning	0.0	0.0	9.1	60.6	30.3	0.0	0.0
Corydon	0.0	2.0	0.0	66.0	32.0	0.0	0.0
Council Blfs	0.0	2.1	6.4	63.8	26.1	1.6	0.0
Cresco	0.0	4.2	2.8	54.9	36.6	1.4	0.0
Creston	0.4	3.5	12.7	58.5	23.6	1.3	0.0
Dallas Ctr	0.0	0.0	0.0	75.0	25.0	0.0	0.0
Dayton	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Decorah	0.0	1.3	6.6	51.3	39.5	1.3	0.0
Denison	0.0	2.1	2.1	63.0	29.2	3.6	0.0
Dewitt	0.0	0.0	15.4	46.2	34.6	3.8	0.0
Dubuque	0.0	2.6	8.2	60.0	27.9	1.1	0.2
Dyersville	0.0	0.0	0.0	62.5	37.5	0.0	0.0
Eagle Grove	0.0	0.0	9.9	54.9	31.0	4.2	0.0
Edgerton	0.2	3.6	7.8	59.8	26.5	2.1	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Eldora	0.0	0.0	8.7	65.2	26.1	0.0	0.0
Elkader	0.0	0.0	2.0	72.5	24.5	1.0	0.0
Emmetsburg	0.0	0.0	6.0	64.2	28.4	1.5	0.0
Esthervill	0.0	1.5	5.8	71.5	19.7	1.5	0.0
Fairfiel	0.0	0.0	2.8	69.0	26.8	1.4	0.0
Forest City	0.0	0.7	4.1	64.1	29.7	1.4	0.0
Fort Dodge	0.2	2.5	7.3	64.4	24.2	1.4	0.0
Ft Madison	0.0	3.5	4.7	61.8	28.8	1.2	0.0
Garner	0.0	0.0	9.9	64.2	25.9	0.0	0.0
Glenwood	0.0	1.6	9.8	70.5	18.0	0.0	0.0
Grand Junction	0.0	0.0	57.1	28.6	14.3	0.0	0.0
Greenfield	0.0	0.0	5.5	60.0	34.5	0.0	0.0
Grinnell	0.0	3.5	5.8	67.4	22.1	1.2	0.0
Grundy Ctr	0.0	0.0	0.0	68.0	32.0	0.0	0.0
Guthrie Ct	0.0	1.9	7.5	56.6	34.0	0.0	0.0
Hampton	0.0	4.3	3.6	52.9	35.5	3.6	0.0
Harlan	0.0	2.8	3.8	67.9	23.6	1.9	0.0
Hawarden	0.0	3.4	3.4	54.2	37.3	1.7	0.0
Hispanic Cln	0.0	0.0	9.1	36.4	50.0	4.5	0.0
Humboldt	0.0	1.5	5.9	52.9	39.7	0.0	0.0
ILH	0.2	2.8	7.1	55.7	31.4	2.8	0.1
Ida Grove	0.0	1.3	5.1	64.1	26.9	2.6	0.0
Independen	0.0	3.1	1.0	61.2	30.6	4.1	0.0
Indianola	0.0	2.9	8.7	61.0	25.6	1.7	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Iowa City	0.0	0.4	3.4	51.4	40.8	4.0	0.0
Iowa Falls	0.0	2.2	4.4	64.4	28.9	0.0	0.0
Jamaica	0.0	8.3	0.0	58.3	33.3	0.0	0.0
Jefferson	0.0	0.0	1.8	75.0	21.4	1.8	0.0
KR	0.0	0.0	0.0	45.0	52.5	2.5	0.0
Keokuk	0.0	3.2	10.3	65.4	21.2	0.0	0.0
Keosauqua	0.0	7.4	3.7	48.1	37.0	0.0	3.7
Knoxville	0.0	1.6	7.8	61.6	25.7	3.3	0.0
LCPH	0.0	0.0	3.0	71.7	23.2	2.0	0.0
Lamoni	0.0	0.0	5.6	83.3	11.1	0.0	0.0
Le Mars	0.0	1.4	5.8	61.6	26.8	4.3	0.0
Lenox	0.0	0.0	3.0	69.7	24.2	3.0	0.0
Leon	0.0	0.0	12.2	65.3	14.3	8.2	0.0
Logan	0.0	2.3	4.7	61.6	30.2	1.2	0.0
Manchester	0.0	0.0	4.4	71.1	22.2	2.2	0.0
Manning	0.0	0.0	50.0	50.0	0.0	0.0	0.0
Maquoketa	0.0	1.1	4.4	73.6	20.9	0.0	0.0
Marengo	0.0	4.0	4.0	68.0	24.0	0.0	0.0
Marion	0.2	0.9	4.2	61.2	31.8	1.6	0.0
Marshalltown	0.0	2.8	6.7	58.2	30.0	2.3	0.0
Mason City	0.0	1.2	5.8	65.3	26.1	1.7	0.0
Monticello	0.0	0.0	7.1	53.6	32.1	7.1	0.0
Mt Ayr	0.0	6.5	3.2	71.0	19.4	0.0	0.0
Mt Pleasan	0.0	1.8	6.0	68.1	21.7	2.4	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Mtown Dtown	0.0	0.0	0.0	33.3	66.7	0.0	0.0
Muscatine	0.0	2.7	6.8	62.1	26.2	2.1	0.0
Nevada	0.0	0.0	0.0	46.7	53.3	0.0	0.0
New Hampto	0.0	0.0	8.9	55.4	35.7	0.0	0.0
Newton	0.0	1.5	7.5	62.4	26.7	1.1	0.8
North Towne	0.0	2.2	7.4	58.0	30.8	1.6	0.0
Northwood	0.0	0.0	3.3	73.3	23.3	0.0	0.0
Norwalk	0.0	2.3	2.3	53.5	41.9	0.0	0.0
Oakridge	0.0	0.0	0.0	56.3	43.8	0.0	0.0
Oelwein	0.0	1.2	5.8	61.6	31.4	0.0	0.0
Onawa	1.4	1.4	4.1	61.6	31.5	0.0	0.0
Orange Cty	0.0	2.6	6.6	67.1	19.7	3.9	0.0
Osage	0.0	1.2	1.2	61.6	36.0	0.0	0.0
Osceola	0.0	1.3	5.1	57.1	34.6	1.9	0.0
Oskaloosa	0.0	2.0	4.0	64.0	26.7	3.3	0.0
Ottumwa	0.0	2.3	7.0	63.6	25.0	2.1	0.1
Pella	0.0	4.4	2.2	57.8	31.1	4.4	0.0
Perry	1.2	1.2	4.8	60.6	30.3	1.8	0.0
Pocahontas	0.0	0.0	0.0	73.9	26.1	0.0	0.0
Pomeroy	0.0	0.0	0.0	100.0	0.0	0.0	0.0
Postville	0.8	2.5	5.7	46.7	42.6	1.6	0.0
Red Oak	0.0	2.6	9.6	58.8	28.9	0.0	0.0
Redfield	0.0	0.0	13.3	60.0	26.7	0.0	0.0
Refugee Cln	0.0	3.8	0.0	34.6	53.8	7.7	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Rock Rapid	0.0	1.9	1.9	52.8	37.7	5.7	0.0
Rock Valle	0.0	4.6	7.3	60.6	25.7	1.8	0.0
Rockwell C	0.0	0.0	0.0	74.4	20.9	4.7	0.0
SDM	0.0	1.9	6.8	59.7	29.2	2.5	0.0
Sac City	0.0	2.7	2.7	56.8	37.8	0.0	0.0
Sheldon	0.0	4.5	4.5	56.4	33.6	0.9	0.0
Shenandoah	0.0	7.6	6.3	63.3	20.3	2.5	0.0
Sibley	0.0	0.0	2.5	55.0	40.0	2.5	0.0
Sidney	0.0	0.0	4.3	56.5	39.1	0.0	0.0
Sigourney	0.0	0.0	2.8	63.9	33.3	0.0	0.0
Sioux Cent	0.0	5.4	13.6	47.6	29.9	2.7	0.7
Sioux City	0.1	2.0	6.4	58.9	30.8	1.9	0.0
Spencer	0.0	0.5	6.7	72.5	20.2	0.0	0.0
Spirit Lak	0.0	1.4	5.5	68.5	20.5	4.1	0.0
Storm Lake	0.0	2.9	3.8	56.8	31.4	5.1	0.0
Stuart	0.0	0.0	3.8	53.8	42.3	0.0	0.0
Tama	0.0	1.4	2.9	69.6	24.6	1.4	0.0
Tama/Settl	0.0	0.0	0.0	50.0	33.3	16.7	0.0
Tipton	0.0	0.0	8.3	66.7	20.8	4.2	0.0
Traer	0.0	0.0	0.0	91.7	8.3	0.0	0.0
UIHC	0.0	3.3	6.7	56.7	33.3	0.0	0.0
Vinton	0.0	0.0	2.7	66.7	26.7	4.0	0.0
WDM	0.0	0.0	3.9	55.1	36.1	4.9	0.0
Wapello	0.0	0.0	0.0	0.0	100.0	0.0	0.0

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 4A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Combined 3 Years

Clinic	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Washington	0.0	0.9	3.7	62.0	33.3	0.0	0.0
Waterloo	0.1	1.7	5.3	61.9	29.0	1.9	0.1
Waukee	0.0	4.6	4.6	53.7	34.3	2.8	0.0
Waukon	0.0	0.0	1.4	79.7	18.8	0.0	0.0
Waverly	0.0	0.0	3.2	58.7	33.3	4.8	0.0
Wbstr Ct	0.0	3.1	5.8	63.4	25.7	2.1	0.0
West Union	0.0	2.1	6.3	64.6	22.9	4.2	0.0
Williamsburg	0.0	0.0	0.0	0.0	100.0	0.0	0.0
Winterset	0.0	0.0	4.8	62.9	30.6	1.6	0.0
Iowa	**0.1**	**2.1**	**6.1**	**60.4**	**29.1**	**2.2**	**0.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Ackley		0.0 (1)	45.5 (10)		27.3 (114)	45.5 (80)		66.7 (153)		66.7 (151)		0.0 (1)		0.0 (1)
Adel		0.0 (1)	77.8 (138)		17.6 (28)	47.1 (90)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Albia		7.3 (145)	41.5 (9)		34.1 (138)	34.1 (24)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Algona		5.5 (129)	68.1 (109)		23.9 (90)	44.3 (70)		6.3 (77)		13.6 (93)		100.0 (26)		100.0 (36)
Allen WHC		3.0 (73)	55.3 (21)		21.9 (63)	46.2 (87)		0.0 (1)		8.3 (83)		100.0 (26)		100.0 (36)
Allison		1.9 (56)	81.1 (143)		18.9 (37)	52.8 (122)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Altoona		5.3 (127)	64.0 (84)		38.9 (146)	33.3 (19)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Ames		4.7 (113)	57.8 (28)		27.1 (112)	41.4 (50)		5.2 (75)		12.0 (91)		100.0 (26)		100.0 (36)
Anamosa		1.5 (51)	58.2 (32)		20.0 (42)	53.8 (130)		40.0 (135)		20.0 (105)		100.0 (26)		100.0 (36)
Ankeny		6.1 (136)	62.2 (69)		18.5 (34)	47.8 (92)		2.7 (62)		0.0 (1)		100.0 (26)		100.0 (36)
Atlantic		4.4 (105)	65.6 (97)		23.9 (90)	60.2 (144)		4.8 (73)		6.3 (79)		0.0 (1)		100.0 (36)
Audubon		0.0 (1)	64.5 (89)		16.7 (24)	50.0 (106)		30.0 (118)		30.8 (123)		100.0 (26)		100.0 (36)
Avoca		0.0 (1)	60.0 (46)		33.3 (134)	33.3 (19)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Bedford		11.1 (149)	66.7 (104)		33.3 (134)	27.8 (9)		33.3 (123)		33.3 (125)		100.0 (26)		0.0 (1)
Belle Plaine		0.0 (1)	84.2 (148)		36.8 (144)	36.8 (33)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Bloomfield		5.3 (127)	57.9 (30)		38.9 (146)	44.4 (72)		60.0 (148)		0.0 (1)		0.0 (1)		0.0 (1)
Boone		0.0 (1)	66.1 (101)		29.8 (126)	49.1 (103)		15.4 (97)		0.0 (1)		100.0 (26)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Brdlwns RP		4.4 (105)	54.1 (19)		23.7 (88)	42.3 (59)		2.5 (60)		5.1 (73)		100.0 (26)		100.0 (36)
Brooklyn		0.0 (1)	81.3 (144)		33.3 (134)	60.0 (140)		50.0 (143)		50.0 (144)		0.0 (1)		0.0 (1)
Burlington		5.1 (124)	57.8 (28)		15.8 (20)	52.7 (121)		6.8 (81)		4.3 (70)		100.0 (26)		100.0 (36)
CHC		4.6 (111)	62.0 (67)		22.1 (65)	48.4 (98)		38.2 (133)		37.3 (134)		100.0 (26)		100.0 (36)
CR Urban		3.6 (86)	60.1 (51)		23.0 (80)	48.1 (94)		40.6 (137)		32.9 (124)		100.0 (26)		100.0 (36)
Carlisle		0.0 (1)	61.1 (62)		22.2 (66)	38.9 (42)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (36)
Carroll		4.7 (113)	65.3 (95)		18.4 (33)	53.5 (129)		25.4 (112)		33.3 (125)		100.0 (26)		100.0 (36)
Cedar Falls		13.0 (151)	69.6 (113)		13.0 (16)	60.9 (145)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Centervill		3.0 (73)	76.1 (136)		27.7 (117)	53.8 (130)		50.0 (143)		20.0 (105)		100.0 (26)		100.0 (36)
Chariton		3.9 (96)	58.8 (35)		37.5 (145)	39.6 (44)		38.5 (134)		14.3 (94)		100.0 (26)		100.0 (36)
Charles City		2.2 (60)	64.2 (85)		22.7 (73)	37.1 (34)		9.5 (86)		3.1 (65)		100.0 (26)		100.0 (36)
Cherokee		7.5 (146)	62.5 (71)		21.5 (60)	44.3 (70)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Clarinda		4.5 (108)	63.6 (82)		25.0 (97)	45.0 (77)		37.5 (130)		26.7 (119)		0.0 (1)		100.0 (36)
Clarion		2.0 (58)	66.3 (102)		31.3 (129)	36.4 (29)		3.8 (68)		0.0 (1)		100.0 (26)		100.0 (36)
Clinton		4.3 (104)	64.5 (89)		21.9 (63)	56.0 (136)		26.5 (115)		25.6 (118)		100.0 (26)		100.0 (36)
Colfax		15.8 (152)	68.4 (111)		21.1 (52)	21.1 (4)		12.5 (90)		66.7 (151)		100.0 (26)		0.0 (1)
Columbus J		2.9 (69)	34.3 (8)		34.3 (139)	31.4 (14)		0.0 (1)		0.0 (1)		100.0 (26)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Corning		4.2 (101)	62.5 (71)		25.0 (97)	45.8 (83)		62.5 (152)		28.6 (120)		100.0 (26)		100.0 (36)
Corydon		2.7 (66)	83.8 (147)		36.1 (143)	50.0 (106)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (36)
Council Blfs		3.8 (93)	63.1 (78)		22.8 (76)	50.1 (113)		8.4 (84)		5.7 (75)		100.0 (26)		100.0 (36)
Cresco		6.8 (143)	57.6 (27)		21.8 (61)	43.6 (64)		14.3 (95)		25.0 (116)		100.0 (26)		100.0 (36)
Creston		5.2 (126)	56.8 (24)		22.7 (73)	49.3 (104)		22.5 (108)		17.2 (100)		100.0 (26)		100.0 (36)
Dallas Ctr		0.0 (1)	25.0 (5)		0.0 (1)	75.0 (148)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Dayton		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Decorah		3.3 (79)	60.0 (46)		5.0 (11)	51.7 (116)		17.6 (99)		33.3 (125)		100.0 (26)		100.0 (36)
Denison		2.5 (64)	63.4 (81)		36.0 (142)	30.1 (11)		2.8 (64)		3.8 (66)		100.0 (26)		100.0 (36)
Dewitt		4.5 (108)	77.3 (137)		35.0 (140)	40.0 (46)		60.0 (148)		0.0 (1)		100.0 (26)		100.0 (36)
Dubuque		3.7 (89)	62.6 (75)		22.3 (68)	47.9 (93)		50.4 (146)		39.3 (138)		100.0 (26)		100.0 (36)
Dyersville		0.0 (1)	83.3 (146)		18.2 (31)	36.4 (29)		33.3 (123)		0.0 (1)		100.0 (26)		0.0 (1)
Eagle Grove		1.7 (53)	54.2 (20)		28.8 (121)	33.9 (23)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Edgerton		3.5 (84)	58.8 (35)		20.4 (49)	48.4 (98)		30.6 (120)		19.4 (103)		100.0 (26)		100.0 (36)
Eldora		0.0 (1)	46.7 (11)		7.1 (12)	35.7 (26)		0.0 (1)		0.0 (1)		100.0 (26)		0.0 (1)
Elkader		1.3 (48)	71.8 (121)		19.2 (39)	53.4 (127)		20.0 (100)		22.7 (113)		100.0 (26)		100.0 (36)
Emmetsburg		0.0 (1)	59.5 (42)		15.8 (20)	50.0 (106)		0.0 (1)		16.7 (99)		100.0 (26)		100.0 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Esthervill		3.1 (76)	65.6 (97)		26.7 (109)	42.2 (57)		4.0 (69)		0.0 (1)		100.0 (26)		100.0 (36)
Fairfiel		0.0 (1)	64.9 (92)		17.0 (26)	50.9 (115)		23.5 (109)		23.1 (114)		0.0 (1)		100.0 (36)
Forest City		5.1 (124)	67.5 (107)		17.0 (26)	50.0 (106)		6.7 (79)		5.9 (76)		100.0 (26)		100.0 (36)
Fort Dodge		2.3 (62)	66.3 (102)		21.8 (61)	48.4 (98)		2.7 (62)		6.3 (79)		100.0 (26)		100.0 (36)
Ft Madison		0.8 (45)	73.0 (124)		18.3 (32)	52.2 (119)		4.3 (70)		0.0 (1)		100.0 (26)		100.0 (36)
Garner		5.0 (121)	66.7 (104)		23.3 (84)	50.0 (106)		12.5 (90)		8.3 (83)		100.0 (26)		100.0 (36)
Glenwood		6.7 (141)	53.3 (15)		23.3 (84)	44.2 (68)		0.0 (1)		5.9 (76)		0.0 (1)		100.0 (36)
Grand Junction		0.0 (1)	28.6 (6)		0.0 (1)	33.3 (19)		0.0 (1)		50.0 (144)		100.0 (26)		0.0 (1)
Greenfield		2.4 (63)	73.8 (126)		20.0 (42)	37.5 (35)		20.0 (100)		0.0 (1)		100.0 (26)		100.0 (36)
Grinnell		5.8 (134)	68.1 (109)		15.6 (18)	60.9 (145)		29.4 (117)		33.3 (125)		100.0 (26)		100.0 (36)
Grundy Ctr		0.0 (1)	92.3 (151)		30.8 (128)	30.8 (12)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (36)
Guthrie Ct		0.0 (1)	75.7 (135)		28.6 (119)	54.3 (132)		25.0 (110)		14.3 (94)		100.0 (26)		100.0 (36)
Hampton		1.7 (53)	78.3 (139)		27.5 (115)	45.0 (77)		0.0 (1)		4.3 (70)		100.0 (26)		100.0 (36)
Harlan		4.9 (120)	67.9 (108)		21.0 (51)	51.9 (117)		4.5 (72)		6.5 (81)		100.0 (26)		100.0 (36)
Hawarden		4.0 (98)	74.0 (129)		23.4 (87)	53.2 (125)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Hispanic Cln		0.0 (1)	72.7 (123)		45.5 (150)	27.3 (7)		0.0 (1)		33.3 (125)		100.0 (26)		0.0 (1)
Humboldt		0.0 (1)	64.4 (88)		32.8 (133)	37.9 (37)		0.0 (1)		0.0 (1)		100.0 (26)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
ILH		4.8 (118)	59.7 (44)		26.3 (108)	40.5 (48)		1.9 (58)		3.8 (66)		100.0 (26)		100.0 (36)
Ida Grove		6.0 (135)	67.2 (106)		35.4 (141)	44.6 (75)		0.0 (1)		0.0 (1)		100.0 (26)		0.0 (1)
Independen		0.0 (1)	72.2 (122)		25.7 (105)	35.7 (26)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Indianola		6.6 (140)	60.3 (54)		29.4 (124)	42.0 (56)		6.5 (78)		0.0 (1)		100.0 (26)		100.0 (36)
Iowa City		3.3 (79)	60.1 (51)		22.3 (68)	41.7 (52)		31.1 (122)		22.4 (112)		100.0 (26)		100.0 (36)
Iowa Falls		5.7 (131)	65.7 (99)		32.4 (132)	32.4 (18)		12.5 (90)		18.2 (101)		100.0 (26)		100.0 (36)
Jamaica		10.0 (148)	90.0 (149)		10.0 (14)	80.0 (149)		33.3 (123)		0.0 (1)		100.0 (26)		100.0 (36)
Jefferson		6.5 (139)	73.9 (128)		19.6 (41)	54.3 (132)		42.9 (139)		8.3 (83)		100.0 (26)		100.0 (36)
KR		3.2 (78)	58.1 (31)		26.7 (109)	36.7 (32)		0.0 (1)		20.0 (105)		100.0 (26)		100.0 (36)
Keokuk		6.7 (141)	65.0 (93)		23.0 (80)	35.4 (25)		0.0 (1)		11.1 (90)		100.0 (26)		100.0 (36)
Keosauqua		0.0 (1)	59.1 (38)		22.7 (73)	36.4 (29)		25.0 (110)		0.0 (1)		0.0 (1)		100.0 (36)
Knoxville		4.6 (111)	61.3 (64)		21.1 (52)	42.2 (57)		2.6 (61)		9.5 (88)		100.0 (26)		100.0 (36)
LCPH		0.0 (1)	58.8 (35)		22.4 (70)	55.2 (135)		58.3 (147)		33.3 (125)		100.0 (26)		100.0 (36)
Lamoni		0.0 (1)	64.3 (86)		25.0 (97)	50.0 (106)		0.0 (1)		50.0 (144)		100.0 (26)		0.0 (1)
Le Mars		4.7 (113)	66.0 (100)		21.2 (56)	44.2 (68)		0.0 (1)		4.0 (68)		100.0 (26)		0.0 (1)
Lenox		4.8 (118)	57.1 (26)		33.3 (134)	38.1 (40)		60.0 (148)		50.0 (144)		100.0 (26)		0.0 (1)
Leon		3.1 (76)	62.5 (71)		51.6 (152)	25.8 (6)		10.0 (87)		0.0 (1)		100.0 (26)		100.0 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Logan		1.6 (52)	59.4 (40)		19.4 (40)	50.0 (106)		0.0 (1)		0.0 (1)		0.0 (1)		100.0 (36)
Manchester		2.8 (68)	75.0 (132)		25.0 (97)	44.4 (72)		44.4 (142)		37.5 (135)		100.0 (26)		100.0 (36)
Manning		0.0 (1)	50.0 (12)		0.0 (1)	100.0 (150)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Maquoketa		7.0 (144)	73.2 (125)		21.1 (52)	50.7 (114)		20.0 (100)		42.9 (141)		100.0 (26)		100.0 (36)
Marengo		6.3 (138)	75.0 (132)		20.0 (42)	40.0 (46)		20.0 (100)		12.5 (92)		100.0 (26)		0.0 (1)
Marion		1.8 (55)	65.0 (93)		20.6 (50)	48.9 (102)		30.5 (119)		19.5 (104)		100.0 (26)		100.0 (36)
Marshalltown		3.6 (86)	59.7 (44)		24.8 (96)	39.2 (43)		13.8 (94)		16.2 (98)		100.0 (26)		100.0 (36)
Mason City		3.8 (93)	64.3 (86)		18.6 (35)	47.5 (91)		3.2 (67)		8.3 (83)		100.0 (26)		100.0 (36)
Monticello		4.2 (101)	58.3 (33)		25.0 (97)	41.7 (52)		60.0 (148)		0.0 (1)		100.0 (26)		100.0 (36)
Mt Ayr		3.7 (89)	51.9 (13)		29.6 (125)	48.1 (94)		22.2 (105)		20.0 (105)		100.0 (26)		0.0 (1)
Mt Pleasan		0.8 (45)	61.5 (66)		23.1 (82)	46.2 (87)		7.0 (82)		0.0 (1)		100.0 (26)		100.0 (36)
Mtown Dtown		33.3 (153)	0.0 (1)		100.0 (153)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Muscatine		3.5 (84)	60.1 (51)		21.3 (58)	48.2 (96)		1.5 (57)		7.3 (82)		100.0 (26)		100.0 (36)
Nevada		0.0 (1)	81.8 (145)		0.0 (1)	54.5 (134)		0.0 (1)		0.0 (1)		100.0 (26)		0.0 (1)
New Hampto		4.2 (101)	60.4 (57)		25.0 (97)	39.6 (44)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Newton		5.0 (121)	60.8 (61)		24.7 (95)	44.0 (67)		26.2 (114)		37.5 (135)		100.0 (26)		100.0 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
North Towne		2.2 (60)	62.7 (76)		18.7 (36)	53.4 (127)		34.1 (128)		43.4 (142)		100.0 (26)		100.0 (36)
Northwood		0.0 (1)	75.0 (132)		0.0 (1)	52.6 (120)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Norwalk		0.0 (1)	60.0 (46)		20.0 (42)	60.0 (140)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Oakridge		0.0 (1)	33.3 (7)		44.4 (149)	44.4 (72)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Oelwein		1.4 (49)	62.3 (70)		17.9 (29)	49.3 (104)		37.5 (130)		35.3 (131)		100.0 (26)		100.0 (36)
Onawa		3.6 (86)	60.7 (60)		16.4 (23)	41.8 (55)		0.0 (1)		0.0 (1)		100.0 (26)		0.0 (1)
Orange Cty		3.7 (89)	53.7 (16)		23.1 (82)	19.2 (3)		6.7 (79)		0.0 (1)		100.0 (26)		100.0 (36)
Osage		0.0 (1)	73.8 (126)		16.7 (24)	53.3 (126)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Osceola		6.1 (136)	60.5 (59)		30.3 (127)	35.8 (28)		41.2 (138)		46.7 (143)		100.0 (26)		100.0 (36)
Oskaloosa		4.5 (108)	58.6 (34)		23.3 (84)	41.7 (52)		30.8 (121)		28.6 (120)		0.0 (1)		100.0 (36)
Ottumwa		4.7 (113)	59.2 (39)		24.3 (94)	43.8 (65)		28.6 (116)		24.8 (115)		100.0 (26)		100.0 (36)
Pella		5.7 (131)	60.0 (46)		11.4 (15)	57.1 (139)		10.0 (87)		14.3 (94)		0.0 (1)		100.0 (36)
Perry		4.4 (105)	61.3 (64)		25.8 (106)	37.9 (37)		2.8 (64)		0.0 (1)		100.0 (26)		100.0 (36)
Pocahontas		5.0 (121)	60.0 (46)		20.0 (42)	25.0 (5)		16.7 (98)		0.0 (1)		0.0 (1)		100.0 (36)
Pomeroy		0.0 (1)	100.0 (152)		0.0 (1)	100.0 (150)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Postville		2.0 (58)	61.2 (63)		32.3 (131)	38.5 (41)		43.8 (141)		40.6 (140)		100.0 (26)		100.0 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Red Oak		3.3 (79)	71.4 (117)		27.6 (116)	46.0 (85)		0.0 (1)		5.6 (74)		100.0 (26)		100.0 (36)
Redfield		0.0 (1)	71.4 (117)		21.4 (59)	42.9 (60)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Refugee Cln		0.0 (1)	71.4 (117)		19.0 (38)	28.6 (10)		40.0 (135)		50.0 (144)		100.0 (26)		0.0 (1)
Rock Rapid		3.0 (73)	63.6 (82)		26.7 (109)	56.7 (138)		0.0 (1)		0.0 (1)		100.0 (26)		0.0 (1)
Rock Valle		3.7 (89)	53.7 (16)		22.5 (71)	37.5 (35)		0.0 (1)		2.9 (63)		100.0 (26)		100.0 (36)
Rockwell C		2.9 (69)	55.9 (22)		24.2 (93)	45.5 (80)		9.1 (85)		30.0 (122)		100.0 (26)		100.0 (36)
SDM		4.7 (113)	60.4 (57)		23.7 (88)	45.3 (79)		2.1 (59)		2.7 (62)		100.0 (26)		100.0 (36)
Sac City		5.7 (131)	74.3 (131)		22.9 (78)	48.6 (101)		22.2 (105)		20.0 (105)		100.0 (26)		100.0 (36)
Sheldon		3.8 (93)	62.0 (67)		28.8 (121)	52.1 (118)		0.0 (1)		5.0 (72)		100.0 (26)		0.0 (1)
Shenandoah		3.3 (79)	71.7 (120)		22.8 (76)	45.6 (82)		35.3 (129)		40.0 (139)		100.0 (26)		100.0 (36)
Sibley		0.0 (1)	80.0 (141)		21.1 (52)	31.6 (16)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Sidney		0.0 (1)	64.7 (91)		20.0 (42)	60.0 (140)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Sigourney		0.0 (1)	59.4 (40)		24.1 (92)	31.0 (13)		33.3 (123)		25.0 (116)		0.0 (1)		0.0 (1)
Sioux Cent		0.9 (47)	53.7 (16)		41.9 (148)	31.4 (14)		5.8 (76)		2.9 (63)		100.0 (26)		100.0 (36)
Sioux City		2.9 (69)	62.5 (71)		22.6 (72)	42.9 (60)		5.1 (74)		5.9 (76)		100.0 (26)		100.0 (36)
Spencer		2.6 (65)	59.6 (43)		29.2 (123)	41.6 (51)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Spirit Lak		1.9 (56)	71.2 (116)		18.0 (30)	56.0 (136)		0.0 (1)		22.2 (111)		100.0 (26)		100.0 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Storm Lake		3.3 (79)	63.1 (78)		25.6 (104)	41.0 (49)		1.1 (56)		14.3 (94)		100.0 (26)		100.0 (36)
Stuart		0.0 (1)	68.4 (111)		15.8 (20)	63.2 (147)		22.2 (105)		62.5 (150)		100.0 (26)		0.0 (1)
Tama		2.7 (66)	69.6 (113)		25.2 (103)	44.7 (76)		25.9 (113)		36.4 (132)		100.0 (26)		100.0 (36)
Tama/Settl		0.0 (1)	80.0 (141)		0.0 (1)	60.0 (140)		0.0 (1)		100.0 (153)		100.0 (26)		0.0 (1)
Tipton		11.8 (150)	52.9 (14)		46.7 (151)	33.3 (19)		33.3 (123)		0.0 (1)		100.0 (26)		100.0 (36)
Traer		0.0 (1)	90.9 (150)		9.1 (13)	27.3 (7)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
UIHC		0.0 (1)	14.3 (4)		28.6 (119)	42.9 (60)		50.0 (143)		50.0 (144)		100.0 (26)		100.0 (36)
Vinton		0.0 (1)	63.1 (78)		28.1 (118)	46.9 (89)		12.5 (90)		0.0 (1)		100.0 (26)		100.0 (36)
WDM		4.0 (98)	65.3 (95)		27.1 (112)	43.8 (65)		4.4 (71)		8.3 (83)		100.0 (26)		100.0 (36)
Wapello		0.0 (1)	0.0 (1)		0.0 (1)	100.0 (150)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Washington		8.9 (147)	57.0 (25)		22.2 (66)	43.1 (63)		20.8 (104)		10.0 (89)		100.0 (26)		100.0 (36)
Waterloo		3.9 (96)	60.3 (54)		21.2 (56)	48.3 (97)		7.2 (83)		4.1 (69)		100.0 (26)		100.0 (36)
Waukee		5.5 (129)	60.3 (54)		13.2 (17)	52.9 (123)		0.0 (1)		0.0 (1)		100.0 (26)		100.0 (36)
Waukon		0.0 (1)	70.4 (115)		26.0 (107)	38.0 (39)		42.9 (139)		21.4 (110)		100.0 (26)		100.0 (36)
Waverly		4.0 (98)	74.0 (129)		20.0 (42)	46.0 (85)		11.1 (89)		0.0 (1)		100.0 (26)		100.0 (36)
Wbstr Ct		1.4 (49)	56.1 (23)		22.9 (78)	45.8 (83)		2.9 (66)		18.2 (101)		100.0 (26)		100.0 (36)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Health Indicators by Clinic

Combined 3 Years

Comparison of Maternal Health Indicators (2,3)

Clinic	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
		Under-weight	Over-weight		< Ideal	> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
West Union		2.9 (69)	62.9 (77)		15.6 (18)	53.1 (124)		14.3 (95)		36.4 (132)		100.0 (26)		100.0 (36)
Williamsburg		0.0 (1)	100.0 (152)		0.0 (1)	100.0 (150)		0.0 (1)		0.0 (1)		0.0 (1)		0.0 (1)
Winterset		0.0 (1)	78.6 (140)		31.7 (130)	31.7 (17)		37.5 (130)		38.5 (137)		100.0 (26)		100.0 (36)
Iowa		**3.8**	**61.2**		**23.2**	**45.2**		**14.7**		**14.2**		**100.0**		**100.0**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley		54.5 (7)	9.1 (74)		0.0 (67)		30.0 (39)		90.9 (87)
Adel		35.3 (45)	5.9 (43)		0.0 (67)		45.5 (16)		100.0 (1)
Albia		24.3 (117)	13.5 (114)		0.0 (67)		23.8 (71)		86.4 (119)
Algona		48.4 (10)	11.0 (96)		0.0 (67)		18.5 (106)		96.6 (42)
Allen WHC		27.7 (93)	31.0 (151)		0.9 (51)		22.4 (81)		87.5 (110)
Allison		38.5 (33)	3.8 (33)		0.0 (67)		27.9 (48)		86.3 (120)
Altoona		32.8 (62)	9.4 (81)		1.4 (41)		21.2 (85)		97.3 (38)
Ames		33.5 (56)	12.3 (104)		1.9 (28)		24.6 (67)		90.0 (98)
Anamosa		27.9 (92)	6.6 (48)		1.4 (41)		34.4 (25)		90.9 (87)
Ankeny		30.2 (74)	7.0 (56)		0.8 (54)		16.2 (116)		93.2 (69)
Atlantic		29.7 (79)	6.8 (52)		0.0 (67)		20.0 (96)		100.0 (1)
Audubon		33.3 (57)	7.4 (59)		0.0 (67)		33.3 (27)		100.0 (1)
Avoca		14.3 (140)	35.7 (153)		0.0 (67)		15.4 (119)		84.6 (131)
Bedford		13.3 (141)	0.0 (1)		0.0 (67)		20.0 (96)		80.0 (138)
Belle Plaine		20.0 (130)	0.0 (1)		0.0 (67)		16.7 (113)		85.7 (122)
Bloomfield		38.5 (33)	15.4 (121)		7.1 (1)		0.0 (141)		100.0 (1)
Boone		25.0 (105)	11.4 (100)		3.6 (7)		30.8 (37)		96.3 (44)
Brdlwns RP		27.0 (97)	12.6 (106)		0.7 (58)		12.2 (128)		87.2 (113)
Brooklyn		36.4 (42)	27.3 (148)		0.0 (67)		50.0 (7)		100.0 (1)
Burlington		32.2 (63)	3.9 (34)		1.0 (47)		21.8 (82)		94.2 (61)
CHC		28.0 (90)	19.9 (136)		1.6 (38)		10.0 (134)		84.8 (130)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
CR Urban		24.9 (111)	9.9 (87)		0.6 (62)		13.0 (127)		88.4 (107)
Carlisle		25.0 (105)	25.0 (145)		0.0 (67)		30.0 (39)		100.0 (1)
Carroll		25.9 (104)	7.6 (62)		0.5 (65)		21.3 (83)		92.2 (77)
Cedar Falls		29.2 (81)	16.7 (125)		0.0 (67)		46.7 (15)		85.0 (126)
Centervill		30.8 (69)	17.3 (129)		3.2 (13)		40.0 (20)		100.0 (1)
Chariton		39.6 (28)	6.3 (46)		0.0 (67)		27.8 (49)		72.2 (146)
Charles City		45.4 (13)	12.8 (108)		0.0 (67)		21.0 (87)		90.8 (94)
Cherokee		39.7 (27)	2.9 (29)		2.2 (23)		29.2 (44)		96.8 (41)
Clarinda		43.2 (17)	2.3 (27)		4.7 (4)		38.9 (21)		94.1 (62)
Clarion		20.6 (129)	22.7 (143)		1.0 (47)		10.2 (133)		80.0 (138)
Clinton		40.5 (24)	8.3 (69)		0.8 (54)		38.9 (21)		87.0 (116)
Colfax		26.3 (102)	15.8 (123)		0.0 (67)		33.3 (27)		100.0 (1)
Columbus J		25.0 (105)	11.1 (97)		0.0 (67)		16.1 (117)		81.3 (136)
Corning		35.0 (49)	0.0 (1)		0.0 (67)		0.0 (141)		100.0 (1)
Corydon		39.5 (29)	5.3 (39)		0.0 (67)		26.7 (53)		86.7 (117)
Council Blfs		36.7 (40)	6.1 (44)		0.6 (62)		21.0 (87)		87.4 (112)
Cresco		42.1 (19)	7.0 (56)		0.0 (67)		23.0 (79)		90.2 (96)
Creston		43.9 (15)	9.8 (85)		1.3 (43)		30.9 (36)		94.3 (60)
Dallas Ctr		0.0 (147)	0.0 (1)		0.0 (67)		0.0 (141)		0.0 (149)
Dayton		0.0 (147)	0.0 (1)		0.0 (67)		0.0 (141)		0.0 (149)
Decorah		22.4 (126)	10.3 (90)		0.0 (67)		18.9 (104)		90.9 (87)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Denison		24.9 (111)	7.7 (63)		0.0 (67)		23.3 (76)		96.4 (43)
Dewitt		40.9 (22)	4.5 (36)		0.0 (67)		55.6 (5)		84.2 (133)
Dubuque		36.6 (41)	13.1 (110)		3.6 (7)		15.7 (118)		91.2 (85)
Dyersville		38.5 (33)	7.7 (63)		0.0 (67)		23.1 (77)		100.0 (1)
Eagle Grove		26.7 (98)	18.3 (132)		0.0 (67)		19.2 (102)		93.1 (70)
Edgerton		33.2 (60)	10.6 (93)		2.0 (26)		12.1 (129)		93.1 (70)
Eldora		58.3 (6)	0.0 (1)		0.0 (67)		37.5 (24)		88.9 (104)
Elkader		23.8 (121)	10.0 (88)		1.2 (45)		33.7 (26)		97.6 (36)
Emmetsburg		39.5 (29)	14.0 (119)		0.0 (67)		65.9 (2)		95.6 (49)
Esthervill		29.8 (77)	7.4 (59)		0.0 (67)		43.8 (17)		92.7 (74)
Fairfiel		31.0 (67)	19.0 (134)		1.7 (33)		33.3 (27)		100.0 (1)
Forest City		35.3 (45)	9.5 (82)		0.9 (51)		20.4 (92)		85.7 (122)
Fort Dodge		28.6 (87)	20.8 (138)		2.4 (20)		17.1 (111)		85.9 (121)
Ft Madison		28.2 (89)	1.6 (25)		0.0 (67)		16.7 (113)		94.5 (58)
Garner		40.0 (25)	21.7 (141)		0.0 (67)		24.1 (70)		82.8 (134)
Glenwood		17.6 (137)	2.9 (29)		1.9 (28)		25.0 (60)		87.5 (110)
Grand Junction		66.7 (3)	0.0 (1)		0.0 (67)		20.0 (96)		80.0 (138)
Greenfield		28.9 (83)	7.9 (66)		0.0 (67)		18.5 (106)		91.7 (81)
Grinnell		24.6 (114)	14.8 (120)		2.9 (16)		10.0 (134)		85.4 (125)
Grundy Ctr		7.7 (144)	15.4 (121)		0.0 (67)		50.0 (7)		100.0 (1)
Guthrie Ct		45.5 (11)	9.1 (74)		0.0 (67)		0.0 (141)		77.8 (142)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Hampton		29.8 (77)	33.0 (152)		1.9 (28)		28.0 (47)		84.9 (129)
Harlan		26.6 (100)	3.1 (31)		0.0 (67)		28.6 (46)		100.0 (1)
Hawarden		34.9 (50)	9.3 (79)		3.6 (7)		26.3 (56)		95.5 (50)
Hispanic Cln		20.0 (130)	5.0 (37)		0.0 (67)		10.0 (134)		94.1 (62)
Humboldt		34.5 (52)	23.6 (144)		3.6 (7)		25.5 (59)		90.9 (87)
ILH		32.0 (64)	11.9 (103)		0.7 (58)		13.8 (123)		88.6 (106)
Ida Grove		38.3 (36)	8.3 (69)		3.2 (13)		22.6 (80)		90.9 (87)
Independen		35.2 (47)	16.7 (125)		0.0 (67)		30.0 (39)		96.2 (45)
Indianola		25.0 (105)	21.4 (140)		0.0 (67)		16.8 (112)		92.3 (75)
Iowa City		18.4 (136)	20.4 (137)		1.7 (33)		20.1 (94)		89.2 (102)
Iowa Falls		41.4 (21)	0.0 (1)		0.0 (67)		19.0 (103)		90.5 (95)
Jamaica		44.4 (14)	0.0 (1)		0.0 (67)		33.3 (27)		100.0 (1)
Jefferson		26.2 (103)	16.7 (125)		0.0 (67)		23.5 (72)		100.0 (1)
KR		25.0 (105)	25.0 (145)		0.0 (67)		47.1 (14)		95.5 (50)
Keokuk		22.1 (128)	2.7 (28)		1.6 (38)		20.4 (92)		94.7 (54)
Keosauqua		36.8 (39)	0.0 (1)		0.0 (67)		33.3 (27)		100.0 (1)
Knoxville		31.3 (66)	12.7 (107)		1.8 (32)		18.1 (109)		94.4 (59)
LCPH		37.7 (38)	6.6 (48)		0.0 (67)		13.3 (125)		94.0 (64)
Lamoni		25.0 (105)	16.7 (125)		0.0 (67)		14.3 (121)		100.0 (1)
Le Mars		20.0 (130)	10.0 (88)		0.0 (67)		33.3 (27)		94.6 (56)
Lenox		40.9 (22)	9.1 (74)		0.0 (67)		54.5 (6)		72.7 (145)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Leon		30.0 (75)	13.3 (112)		0.0 (67)		23.5 (72)		93.8 (65)
Logan		31.5 (65)	1.9 (26)		0.0 (67)		33.3 (27)		80.0 (138)
Manchester		24.3 (117)	8.1 (68)		0.0 (67)		5.7 (140)		100.0 (1)
Manning		100.0 (1)	0.0 (1)		0.0 (67)		0.0 (141)		0.0 (149)
Maquoketa		38.2 (37)	7.4 (59)		0.0 (67)		33.3 (27)		87.1 (115)
Marengo		11.1 (143)	0.0 (1)		5.0 (3)		0.0 (141)		66.7 (148)
Marion		28.4 (88)	12.8 (108)		0.3 (66)		20.6 (91)		93.3 (67)
Marshalltown		38.8 (32)	6.2 (45)		0.7 (58)		14.3 (121)		87.9 (109)
Mason City		39.2 (31)	16.5 (124)		0.9 (51)		21.3 (83)		86.5 (118)
Monticello		50.0 (8)	0.0 (1)		0.0 (67)		23.1 (77)		94.7 (54)
Mt Ayr		33.3 (57)	0.0 (1)		0.0 (67)		25.0 (60)		100.0 (1)
Mt Pleasan		23.5 (123)	6.7 (50)		1.7 (33)		26.7 (53)		87.2 (113)
Mtown Dtown		0.0 (147)	0.0 (1)		0.0 (67)		0.0 (141)		0.0 (149)
Muscatine		33.0 (61)	8.3 (69)		1.7 (33)		17.9 (110)		93.3 (67)
Nevada		45.5 (11)	18.2 (130)		0.0 (67)		42.9 (18)		85.7 (122)
New Hampto		26.5 (101)	11.8 (102)		0.0 (67)		62.1 (3)		94.6 (56)
Newton		24.5 (115)	11.3 (99)		2.0 (26)		13.5 (124)		90.2 (96)
North Towne		24.5 (115)	6.9 (54)		1.0 (47)		20.1 (94)		91.8 (80)
Northwood		50.0 (8)	18.8 (133)		0.0 (67)		23.5 (72)		95.0 (53)
Norwalk		27.3 (95)	9.1 (74)		0.0 (67)		21.1 (86)		85.0 (126)
Oakridge		30.0 (75)	0.0 (1)		0.0 (67)		25.0 (60)		100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Oelwein		27.4 (94)	9.7 (84)		0.0 (67)		26.8 (52)		91.4 (82)
Onawa		23.4 (124)	8.5 (72)		0.0 (67)		10.0 (134)		100.0 (1)
Orange Cty		34.0 (53)	10.6 (93)		3.3 (11)		38.5 (23)		92.9 (73)
Osage		27.3 (95)	18.2 (130)		0.0 (67)		29.1 (45)		89.6 (99)
Osceola		28.0 (90)	9.3 (79)		0.8 (54)		24.6 (67)		95.2 (52)
Oskaloosa		35.5 (44)	8.6 (73)		2.6 (18)		20.0 (96)		92.3 (75)
Ottumwa		28.9 (83)	26.7 (147)		2.1 (24)		20.7 (90)		75.9 (144)
Pella		22.2 (127)	22.2 (142)		0.0 (67)		20.8 (89)		96.0 (47)
Perry		23.3 (125)	12.4 (105)		2.9 (16)		10.5 (131)		93.4 (66)
Pocahontas		43.8 (16)	6.3 (46)		0.0 (67)		26.7 (53)		88.2 (108)
Pomeroy		0.0 (147)	0.0 (1)		0.0 (67)		0.0 (141)		100.0 (1)
Postville		16.0 (138)	21.3 (139)		0.0 (67)		19.3 (101)		84.6 (131)
Red Oak		30.6 (70)	5.6 (41)		1.0 (47)		25.0 (60)		76.5 (143)
Redfield		33.3 (57)	0.0 (1)		0.0 (67)		50.0 (7)		100.0 (1)
Refugee Cln		0.0 (147)	5.0 (37)		0.0 (67)		6.7 (138)		72.2 (146)
Rock Rapid		24.1 (120)	6.9 (54)		0.0 (67)		26.3 (56)		100.0 (1)
Rock Valle		35.1 (48)	13.5 (114)		2.4 (20)		12.0 (130)		81.5 (135)
Rockwell C		40.0 (25)	5.7 (42)		0.0 (67)		33.3 (27)		100.0 (1)
SDM		29.1 (82)	11.4 (100)		0.8 (54)		18.9 (104)		89.0 (103)
Sac City		31.0 (67)	10.3 (90)		0.0 (67)		18.2 (108)		90.9 (87)
Sheldon		26.7 (98)	6.7 (50)		2.4 (20)		48.6 (11)		96.2 (45)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Shenandoah		41.7 (20)	3.3 (32)		3.2 (13)		27.0 (51)		97.1 (40)
Sibley		23.8 (121)	0.0 (1)		3.8 (6)		57.9 (4)		90.9 (87)
Sidney		6.7 (145)	13.3 (112)		5.3 (2)		25.0 (60)		100.0 (1)
Sigourney		30.4 (72)	4.3 (35)		3.3 (11)		16.7 (113)		100.0 (1)
Sioux Cent		11.9 (142)	7.9 (66)		0.0 (67)		5.9 (139)		97.3 (38)
Sioux City		28.9 (83)	7.3 (58)		0.7 (58)		30.2 (38)		93.0 (72)
Spencer		24.8 (113)	9.5 (82)		2.1 (24)		49.3 (10)		91.3 (84)
Spirit Lak		29.4 (80)	9.8 (85)		1.7 (33)		47.8 (13)		100.0 (1)
Storm Lake		19.1 (135)	6.8 (52)		0.0 (67)		41.7 (19)		95.8 (48)
Stuart		5.6 (146)	11.1 (97)		0.0 (67)		30.0 (39)		100.0 (1)
Tama		33.7 (54)	5.4 (40)		4.5 (5)		15.0 (120)		91.4 (82)
Tama/Settl		0.0 (147)	0.0 (1)		0.0 (67)		0.0 (141)		100.0 (1)
Tipton		20.0 (130)	30.0 (149)		0.0 (67)		0.0 (141)		100.0 (1)
Traer		60.0 (5)	0.0 (1)		0.0 (67)		25.0 (60)		100.0 (1)
UIHC		20.0 (130)	0.0 (1)		0.0 (67)		0.0 (141)		100.0 (1)
Vinton		42.3 (18)	13.5 (114)		0.0 (67)		24.3 (69)		97.6 (36)
WDM		33.6 (55)	9.2 (78)		0.6 (62)		10.3 (132)		92.2 (77)
Wapello		0.0 (147)	0.0 (1)		0.0 (67)		100.0 (1)		100.0 (1)
Washington		14.5 (139)	13.2 (111)		0.0 (67)		19.5 (100)		88.9 (104)
Waterloo		30.4 (72)	19.6 (135)		1.5 (40)		23.4 (75)		89.5 (100)
Waukee		35.8 (43)	10.4 (92)		1.2 (45)		25.0 (60)		89.4 (101)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Maternal Behavioral Indicators by Clinic

Combined 3 Years

Comparison of Maternal Behavioral Indicators (2,3)

Clinic	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Waukon		34.6 (51)	7.7 (63)		1.9 (28)		26.3 (56)		91.2 (85)
Waverly		62.2 (4)	13.5 (114)		0.0 (67)		48.5 (12)		100.0 (1)
Wbstr Ct		28.7 (86)	30.7 (150)		1.3 (43)		13.3 (125)		80.8 (137)
West Union		24.3 (117)	10.8 (95)		2.6 (18)		29.7 (43)		91.9 (79)
Williamsburg		100.0 (1)	0.0 (1)		0.0 (67)		0.0 (141)		0.0 (149)
Winterset		30.6 (70)	13.9 (118)		0.0 (67)		27.3 (50)		85.0 (126)
Iowa		**30.3**	**12.3**		**1.1**		**21.0**		**89.9**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indicators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley		18.2 (29)		18.2 (84)		0.0 (125)		0.0 (1)		0.0 (1)
Adel		20.0 (35)		25.0 (120)		0.0 (125)		0.0 (1)		0.0 (1)
Albia		41.2 (126)		33.3 (145)		7.7 (123)		15.8 (143)		0.0 (1)
Algona		33.0 (85)		23.3 (113)		28.1 (59)		5.5 (63)		1.0 (74)
Allen WHC		23.3 (46)		15.5 (65)		32.6 (50)		8.3 (95)		1.8 (91)
Allison		33.9 (97)		23.1 (110)		31.6 (52)		5.5 (63)		1.7 (88)
Altoona		25.5 (50)		9.8 (28)		55.6 (6)		1.9 (38)		1.4 (81)
Ames		24.3 (48)		13.8 (54)		27.5 (62)		8.7 (99)		1.5 (85)
Anamosa		49.1 (141)		27.7 (132)		25.8 (67)		6.7 (77)		4.0 (122)
Ankeny		13.9 (21)		10.4 (32)		25.0 (70)		1.3 (32)		1.0 (74)
Atlantic		43.4 (135)		31.4 (143)		27.3 (63)		5.3 (59)		0.0 (1)
Audubon		32.1 (80)		14.7 (61)		40.0 (20)		14.3 (137)		8.8 (138)
Avoca		37.5 (112)		11.1 (39)		50.0 (9)		12.5 (129)		0.0 (1)
Bedford		70.6 (150)		20.0 (97)		40.0 (20)		17.6 (147)		6.7 (131)
Belle Plaine		31.3 (75)		30.0 (139)		25.0 (70)		6.3 (72)		0.0 (1)
Bloomfield		14.3 (22)		10.0 (29)		50.0 (9)		0.0 (1)		12.5 (143)
Boone		37.3 (111)		31.7 (144)		19.0 (97)		3.8 (50)		0.0 (1)
Brdlwns RP		18.5 (32)		12.3 (44)		25.5 (68)		4.5 (56)		1.4 (81)
Brooklyn		30.8 (72)		15.4 (63)		25.0 (70)		6.7 (77)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Ind cators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Burlington		45.8 (138)		22.2 (107)		40.3 (19)		14.3 (137)		2.8 (112)
CHC		33.8 (96)		18.3 (85)		19.4 (95)		13.3 (133)		5.3 (125)
CR Urban		26.5 (56)		14.4 (59)		32.0 (51)		7.0 (84)		0.9 (72)
Carlisle		16.7 (28)		27.3 (130)		0.0 (125)		0.0 (1)		16.7 (147)
Carroll		38.1 (117)		19.1 (92)		44.3 (16)		5.1 (58)		3.2 (118)
Cedar Falls		12.5 (19)		0.0 (1)		0.0 (125)		7.7 (89)		0.0 (1)
Centervill		42.1 (128)		23.5 (115)		0.0 (125)		12.5 (129)		0.0 (1)
Chariton		25.5 (50)		14.5 (60)		38.5 (27)		5.6 (66)		0.0 (1)
Charles City		33.3 (87)		18.3 (85)		34.0 (40)		4.8 (57)		0.6 (64)
Cherokee		35.1 (104)		21.1 (104)		20.8 (90)		10.8 (118)		5.4 (126)
Clarinda		42.2 (131)		27.3 (130)		27.8 (61)		13.3 (133)		2.6 (104)
Clarion		9.9 (14)		10.9 (38)		27.3 (63)		6.9 (82)		0.9 (72)
Clinton		42.1 (128)		23.7 (116)		28.0 (60)		6.5 (75)		1.1 (77)
Colfax		25.0 (49)		13.6 (53)		0.0 (125)		0.0 (1)		0.0 (1)
Columbus J		0.0 (1)		0.0 (1)		0.0 (125)		0.0 (1)		0.0 (1)
Corning		33.3 (87)		29.6 (138)		0.0 (125)		19.0 (148)		8.0 (137)
Corydon		32.0 (78)		30.8 (142)		11.1 (119)		3.6 (47)		3.7 (121)
Council Blfs		36.7 (109)		23.0 (109)		23.4 (81)		6.2 (71)		0.6 (64)
Cresco		29.7 (66)		23.2 (112)		18.8 (99)		13.6 (135)		7.1 (132)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Ind cators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Creston		31.5 (77)		20.3 (101)		22.7 (83)		7.5 (88)		0.7 (67)
Dallas Ctr		0.0 (1)		20.0 (97)		0.0 (125)		0.0 (1)		0.0 (1)
Dayton		0.0 (1)		0.0 (1)		0.0 (125)		0.0 (1)		0.0 (1)
Decorah		32.8 (83)		15.9 (68)		33.3 (41)		17.2 (146)		11.1 (140)
Denison		11.4 (17)		8.0 (21)		14.3 (108)		1.2 (31)		0.0 (1)
Dewitt		33.3 (87)		26.1 (125)		14.3 (108)		14.3 (137)		0.0 (1)
Dubuque		15.5 (26)		8.3 (22)		22.9 (82)		0.1 (29)		0.0 (1)
Dyersville		35.7 (107)		13.3 (51)		0.0 (125)		0.0 (1)		0.0 (1)
Eagle Grove		20.3 (38)		10.8 (36)		42.9 (18)		3.1 (43)		3.1 (117)
Edgerton		26.9 (59)		16.1 (70)		22.5 (84)		10.9 (119)		1.8 (91)
Eldora		50.0 (142)		58.8 (152)		12.5 (115)		31.3 (152)		11.8 (141)
Elkader		38.8 (119)		20.7 (103)		45.5 (14)		16.9 (144)		12.2 (142)
Emmetsburg		34.8 (102)		23.8 (117)		35.7 (36)		8.7 (99)		14.3 (146)
Esthervill		20.6 (39)		15.0 (62)		22.2 (85)		14.4 (141)		13.7 (145)
Fairfiel		40.8 (125)		18.5 (87)		10.5 (121)		4.1 (52)		0.0 (1)
Forest City		29.7 (66)		12.4 (46)		37.5 (29)		5.3 (59)		0.8 (69)
Fort Dodge		40.7 (124)		21.7 (105)		36.8 (33)		9.5 (111)		2.4 (102)
Ft Madison		42.2 (131)		19.3 (95)		45.1 (15)		8.7 (99)		0.0 (1)
Garner		30.6 (70)		16.7 (75)		36.8 (33)		8.2 (93)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Ind cators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Glenwood		47.2 (139)		16.1 (70)		35.7 (36)		0.0 (1)		0.0 (1)
Grand Junction		71.4 (151)		16.7 (75)		75.0 (2)		0.0 (1)		0.0 (1)
Greenfield		18.6 (33)		10.8 (36)		28.6 (56)		2.4 (41)		2.9 (115)
Grinnell		25.9 (55)		20.6 (102)		25.0 (70)		16.9 (144)		7.6 (135)
Grundy Ctr		6.7 (11)		14.3 (56)		0.0 (125)		0.0 (1)		0.0 (1)
Guthrie Ct		40.0 (121)		28.6 (134)		14.3 (108)		8.6 (98)		2.0 (95)
Hampton		15.3 (24)		13.2 (50)		22.2 (85)		1.8 (37)		0.8 (69)
Harlan		42.0 (127)		27.1 (128)		20.0 (93)		4.3 (55)		0.0 (1)
Hawarden		29.8 (68)		13.5 (52)		38.5 (27)		2.1 (39)		2.7 (109)
Hispanic Cln		0.0 (1)		0.0 (1)		0.0 (125)		0.0 (1)		0.0 (1)
Humboldt		32.2 (81)		18.5 (87)		40.0 (20)		6.8 (80)		1.7 (88)
ILH		26.8 (58)		19.8 (96)		21.6 (89)		5.7 (68)		1.4 (81)
Ida Grove		35.5 (106)		25.9 (123)		9.5 (122)		6.6 (76)		5.5 (127)
Independen		37.7 (114)		26.4 (126)		14.3 (108)		8.3 (95)		1.9 (94)
Indianola		37.9 (116)		17.3 (81)		33.3 (41)		10.3 (115)		2.8 (112)
Iowa City		13.6 (20)		4.4 (15)		37.5 (29)		3.2 (46)		0.3 (63)
Iowa Falls		22.6 (44)		6.7 (17)		16.7 (103)		9.4 (108)		0.0 (1)
Jamaica		50.0 (142)		20.0 (97)		40.0 (20)		12.5 (129)		0.0 (1)
Jefferson		34.1 (98)		16.4 (73)		25.0 (70)		9.1 (104)		1.8 (91)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Ind cators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
KR		15.4 (25)		13.0 (48)		40.0 (20)		11.5 (123)		0.0 (1)
Keokuk		44.4 (136)		25.7 (122)		25.9 (66)		19.0 (148)		2.8 (112)
Keosauqua		30.0 (69)		28.6 (134)		0.0 (125)		15.0 (142)		7.1 (132)
Knoxville		39.2 (120)		23.1 (110)		20.8 (90)		9.1 (104)		2.3 (100)
LCPH		31.3 (75)		10.3 (31)		33.3 (41)		1.5 (35)		0.0 (1)
Lamoni		40.0 (121)		11.1 (39)		25.0 (70)		6.7 (77)		0.0 (1)
Le Mars		21.3 (41)		8.3 (22)		40.0 (20)		1.1 (30)		1.1 (77)
Lenox		14.3 (22)		0.0 (1)		33.3 (41)		0.0 (1)		0.0 (1)
Leon		34.4 (100)		16.2 (72)		18.2 (100)		3.1 (43)		5.6 (128)
Logan		25.5 (50)		17.0 (78)		27.3 (63)		5.5 (63)		0.0 (1)
Manchester		32.4 (82)		15.8 (67)		0.0 (125)		0.0 (1)		0.0 (1)
Manning		0.0 (1)		0.0 (1)		0.0 (125)		0.0 (1)		0.0 (1)
Maquoketa		50.7 (145)		24.3 (118)		23.5 (80)		8.7 (99)		4.0 (122)
Marengo		60.0 (148)		50.0 (150)		28.6 (56)		10.0 (113)		0.0 (1)
Marion		33.0 (85)		18.9 (91)		35.8 (35)		7.2 (87)		1.3 (79)
Marshalltown		16.5 (27)		11.4 (41)		17.7 (101)		5.6 (66)		1.6 (87)
Mason City		33.5 (95)		19.1 (92)		25.3 (69)		6.3 (72)		1.3 (79)
Monticello		27.3 (60)		21.7 (105)		0.0 (125)		9.1 (104)		0.0 (1)
Mt Ayr		45.5 (137)		14.3 (56)		11.1 (119)		9.1 (104)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Ind cators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Mt Pleasan		32.0 (78)		17.4 (82)		31.6 (52)		8.2 (93)		0.7 (67)
Mtown Dtown		0.0 (1)		0.0 (1)		0.0 (125)		0.0 (1)		0.0 (1)
Muscatine		27.4 (61)		16.4 (73)		32.7 (49)		11.5 (123)		2.3 (100)
Nevada		33.3 (87)		7.7 (20)		0.0 (125)		10.0 (113)		0.0 (1)
New Hampto		20.0 (35)		17.1 (80)		25.0 (70)		12.2 (127)		0.0 (1)
Newton		35.8 (108)		27.2 (129)		19.4 (95)		8.0 (91)		1.4 (81)
North Towne		25.6 (54)		13.0 (48)		46.2 (13)		3.7 (49)		2.1 (96)
Northwood		21.1 (40)		7.4 (19)		50.0 (9)		0.0 (1)		0.0 (1)
Norwalk		37.5 (112)		25.9 (123)		0.0 (125)		4.2 (54)		3.3 (120)
Oakridge		20.0 (35)		40.0 (149)		0.0 (125)		0.0 (1)		0.0 (1)
Oelwein		43.3 (134)		28.2 (133)		22.2 (85)		26.8 (151)		21.1 (152)
Onawa		30.6 (70)		28.6 (134)		6.3 (124)		4.1 (52)		0.0 (1)
Orange Cty		11.5 (18)		2.0 (12)		20.0 (93)		3.9 (51)		2.1 (96)
Osage		23.9 (47)		14.1 (55)		37.5 (29)		3.1 (43)		2.6 (104)
Osceola		31.0 (74)		23.4 (114)		15.2 (107)		7.0 (84)		1.0 (74)
Oskaloosa		35.2 (105)		24.3 (118)		13.3 (113)		11.4 (121)		2.9 (115)
Ottumwa		27.5 (62)		17.0 (78)		12.6 (114)		9.4 (108)		2.2 (99)
Pella		33.3 (87)		16.7 (75)		33.3 (41)		3.6 (47)		0.0 (1)
Perry		19.0 (34)		8.7 (25)		35.7 (36)		1.6 (36)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Ind cators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Pocahontas		47.4 (140)		30.4 (140)		22.2 (85)		5.3 (59)		0.0 (1)
Pomeroy		100.0 (152)		0.0 (1)		100.0 (1)		0.0 (1)		0.0 (1)
Postville		8.0 (12)		7.3 (18)		25.0 (70)		6.1 (70)		3.2 (118)
Red Oak		60.8 (149)		35.5 (147)		18.9 (98)		8.1 (92)		0.0 (1)
Redfield		50.0 (142)		50.0 (150)		0.0 (125)		0.0 (1)		25.0 (153)
Refugee Cln		0.0 (1)		0.0 (1)		0.0 (125)		0.0 (1)		0.0 (1)
Rock Rapid		18.2 (29)		9.3 (27)		60.0 (4)		3.0 (42)		0.0 (1)
Rock Valle		4.0 (10)		4.1 (14)		33.3 (41)		1.3 (32)		0.0 (1)
Rockwell C		42.9 (133)		28.6 (134)		12.5 (115)		14.3 (137)		2.4 (102)
SDM		26.6 (57)		16.0 (69)		24.7 (79)		7.7 (89)		2.6 (104)
Sac City		55.2 (147)		30.6 (141)		38.9 (26)		6.9 (82)		5.7 (130)
Sheldon		29.1 (64)		10.4 (32)		52.0 (8)		9.9 (112)		7.8 (136)
Shenandoah		51.6 (146)		26.6 (127)		16.7 (103)		9.4 (108)		1.7 (88)
Sibley		18.2 (29)		8.3 (22)		25.0 (70)		13.6 (135)		2.7 (109)
Sidney		29.4 (65)		14.3 (56)		33.3 (41)		11.8 (126)		0.0 (1)
Sigourney		38.5 (118)		36.8 (148)		0.0 (125)		11.1 (120)		0.0 (1)
Sioux Cent		9.3 (13)		4.8 (16)		37.5 (29)		0.0 (1)		0.0 (1)
Sioux City		21.8 (43)		11.8 (42)		29.1 (55)		5.4 (62)		1.5 (85)
Spencer		36.7 (109)		19.1 (92)		31.4 (54)		11.4 (121)		13.2 (144)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Ind cators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Spirit Lak		30.8 (72)		18.8 (89)		50.0 (9)		11.5 (123)		19.7 (149)
Storm Lake		10.2 (16)		3.2 (13)		58.3 (5)		5.7 (68)		4.9 (124)
Stuart		42.1 (128)		25.0 (120)		12.5 (115)		10.5 (117)		0.0 (1)
Tama		23.0 (45)		10.6 (35)		15.8 (105)		6.8 (80)		2.6 (104)
Tama/Settl		0.0 (1)		20.0 (97)		0.0 (125)		0.0 (1)		20.0 (150)
Tipton		33.3 (87)		33.3 (145)		0.0 (125)		8.3 (95)		5.6 (128)
Traer		33.3 (87)		10.0 (29)		75.0 (2)		0.0 (1)		10.0 (139)
UIHC		40.0 (121)		17.6 (83)		0.0 (125)		0.0 (1)		7.1 (132)
Vinton		34.5 (101)		11.9 (43)		35.3 (39)		12.3 (128)		0.0 (1)
WDM		21.4 (42)		10.4 (32)		17.6 (102)		0.0 (1)		0.8 (69)
Wapello		100.0 (152)		100.0 (153)		0.0 (125)		100.0 (153)		0.0 (1)
Washington		32.9 (84)		22.7 (108)		11.8 (118)		8.9 (103)		0.0 (1)
Waterloo		33.3 (87)		18.8 (89)		20.3 (92)		10.3 (115)		2.1 (96)
Waukee		10.0 (15)		0.0 (1)		33.3 (41)		1.4 (34)		0.0 (1)
Waukon		25.5 (50)		12.3 (44)		28.6 (56)		7.0 (84)		18.2 (148)
Waverly		28.3 (63)		9.1 (26)		14.3 (108)		2.2 (40)		2.7 (109)
Wbstr Ct		34.8 (102)		15.6 (66)		43.1 (17)		6.3 (72)		0.6 (64)
West Union		37.8 (115)		15.4 (63)		15.4 (106)		22.5 (150)		20.5 (151)
Williamsburg		0.0 (1)		0.0 (1)		0.0 (125)		0.0 (1)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Smoking Indicators by Clinic

Combined 3 Years

Comparison of Smoking Indi ators (2,3)

Clinic	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Winterset		34.2 (99)		12.8 (47)		54.5 (7)		13.2 (132)		2.6 (104)
Iowa		**27.5**		**16.0**		**26.9**		**7.1**		**2.1**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Ackley		0.0 (1)	36.4 (152)		9.1 (61)		0.0 (1)		63.6 (121)
Adel		3.2 (30)	12.9 (132)		0.0 (1)		0.0 (1)		64.5 (119)
Albia		3.9 (35)	9.8 (99)		9.3 (66)		0.0 (1)		62.7 (122)
Algona		7.0 (79)	14.0 (139)		10.0 (78)		2.2 (75)		73.7 (60)
Allen WHC		6.6 (70)	6.2 (46)		7.9 (49)		3.5 (105)		70.2 (82)
Allison		15.2 (146)	9.1 (92)		15.8 (133)		6.3 (136)		78.8 (35)
Altoona		2.0 (25)	5.9 (43)		4.8 (21)		0.0 (1)		73.8 (57)
Ames		9.0 (113)	10.0 (103)		10.7 (91)		3.8 (111)		80.2 (27)
Anamosa		6.5 (67)	7.8 (75)		9.7 (75)		1.5 (61)		68.8 (92)
Ankeny		4.8 (46)	10.3 (109)		4.9 (24)		0.9 (55)		80.8 (25)
Atlantic		6.6 (70)	11.3 (122)		10.9 (94)		3.3 (95)		57.5 (135)
Audubon		8.3 (102)	11.1 (120)		5.9 (28)		3.1 (91)		72.2 (66)
Avoca		5.9 (57)	0.0 (1)		5.9 (28)		6.3 (136)		70.6 (79)
Bedford		4.5 (43)	13.6 (136)		0.0 (1)		0.0 (1)		59.1 (132)
Belle Plaine		0.0 (1)	14.3 (140)		9.5 (71)		0.0 (1)		71.4 (73)
Bloomfield		0.0 (1)	0.0 (1)		6.7 (33)		0.0 (1)		60.0 (128)
Boone		9.3 (120)	8.0 (77)		10.2 (80)		7.5 (145)		64.0 (120)
Brdlwns RP		8.5 (112)	6.6 (51)		9.6 (73)		3.5 (105)		79.6 (29)
Brooklyn		0.0 (1)	6.7 (52)		20.0 (147)		0.0 (1)		68.8 (92)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Burlington		6.7 (73)	9.0 (90)		7.5 (38)		3.9 (114)		55.3 (139)
CHC		11.4 (138)	8.2 (80)		11.4 (103)		6.4 (140)		65.5 (113)
CR Urban		7.2 (85)	8.9 (88)		7.7 (41)		3.5 (105)		71.2 (76)
Carlisle		5.6 (52)	2.8 (18)		16.7 (136)		0.0 (1)		72.2 (66)
Carroll		7.0 (79)	8.7 (86)		9.2 (64)		1.9 (68)		72.2 (66)
Cedar Falls		8.1 (101)	8.1 (78)		11.8 (106)		0.0 (1)		83.8 (14)
Centervill		7.9 (99)	9.2 (95)		13.2 (119)		3.4 (100)		55.3 (139)
Chariton		5.9 (57)	14.7 (141)		16.7 (136)		0.0 (1)		70.6 (79)
Charles City		9.7 (126)	10.3 (109)		12.3 (110)		6.7 (141)		68.8 (92)
Cherokee		11.3 (137)	5.7 (41)		9.2 (64)		9.0 (148)		82.1 (17)
Clarinda		3.6 (32)	5.4 (37)		10.6 (89)		0.0 (1)		72.2 (66)
Clarion		6.6 (70)	9.9 (101)		13.3 (121)		1.1 (57)		72.7 (64)
Clinton		9.6 (125)	7.3 (61)		12.4 (111)		3.3 (95)		69.7 (85)
Colfax		3.4 (31)	13.8 (138)		13.0 (115)		0.0 (1)		72.4 (65)
Columbus J		7.1 (82)	2.4 (17)		10.5 (87)		5.9 (133)		76.2 (45)
Corning		0.0 (1)	12.5 (129)		19.2 (146)		0.0 (1)		59.4 (131)
Corydon		8.3 (102)	20.8 (149)		12.5 (112)		2.9 (85)		66.7 (105)
Council Blfs		9.4 (123)	7.6 (68)		12.5 (112)		3.9 (114)		73.6 (61)
Cresco		9.0 (113)	13.4 (135)		14.3 (125)		0.0 (1)		79.1 (34)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Creston		6.5 (67)	8.8 (87)		8.4 (54)		2.0 (71)		71.9 (70)
Dallas Ctr		0.0 (1)	12.5 (129)		0.0 (1)		0.0 (1)		100.0 (1)
Dayton		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (152)
Decorah		9.3 (120)	12.0 (125)		7.8 (46)		3.4 (100)		85.5 (9)
Denison		8.4 (110)	4.3 (31)		12.6 (114)		1.8 (66)		65.2 (118)
Dewitt		19.2 (149)	0.0 (1)		26.1 (152)		0.0 (1)		65.4 (114)
Dubuque		6.9 (78)	7.6 (68)		7.7 (41)		3.2 (94)		54.5 (142)
Dyersville		0.0 (1)	6.7 (52)		0.0 (1)		0.0 (1)		46.7 (148)
Eagle Grove		7.1 (82)	7.1 (55)		13.4 (122)		1.7 (65)		75.7 (48)
Edgerton		9.1 (117)	8.5 (85)		10.9 (94)		3.3 (95)		67.7 (100)
Eldora		13.6 (142)	4.5 (33)		23.5 (151)		0.0 (1)		54.5 (142)
Elkader		4.3 (39)	10.6 (114)		4.8 (21)		2.5 (78)		83.0 (16)
Emmetsburg		14.5 (145)	8.1 (78)		6.5 (32)		7.0 (143)		66.1 (110)
Esthervill		7.0 (79)	10.9 (118)		8.5 (55)		2.1 (73)		65.4 (114)
Fairfiel		7.6 (96)	12.1 (127)		8.5 (55)		3.7 (108)		68.7 (95)
Forest City		6.5 (67)	13.0 (133)		11.0 (99)		1.8 (66)		65.9 (111)
Fort Dodge		7.4 (91)	7.4 (63)		9.7 (75)		1.4 (60)		67.9 (99)
Ft Madison		9.8 (128)	10.4 (112)		7.6 (40)		4.1 (118)		50.9 (145)
Garner		5.1 (50)	7.6 (68)		6.1 (31)		0.0 (1)		69.6 (86)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Glenwood		6.8 (75)	3.4 (22)		7.3 (37)		2.0 (71)		62.7 (122)
Grand Junction		33.3 (152)	0.0 (1)		0.0 (1)		33.3 (152)		66.7 (105)
Greenfield		0.0 (1)	12.8 (131)		2.5 (14)		0.0 (1)		76.6 (43)
Grinnell		3.8 (33)	6.3 (48)		8.7 (57)		1.6 (63)		67.5 (102)
Grundy Ctr		8.0 (100)	12.0 (125)		19.0 (145)		0.0 (1)		52.0 (144)
Guthrie Ct		6.0 (60)	0.0 (1)		10.4 (83)		2.3 (76)		74.0 (56)
Hampton		3.9 (35)	14.7 (141)		11.8 (106)		0.0 (1)		70.0 (84)
Harlan		5.8 (55)	4.8 (35)		14.6 (126)		1.2 (58)		68.3 (98)
Hawarden		7.4 (91)	11.1 (120)		11.3 (102)		4.3 (121)		81.5 (22)
Hispanic Cln		5.0 (49)	10.0 (103)		0.0 (1)		5.0 (128)		75.0 (50)
Humboldt		4.9 (48)	8.2 (80)		10.9 (94)		0.0 (1)		82.0 (19)
ILH		8.4 (110)	8.3 (83)		10.4 (83)		3.3 (95)		74.3 (55)
Ida Grove		4.3 (39)	4.3 (31)		13.2 (119)		0.0 (1)		78.3 (37)
Independen		7.4 (91)	14.9 (143)		7.7 (41)		2.8 (82)		61.7 (125)
Indianola		12.6 (140)	10.8 (115)		9.3 (66)		6.8 (142)		75.3 (49)
Iowa City		10.3 (130)	7.4 (63)		10.0 (78)		4.9 (127)		85.1 (10)
Iowa Falls		11.1 (135)	2.2 (16)		10.5 (87)		2.9 (85)		77.8 (40)
Jamaica		18.2 (148)	9.1 (92)		9.1 (61)		10.0 (150)		90.9 (5)
Jefferson		5.6 (52)	3.7 (25)		15.1 (131)		0.0 (1)		75.9 (47)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
KR		7.7 (98)	10.3 (109)		17.1 (139)		3.4 (100)		82.1 (17)
Keokuk		9.7 (126)	3.4 (22)		10.7 (91)		6.1 (135)		46.2 (149)
Keosauqua		3.8 (33)	15.4 (144)		4.0 (19)		0.0 (1)		57.7 (133)
Knoxville		6.8 (75)	7.7 (72)		8.3 (51)		3.0 (89)		66.8 (104)
LCPH		10.5 (131)	7.4 (63)		15.2 (132)		1.5 (61)		65.6 (112)
Lamoni		0.0 (1)	25.0 (151)		7.7 (41)		0.0 (1)		81.3 (24)
Le Mars		6.0 (60)	6.0 (44)		11.8 (106)		1.0 (56)		84.3 (13)
Lenox		6.1 (62)	6.1 (45)		3.6 (16)		3.7 (108)		57.6 (134)
Leon		8.3 (102)	6.3 (48)		8.3 (51)		3.0 (89)		60.4 (127)
Logan		13.4 (141)	9.8 (99)		11.4 (103)		7.1 (144)		78.0 (39)
Manchester		9.1 (117)	9.1 (92)		13.5 (123)		6.3 (136)		38.6 (151)
Manning		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		50.0 (146)
Maquoketa		9.0 (113)	5.6 (39)		16.4 (135)		3.3 (95)		57.3 (136)
Marengo		8.3 (102)	20.8 (149)		4.5 (20)		9.5 (149)		70.8 (77)
Marion		6.3 (64)	10.0 (103)		10.6 (89)		2.8 (82)		67.1 (103)
Marshalltown		7.1 (82)	7.8 (75)		9.3 (66)		3.8 (111)		74.5 (54)
Mason City		8.3 (102)	8.2 (80)		12.0 (109)		4.0 (117)		65.4 (114)
Monticello		0.0 (1)	3.8 (26)		7.7 (41)		0.0 (1)		76.9 (42)
Mt Ayr		0.0 (1)	3.2 (21)		3.7 (17)		0.0 (1)		67.7 (100)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Mt Pleasan		10.8 (133)	10.8 (115)		9.4 (70)		4.3 (121)		68.4 (97)
Mtown Dtown		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		66.7 (105)
Muscatine		7.6 (96)	7.2 (59)		11.7 (105)		3.1 (91)		77.6 (41)
Nevada		0.0 (1)	7.7 (72)		0.0 (1)		0.0 (1)		78.6 (36)
New Hampto		7.3 (87)	10.9 (118)		6.0 (30)		4.3 (121)		80.0 (28)
Newton		9.0 (113)	11.9 (123)		8.8 (59)		4.3 (121)		69.1 (88)
North Towne		8.3 (102)	9.7 (98)		9.1 (61)		4.7 (125)		71.6 (72)
Northwood		7.4 (91)	7.4 (63)		9.5 (71)		5.3 (129)		81.5 (22)
Norwalk		14.3 (143)	7.1 (55)		21.9 (150)		0.0 (1)		71.4 (73)
Oakridge		0.0 (1)	0.0 (1)		16.7 (136)		0.0 (1)		46.2 (149)
Oelwein		6.2 (63)	13.6 (136)		9.9 (77)		4.7 (125)		65.4 (114)
Onawa		7.2 (85)	2.9 (19)		7.8 (46)		3.4 (100)		60.9 (126)
Orange Cty		4.1 (38)	10.8 (115)		7.5 (38)		1.6 (63)		86.5 (8)
Osage		2.4 (27)	7.1 (55)		5.8 (27)		0.0 (1)		75.0 (50)
Osceola		8.3 (102)	8.3 (83)		13.9 (124)		1.9 (68)		73.8 (57)
Oskaloosa		7.3 (87)	7.3 (61)		10.4 (83)		1.9 (68)		56.9 (138)
Ottumwa		5.6 (52)	7.2 (59)		7.8 (46)		2.1 (73)		73.4 (63)
Pella		4.4 (42)	15.6 (146)		5.6 (26)		2.9 (85)		68.9 (90)
Perry		6.4 (66)	5.1 (36)		10.4 (83)		3.1 (91)		80.3 (26)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Compar son of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Pocahontas		4.3 (39)	13.0 (133)		15.0 (130)		0.0 (1)		78.3 (37)
Pomeroy		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		100.0 (1)
Postville		8.3 (102)	5.8 (42)		11.1 (100)		4.2 (119)		79.2 (32)
Red Oak		4.6 (44)	10.1 (107)		9.6 (73)		0.0 (1)		70.6 (79)
Redfield		0.0 (1)	0.0 (1)		6.7 (33)		0.0 (1)		60.0 (128)
Refugee Cln		5.9 (57)	17.6 (148)		13.0 (115)		5.9 (133)		96.0 (4)
Rock Rapid		2.0 (25)	10.2 (108)		14.6 (126)		0.0 (1)		73.5 (62)
Rock Valle		5.8 (55)	7.7 (72)		14.7 (128)		3.8 (111)		84.8 (11)
Rockwell C		14.3 (143)	7.1 (55)		10.3 (81)		8.6 (147)		54.8 (141)
SDM		7.5 (95)	8.9 (88)		10.9 (94)		3.4 (100)		75.0 (50)
Sac City		11.8 (139)	2.9 (19)		20.6 (149)		0.0 (1)		61.8 (124)
Sheldon		10.6 (132)	3.8 (26)		14.8 (129)		1.3 (59)		84.6 (12)
Shenandoah		9.3 (120)	4.0 (28)		18.2 (142)		3.7 (108)		70.7 (78)
Sibley		2.6 (28)	10.5 (113)		17.2 (140)		0.0 (1)		81.6 (20)
Sidney		0.0 (1)	9.5 (97)		17.6 (141)		0.0 (1)		76.2 (45)
Sigourney		10.0 (129)	16.7 (147)		7.1 (35)		7.7 (146)		66.7 (105)
Sioux Cent		6.3 (64)	5.6 (39)		8.3 (51)		2.5 (78)		88.3 (7)
Sioux City		7.3 (87)	6.8 (54)		11.2 (101)		2.8 (82)		70.1 (83)
Spencer		4.7 (45)	9.9 (101)		8.7 (57)		0.0 (1)		73.8 (57)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Spirit Lak		2.9 (29)	7.4 (63)		10.3 (81)		0.0 (1)		76.5 (44)
Storm Lake		9.5 (124)	6.2 (46)		13.0 (115)		4.2 (119)		68.5 (96)
Stuart		24.0 (151)	4.0 (28)		18.2 (142)		5.6 (130)		48.0 (147)
Tama		7.3 (87)	12.1 (127)		7.9 (49)		2.9 (85)		66.4 (109)
Tama/Settl		20.0 (150)	0.0 (1)		20.0 (147)		0.0 (1)		60.0 (128)
Tipton		0.0 (1)	0.0 (1)		5.3 (25)		0.0 (1)		57.1 (137)
Traer		16.7 (147)	0.0 (1)		18.2 (142)		11.1 (151)		83.3 (15)
UIHC		86.2 (153)	3.4 (22)		70.0 (153)		33.3 (152)		89.7 (6)
Vinton		1.4 (24)	15.5 (145)		9.0 (60)		0.0 (1)		69.0 (89)
WDM		11.2 (136)	9.0 (90)		13.0 (115)		5.7 (131)		79.3 (31)
Wapello		0.0 (1)	0.0 (1)		0.0 (1)		0.0 (1)		0.0 (152)
Washington		5.1 (50)	4.1 (30)		7.1 (35)		0.0 (1)		81.6 (20)
Waterloo		9.2 (119)	7.6 (68)		10.7 (91)		3.9 (114)		71.4 (73)
Waukee		4.0 (37)	11.9 (123)		4.8 (21)		2.5 (78)		79.4 (30)
Waukon		0.0 (1)	9.4 (96)		3.8 (18)		0.0 (1)		71.9 (70)
Waverly		6.7 (73)	10.0 (103)		9.3 (66)		6.3 (136)		68.9 (90)
Wbstr Ct		10.9 (134)	5.5 (38)		16.3 (134)		5.8 (132)		79.2 (32)
West Union		6.8 (75)	4.5 (33)		2.6 (15)		2.7 (81)		75.0 (50)
Williamsburg		0.0 (1)	100.0 (153)		0.0 (1)		0.0 (1)		100.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 8A

2018 - 2020 Pregnancy Nutrition Surveillance (1)

Iowa

Comparison of Infant Health Indicators by Clinic

Combined 3 Years

Comparison of Infant Health Indicators (2,3)

Clinic	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
		Low (4)	High (5)						
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Winterset		4.8 (46)	6.5 (50)		10.9 (94)		2.4 (77)		69.4 (87)
Iowa		**8.0**	**8.0**		**10.2**		**3.3**		**66.9**

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.