WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Ackley		60.9	0.0	39.1	0.0	0.0	0.0	0.0
Adel		80.8	3.8	7.7	0.0	0.0	7.7	0.0
Albia		93.8	1.8	3.6	0.0	0.0	0.9	0.0
Algona		73.6	0.8	22.0	0.0	0.0	3.7	0.0
Allen WHC		51.3	28.4	6.9	0.0	2.7	10.7	0.0
Allison		87.3	0.8	7.9	0.0	0.0	4.0	0.0
Altoona		65.2	17.4	7.0	0.0	3.0	7.0	0.5
Ames		60.7	11.9	14.7	0.4	4.0	8.3	0.0
Anamosa		87.4	4.9	4.9	0.0	0.7	2.1	0.0
Ankeny		62.9	15.8	12.1	0.3	3.2	5.7	0.0
Atlantic		82.1	0.5	14.4	0.0	0.0	3.1	0.0
Audubon		88.0	1.1	7.6	0.0	2.2	1.1	0.0
Avoca		76.5	2.9	8.8	0.0	0.0	11.8	0.0
Bedford		87.5	0.0	4.2	0.0	2.1	6.3	0.0
Belle Plaine		80.0	1.8	9.1	0.0	0.0	9.1	0.0
Bloomfield		93.8	1.6	3.1	0.0	0.0	1.6	0.0
Boone		83.0	1.4	8.5	0.0	0.9	6.1	0.0
Brdlwns RP		32.0	25.5	23.5	0.1	11.7	7.1	0.0
Brooklyn		86.4	0.0	13.6	0.0	0.0	0.0	0.0
Burlington		64.7	18.5	5.2	0.0	0.6	11.0	0.0
CHC		39.1	24.3	19.1	0.1	1.4	16.0	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution

Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
CR Urban		38.8	38.0	8.5	0.2	1.5	12.9	0.2
Carlisle		80.7	1.1	8.0	0.0	1.1	9.1	0.0
Carroll		77.2	7.7	12.1	0.2	0.2	2.3	0.2
Cedar Falls		73.3	5.9	5.9	0.0	3.0	11.9	0.0
Centervill		90.7	0.5	5.5	0.0	0.5	2.7	0.0
Chariton		87.5	2.0	9.9	0.0	0.0	0.7	0.0
Charles City		72.0	8.0	11.3	0.0	2.3	6.4	0.0
Cherokee		73.4	1.6	20.3	0.5	1.0	2.6	0.5
Clarinda		94.0	2.3	2.3	0.0	0.0	1.5	0.0
Clarion		30.8	0.0	67.8	0.0	0.0	1.4	0.0
Clinton		67.4	7.2	13.3	0.0	0.2	12.0	0.0
Colfax		82.5	1.8	12.3	0.0	1.8	1.8	0.0
Columbus J		18.9	0.0	39.2	0.0	40.5	1.4	0.0
Corning		88.1	1.7	8.5	0.0	0.0	0.0	1.7
Corydon		94.2	0.0	2.9	0.0	2.9	0.0	0.0
Council Blfs		66.9	4.0	20.4	0.1	0.3	8.3	0.0
Cresco		87.4	1.3	9.9	0.0	0.0	1.3	0.0
Creston		84.8	0.9	10.3	0.3	0.0	3.8	0.0
Dallas Ctr		90.0	0.0	10.0	0.0	0.0	0.0	0.0
Dayton		50.0	0.0	50.0	0.0	0.0	0.0	0.0
Decorah		78.4	0.6	17.5	0.0	1.2	2.3	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Denison		24.9	3.6	63.9	0.5	5.9	1.2	0.0
Dewitt		78.9	5.3	10.5	0.0	0.0	5.3	0.0
Dubuque		68.5	14.5	7.1	0.3	0.5	9.1	0.0
Dyersville		80.8	7.7	3.8	3.8	0.0	3.8	0.0
Eagle Grove		29.9	0.0	68.0	0.0	0.0	2.0	0.0
Edgerton		43.3	22.7	15.3	0.7	1.3	16.6	0.1
Eldora		86.6	0.0	9.8	0.0	0.0	3.7	0.0
Elkader		87.9	0.5	8.4	0.0	0.0	3.2	0.0
Emmetsburg		83.5	4.1	10.7	0.0	0.0	1.7	0.0
Esthervill		54.1	2.5	41.8	0.0	0.0	1.6	0.0
Fairfiel		80.2	3.6	10.2	0.0	0.6	5.4	0.0
Forest City		67.9	1.7	20.3	0.0	0.0	10.1	0.0
Fort Dodge		60.7	7.0	18.5	0.3	0.1	12.9	0.4
Ft Madison		86.0	3.8	6.1	0.0	0.3	3.8	0.0
Garner		68.1	0.7	27.4	0.0	0.0	3.7	0.0
Glenwood		89.6	0.0	7.3	0.0	0.0	3.1	0.0
Grand Junction		75.0	0.0	25.0	0.0	0.0	0.0	0.0
Greenfield		88.2	0.0	7.3	0.0	0.9	3.6	0.0
Grinnell		79.5	5.3	9.3	2.0	0.7	3.3	0.0
Grundy Ctr		84.1	2.9	2.9	0.0	0.0	10.1	0.0
Guthrie Ct		86.5	2.7	8.1	0.0	0.0	2.7	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Hampton		44.9	0.0	53.3	0.0	0.0	1.8	0.0
Harlan		73.0	4.0	17.8	0.0	0.0	5.2	0.0
Hawarden		49.2	1.7	45.0	2.5	0.0	1.7	0.0
Hispanic Cln		2.2	2.2	95.7	0.0	0.0	0.0	0.0
Humboldt		69.5	3.5	24.1	0.0	0.0	2.8	0.0
ILH		30.4	16.0	41.2	0.1	5.2	7.1	0.0
Ida Grove		83.0	0.7	15.0	0.0	0.0	1.4	0.0
Independen		86.7	1.4	5.7	0.0	0.0	6.2	0.0
Indianola		87.3	2.0	5.4	0.3	0.0	5.1	0.0
Iowa City		27.2	47.3	18.2	0.2	1.3	5.7	0.0
Iowa Falls		75.6	4.1	15.4	0.0	0.0	4.9	0.0
Jamaica		58.6	3.4	34.5	0.0	0.0	3.4	0.0
Jefferson		82.8	2.0	13.1	1.0	0.0	1.0	0.0
KR		46.7	32.7	6.5	0.0	1.9	12.1	0.0
Keokuk		82.0	4.8	5.1	0.0	0.0	8.1	0.0
Keosauqua		96.9	0.0	1.5	0.0	0.0	1.5	0.0
Knoxville		87.4	3.5	5.0	0.4	0.2	3.3	0.2
LCPH		47.5	27.6	9.5	0.1	0.4	14.7	0.0
Lake City		88.9	0.0	11.1	0.0	0.0	0.0	0.0
Lamoni		83.3	2.4	7.1	0.0	0.0	7.1	0.0
Le Mars		61.7	9.4	23.1	0.6	0.6	4.5	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution

Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Lenox		48.1	0.0	51.9	0.0	0.0	0.0	0.0
Leon		90.7	0.0	7.4	0.0	0.0	1.9	0.0
Logan		88.0	0.8	10.5	0.0	0.0	0.8	0.0
Manchester		84.0	2.5	6.2	0.0	0.0	7.4	0.0
Manning		91.7	0.0	8.3	0.0	0.0	0.0	0.0
Maquoketa		87.8	3.3	3.9	0.0	0.0	5.0	0.0
Marengo		75.0	0.0	18.8	0.0	0.0	6.3	0.0
Marion		71.1	6.6	9.2	0.0	1.4	11.7	0.1
Marshalltown		37.5	3.7	48.2	0.8	7.8	2.0	0.0
Mason City		67.3	6.2	16.3	0.4	0.7	9.2	0.0
Monticello		85.2	5.6	5.6	0.0	0.0	3.7	0.0
Mt Ayr		90.0	0.0	7.1	0.0	0.0	2.9	0.0
Mt Pleasan		80.4	1.8	14.5	0.0	0.4	2.9	0.0
Mtown Dtown		0.0	0.0	75.0	0.0	25.0	0.0	0.0
Muscatine		49.4	9.0	39.2	0.0	1.3	1.1	0.0
Nevada		65.5	0.0	27.3	0.0	0.0	7.3	0.0
New Hampto		72.4	2.4	20.5	0.0	0.0	4.7	0.0
Newton		83.9	4.6	6.5	0.2	1.2	3.6	0.0
Northwood		82.3	4.8	8.1	0.0	0.0	4.8	0.0
Norwalk		82.4	1.4	10.8	0.0	0.0	5.4	0.0
Oakridge		11.0	78.1	5.5	1.4	0.0	4.1	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution								
Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Oelwein		85.8	0.7	7.1	0.0	0.0	6.4	0.0
Onawa		86.7	0.7	7.7	2.8	0.0	2.1	0.0
Orange Cty		48.5	2.4	47.9	0.0	0.0	1.2	0.0
Osage		87.3	0.6	9.9	0.0	0.0	2.2	0.0
Osceola		60.0	0.4	34.4	0.0	1.8	3.5	0.0
Oskaloosa		84.9	4.1	5.1	0.0	1.0	4.8	0.0
Ottumwa		66.2	7.3	17.9	0.1	2.1	6.6	0.0
Pella		89.6	0.0	7.8	0.0	0.0	2.6	0.0
Perry		33.5	10.5	51.1	0.0	2.5	2.5	0.0
Pocahontas		78.7	11.5	8.2	0.0	0.0	1.6	0.0
Postville		31.8	11.7	56.5	0.0	0.0	0.0	0.0
Red Oak		83.9	0.0	12.8	0.0	0.0	3.2	0.0
Redfield		90.0	0.0	0.0	0.0	0.0	10.0	0.0
Refugee Cln		1.2	98.8	0.0	0.0	0.0	0.0	0.0
Rock Rapid		80.8	0.0	12.0	0.8	0.0	6.4	0.0
Rock Valle		39.3	0.0	59.7	0.0	0.0	1.0	0.0
Rockwell C		91.0	0.0	7.0	0.0	0.0	2.0	0.0
SDM		35.8	17.3	32.4	0.2	8.6	5.8	0.0
Sac City		87.3	1.6	11.1	0.0	0.0	0.0	0.0
Sheldon		63.6	2.7	26.8	0.0	0.0	6.8	0.0
Shenandoah		86.0	0.0	11.5	0.0	0.0	2.5	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution

Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Sibley		47.9	0.0	52.1	0.0	0.0	0.0	0.0
Sidney		96.5	0.0	3.5	0.0	0.0	0.0	0.0
Sigourney		90.7	0.0	2.3	0.0	0.0	7.0	0.0
Sioux Cent		25.6	0.6	70.4	0.0	1.2	2.2	0.0
Sioux City		30.6	14.2	40.0	3.6	1.5	9.9	0.2
Spencer		78.8	2.3	16.4	0.0	0.8	1.7	0.0
Spirit Lak		81.4	0.0	12.4	0.0	0.6	5.6	0.0
Storm Lake		13.9	5.2	58.5	0.0	12.2	10.2	0.0
Stuart		75.0	13.5	9.6	1.9	0.0	0.0	0.0
Tama		40.2	0.5	44.7	8.7	0.0	5.9	0.0
Tama/Settl		9.5	0.0	14.3	61.9	0.0	14.3	0.0
Tipton		82.6	4.3	6.5	0.0	0.0	6.5	0.0
Traer		92.0	0.0	8.0	0.0	0.0	0.0	0.0
UIHC		66.7	9.1	18.2	0.0	6.1	0.0	0.0
Vinton		90.1	0.6	7.5	0.0	0.0	1.9	0.0
WDM		43.4	24.2	22.0	0.7	3.9	5.9	0.0
Washington		65.0	2.0	31.5	0.0	0.0	1.5	0.0
Waterloo		44.1	25.4	11.2	0.1	4.2	14.9	0.1
Waukee		64.6	10.3	15.6	0.0	2.7	6.8	0.0
Waukon		87.0	0.0	10.9	0.0	0.0	2.2	0.0
Waverly		86.9	3.3	1.3	0.0	1.3	7.2	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Racial and Ethnic Distribution

Clinic	Number Included in Analysis (3,4)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wbstr Ct		52.2	2.0	41.0	0.0	2.7	2.0	0.0
West Union		92.8	1.2	4.8	0.0	0.0	1.2	0.0
Williamsburg		33.3	0.0	55.6	0.0	0.0	11.1	0.0
Winterset		89.1	0.0	7.7	0.0	1.3	1.9	0.0
Iowa		**52.7**	**14.6**	**21.5**	**0.4**	**3.1**	**7.7**	**0.0**
National (Prior Year)	**712,311**	**37.6**	**23.5**	**23.2**	**1.4**	**2.6**	**6.5**	**5.1**

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Ackley		39.1	13.0	47.8
Adel		34.6	20.5	44.9
Albia		40.2	23.2	36.6
Algona		43.1	15.0	41.9
Allen WHC		48.7	17.3	34.0
Allison		37.3	15.9	46.8
Altoona		33.8	21.9	44.3
Ames		34.3	22.6	43.1
Anamosa		39.2	22.4	38.5
Ankeny		36.8	22.7	40.5
Atlantic		42.1	20.0	37.9
Audubon		25.0	22.8	52.2
Avoca		35.3	14.7	50.0
Bedford		41.7	12.5	45.8
Belle Plaine		36.4	14.5	49.1
Bloomfield		40.6	20.3	39.1
Boone		34.9	19.3	45.8
Brdlwns RP		37.9	20.8	41.3
Brooklyn		22.7	22.7	54.5
Burlington		48.7	18.2	33.0
CHC		41.9	18.9	39.1

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
CR Urban		42.1	16.3	41.6
Carlisle		42.0	17.0	40.9
Carroll		43.5	14.0	42.5
Cedar Falls		33.7	16.8	49.5
Centervill		39.0	20.9	40.1
Chariton		39.5	17.8	42.8
Charles City		37.9	17.4	44.7
Cherokee		33.9	20.8	45.3
Clarinda		41.4	15.0	43.6
Clarion		41.7	14.7	43.6
Clinton		46.6	17.1	36.3
Colfax		36.8	15.8	47.4
Columbus J		43.2	9.5	47.3
Corning		42.4	18.6	39.0
Corydon		42.7	14.6	42.7
Council Blfs		38.9	19.2	42.0
Cresco		37.1	23.2	39.7
Creston		40.5	17.6	41.9
Dallas Ctr		20.0	35.0	45.0
Dayton		50.0	50.0	0.0
Decorah		39.2	15.8	45.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Denison		40.9	15.8	43.3
Dewitt		54.4	14.0	31.6
Dubuque		45.7	15.9	38.4
Dyersville		46.2	11.5	42.3
Eagle Grove		34.0	19.7	46.3
Edgerton		46.1	15.1	38.7
Eldora		28.0	28.0	43.9
Elkader		49.5	15.8	34.7
Emmetsburg		43.0	14.0	43.0
Esthervill		38.1	18.0	43.9
Fairfiel		40.1	16.2	43.7
Forest City		43.5	12.2	44.3
Fort Dodge		45.9	18.5	35.6
Ft Madison		41.4	14.0	44.6
Garner		35.6	20.0	44.4
Glenwood		43.8	13.5	42.7
Grand Junction		18.8	31.3	50.0
Greenfield		40.9	20.0	39.1
Grinnell		31.8	23.2	45.0
Grundy Ctr		33.3	18.8	47.8
Guthrie Ct		33.3	11.7	55.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Hampton		40.8	15.1	44.1
Harlan		39.7	18.4	42.0
Hawarden		31.7	22.5	45.8
Hispanic Cln		30.4	15.2	54.3
Humboldt		37.6	14.9	47.5
ILH		35.3	20.9	43.9
Ida Grove		35.4	16.3	48.3
Independen		33.2	24.6	42.2
Indianola		38.6	16.3	45.1
Iowa City		39.5	18.8	41.8
Iowa Falls		35.0	22.8	42.3
Jamaica		24.1	6.9	69.0
Jefferson		35.4	15.2	49.5
KR		29.0	19.6	51.4
Keokuk		46.7	16.2	37.1
Keosauqua		43.1	18.5	38.5
Knoxville		38.7	18.6	42.7
LCPH		46.5	16.8	36.7
Lake City		66.7	11.1	22.2
Lamoni		35.7	14.3	50.0
Le Mars		38.0	17.9	44.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Lenox		38.5	11.5	50.0
Leon		45.4	14.8	39.8
Logan		37.6	21.1	41.4
Manchester		32.1	25.9	42.0
Manning		25.0	8.3	66.7
Maquoketa		38.7	16.0	45.3
Marengo		35.9	18.8	45.3
Marion		38.8	18.0	43.2
Marshalltown		38.9	19.5	41.6
Mason City		40.7	19.3	40.0
Monticello		40.7	20.4	38.9
Mt Ayr		42.9	15.7	41.4
Mt Pleasan		38.5	17.1	44.4
Mtown Dtown		0.0	75.0	25.0
Muscatine		42.3	18.5	39.2
Nevada		30.9	25.5	43.6
New Hampto		27.6	25.2	47.2
Newton		40.5	18.0	41.5
Northwood		32.3	17.7	50.0
Norwalk		29.7	25.7	44.6
Oakridge		17.8	19.2	63.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Oelwein		42.6	14.9	42.6
Onawa		40.6	18.2	41.3
Orange Cty		33.1	23.1	43.8
Osage		39.2	17.1	43.6
Osceola		41.1	19.3	39.6
Oskaloosa		43.2	15.4	41.4
Ottumwa		45.1	18.2	36.6
Pella		38.3	14.8	47.0
Perry		34.2	19.4	46.5
Pocahontas		31.1	18.0	50.8
Postville		37.7	16.1	46.2
Red Oak		36.7	16.5	46.8
Redfield		40.0	33.3	26.7
Refugee Cln		33.7	16.3	50.0
Rock Rapid		41.6	16.8	41.6
Rock Valle		40.3	17.3	42.4
Rockwell C		28.0	21.0	51.0
SDM		35.7	22.3	42.1
Sac City		30.2	19.0	50.8
Sheldon		39.1	17.3	43.6
Shenandoah		40.1	15.9	43.9

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Sibley		38.4	13.7	47.9
Sidney		38.6	8.8	52.6
Sigourney		36.0	12.8	51.2
Sioux Cent		36.1	19.4	44.4
Sioux City		43.0	17.1	39.9
Spencer		40.7	16.9	42.4
Spirit Lak		46.6	19.3	34.2
Storm Lake		39.0	17.8	43.2
Stuart		36.5	23.1	40.4
Tama		37.9	16.9	45.2
Tama/Settl		33.3	14.3	52.4
Tipton		34.8	10.9	54.3
Traer		20.0	4.0	76.0
UIHC		54.5	24.2	21.2
Vinton		32.9	19.3	47.8
WDM		31.6	20.7	47.7
Washington		37.1	20.8	42.1
Waterloo		41.7	18.0	40.3
Waukee		28.5	27.0	44.5
Waukon		42.8	21.0	36.2
Waverly		28.8	22.9	48.4

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Comparison of Age Distribution				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Wbstr Ct		43.3	15.7	41.0
West Union		43.4	24.1	32.5
Williamsburg		11.1	11.1	77.8
Winterset		43.6	17.9	38.5
Iowa		**40.1**	**18.5**	**41.4**
National (Prior Year)	**712,311**	**35.1**	**21.8**	**43.1**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ackley		14.3 (118)	28.6 (148)		4.5 (14)	0.0 (1)	13.6 (51)		16.7 (75)	16.7 (66)		14.3 (119)
Adel		5.9 (46)	17.6 (142)		6.9 (44)	1.4 (12)	16.7 (94)		21.6 (128)	21.6 (115)		0.0 (1)
Albia		3.6 (29)	14.3 (133)		13.7 (114)	6.9 (130)	16.7 (94)		20.5 (125)	20.5 (104)		0.0 (1)
Algona		7.5 (55)	7.5 (84)		6.6 (37)	1.8 (18)	15.9 (84)		27.7 (147)	16.8 (67)		11.8 (101)
Allen WHC		11.7 (102)	4.3 (47)		13.1 (108)	5.3 (117)	19.4 (126)		12.5 (26)	20.5 (104)		14.8 (125)
Allison		10.5 (91)	15.8 (141)		12.4 (97)	4.1 (72)	9.1 (8)		10.2 (15)	15.3 (59)		21.1 (139)
Altoona		11.1 (96)	3.7 (40)		8.9 (66)	1.6 (15)	12.1 (36)		27.1 (146)	12.9 (33)		7.1 (59)
Ames		9.0 (74)	6.7 (67)		13.3 (110)	4.7 (94)	15.7 (77)		15.2 (55)	15.2 (57)		7.5 (66)
Anamosa		13.3 (113)	3.3 (37)		13.0 (107)	2.3 (29)	14.5 (59)		17.5 (89)	14.0 (48)		12.5 (106)
Ankeny		15.5 (127)	9.5 (108)		9.2 (70)	2.8 (37)	9.5 (11)		15.6 (64)	13.3 (41)		11.3 (100)
Atlantic		8.1 (60)	0.0 (1)		14.4 (125)	4.8 (98)	16.0 (85)		18.7 (108)	14.7 (52)		29.2 (147)
Audubon		7.7 (58)	30.8 (150)		3.4 (12)	1.1 (11)	10.3 (15)		22.0 (129)	10.0 (16)		4.5 (40)
Avoca		0.0 (1)	0.0 (1)		11.8 (96)	8.8 (145)	5.9 (4)		16.7 (75)	5.6 (4)		10.0 (87)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Bedford		11.1 (96)	0.0 (1)		11.4 (88)	2.3 (29)	18.2 (109)		14.3 (40)	23.8 (127)		11.1 (96)
Belle Plaine		0.0 (1)	0.0 (1)		5.7 (19)	3.8 (63)	13.2 (47)		14.3 (40)	21.4 (113)		10.0 (87)
Bloomfield		7.1 (52)	14.3 (133)		10.0 (77)	5.0 (106)	20.0 (130)		14.3 (40)	23.8 (127)		6.7 (54)
Boone		20.8 (141)	4.2 (44)		12.4 (97)	7.4 (137)	11.4 (28)		9.4 (11)	10.4 (18)		0.0 (1)
Brdlwns RP		10.6 (94)	6.7 (67)		8.1 (58)	2.6 (35)	11.5 (29)		15.2 (55)	13.0 (35)		9.0 (79)
Brooklyn		0.0 (1)	33.3 (151)		13.6 (112)	4.5 (86)	13.6 (51)		23.1 (134)	7.7 (7)		0.0 (1)
Burlington		9.3 (80)	8.4 (97)		9.0 (68)	5.0 (106)	14.7 (62)		15.4 (60)	18.2 (82)		9.7 (85)
CHC		14.3 (118)	5.9 (58)		12.4 (97)	3.6 (58)	15.7 (77)		19.5 (119)	17.5 (76)		20.7 (137)
CR Urban		9.8 (87)	7.8 (91)		8.3 (61)	2.3 (29)	10.6 (21)		15.5 (63)	11.0 (21)		19.4 (135)
Carlisle		16.7 (130)	8.3 (93)		16.9 (136)	7.2 (136)	21.7 (137)		11.1 (20)	27.8 (141)		11.1 (96)
Carroll		9.4 (81)	10.4 (114)		7.1 (48)	2.9 (39)	10.3 (15)		12.9 (30)	8.9 (10)		3.7 (32)
Cedar Falls		9.1 (75)	18.2 (143)		15.1 (126)	3.2 (47)	18.3 (111)		14.3 (40)	12.2 (29)		13.8 (117)
Centervill		4.9 (37)	14.6 (136)		5.8 (22)	4.7 (94)	15.2 (68)		14.3 (40)	16.9 (69)		5.7 (47)
Chariton		3.1 (26)	9.4 (106)		15.2 (127)	5.8 (121)	15.2 (68)		17.2 (85)	17.2 (72)		8.7 (77)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Charles City		13.6 (114)	18.6 (144)		7.8 (55)	3.4 (56)	13.7 (53)		15.6 (64)	12.8 (31)		13.6 (115)
Cherokee		4.5 (35)	9.1 (102)		10.3 (79)	4.9 (99)	17.4 (102)		13.3 (33)	23.3 (124)		6.7 (54)
Clarinda		10.5 (91)	0.0 (1)		17.6 (140)	6.4 (129)	20.0 (130)		19.0 (112)	17.2 (72)		7.7 (68)
Clarion		10.4 (90)	4.2 (44)		12.4 (97)	3.0 (43)	18.3 (111)		20.4 (124)	17.2 (72)		1.7 (25)
Clinton		6.6 (51)	6.6 (66)		9.7 (76)	2.9 (39)	12.7 (41)		15.0 (51)	13.8 (47)		11.2 (99)
Colfax		15.4 (124)	7.7 (87)		6.3 (26)	2.1 (24)	10.4 (19)		9.1 (9)	13.6 (44)		0.0 (1)
Columbus J		18.8 (137)	6.3 (62)		13.8 (116)	4.6 (90)	23.1 (142)		16.7 (75)	20.0 (98)		8.3 (71)
Corning		14.3 (118)	7.1 (73)		9.6 (75)	5.8 (121)	13.5 (50)		22.7 (133)	22.7 (122)		10.0 (87)
Corydon		4.0 (30)	12.0 (123)		11.7 (93)	8.5 (144)	17.0 (98)		28.2 (148)	25.6 (134)		0.0 (1)
Council Blfs		11.4 (101)	7.1 (73)		10.5 (81)	4.6 (90)	16.5 (91)		20.1 (123)	20.0 (98)		19.0 (134)
Cresco		7.4 (54)	14.8 (137)		10.6 (82)	4.9 (99)	18.3 (111)		15.2 (55)	28.8 (145)		7.1 (59)
Creston		8.2 (62)	11.0 (118)		15.2 (127)	3.9 (67)	24.8 (145)		19.3 (115)	23.7 (126)		13.2 (111)
Dallas Ctr		100.0 (151)	0.0 (1)		31.6 (150)	10.5 (148)	15.8 (81)		10.0 (13)	20.0 (98)		0.0 (1)
Dayton		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Decorah		0.0 (1)	14.3 (133)		8.2 (59)	3.8 (63)	15.8 (81)		19.2 (113)	13.7 (45)		2.1 (27)
Denison		10.8 (95)	5.8 (57)		15.3 (131)	5.0 (106)	17.8 (103)		16.4 (73)	20.0 (98)		26.4 (144)
Dewitt		10.5 (91)	10.5 (116)		5.3 (18)	0.0 (1)	12.3 (38)		26.3 (144)	21.1 (109)		14.3 (119)
Dubuque		8.3 (65)	8.3 (93)		7.0 (46)	4.3 (80)	10.4 (19)		18.3 (103)	10.0 (16)		12.1 (104)
Dyersville		42.9 (147)	0.0 (1)		15.4 (132)	0.0 (1)	19.2 (124)		9.1 (9)	18.2 (82)		0.0 (1)
Eagle Grove		3.4 (28)	6.9 (70)		9.4 (71)	2.9 (39)	21.7 (137)		23.8 (139)	27.0 (139)		10.4 (94)
Edgerton		12.6 (111)	9.9 (112)		15.2 (127)	3.1 (46)	18.7 (118)		15.6 (64)	20.7 (106)		15.3 (127)
Eldora		11.8 (103)	0.0 (1)		7.4 (51)	4.9 (99)	16.0 (85)		22.0 (129)	22.0 (118)		27.8 (146)
Elkader		12.3 (106)	12.3 (127)		12.6 (104)	6.9 (130)	14.9 (65)		26.5 (145)	13.2 (39)		9.5 (84)
Emmetsburg		6.5 (50)	9.7 (109)		6.9 (44)	6.0 (126)	12.9 (45)		25.5 (141)	9.1 (11)		13.5 (113)
Esthervill		6.0 (47)	8.0 (92)		11.5 (90)	4.9 (99)	17.3 (100)		22.6 (131)	18.9 (90)		13.7 (116)
Fairfiel		4.9 (37)	7.3 (80)		6.3 (26)	7.5 (140)	15.0 (66)		26.0 (143)	13.7 (45)		5.9 (48)
Forest City		5.7 (45)	9.4 (106)		3.2 (11)	5.9 (123)	12.7 (41)		14.7 (49)	16.8 (67)		7.3 (63)
Fort Dodge		6.0 (47)	9.0 (101)		17.6 (140)	2.5 (34)	23.9 (144)		16.4 (73)	28.4 (143)		1.9 (26)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ft Madison		5.6 (44)	9.7 (109)		11.2 (87)	2.2 (26)	17.3 (100)		15.0 (51)	22.0 (118)		22.2 (141)
Garner		14.8 (123)	7.4 (82)		3.1 (10)	2.3 (29)	9.2 (9)		23.5 (135)	13.2 (39)		8.3 (71)
Glenwood		21.7 (142)	4.3 (47)		23.3 (147)	8.9 (146)	22.2 (139)		13.9 (38)	27.8 (141)		10.0 (87)
Grand Junction		0.0 (1)	0.0 (1)		25.0 (148)	0.0 (1)	6.3 (5)		33.3 (151)	11.1 (23)		0.0 (1)
Greenfield		4.3 (33)	13.0 (130)		11.7 (93)	4.9 (99)	12.6 (40)		12.8 (28)	15.4 (61)		12.5 (106)
Grinnell		12.5 (107)	20.8 (145)		14.2 (122)	3.5 (57)	15.6 (75)		21.0 (126)	8.1 (8)		10.0 (87)
Grundy Ctr		9.1 (75)	9.1 (102)		15.2 (127)	3.0 (43)	12.1 (36)		9.4 (11)	15.6 (63)		5.9 (48)
Guthrie Ct		0.0 (1)	10.5 (116)		2.9 (8)	2.9 (39)	14.6 (61)		15.3 (59)	18.6 (85)		0.0 (1)
Hampton		5.2 (41)	6.9 (70)		7.1 (48)	4.3 (80)	15.8 (81)		12.8 (28)	17.1 (70)		3.9 (34)
Harlan		12.2 (105)	4.9 (51)		14.1 (121)	4.3 (80)	17.2 (99)		15.2 (55)	21.5 (114)		18.2 (131)
Hawarden		28.0 (145)	12.0 (123)		6.3 (26)	6.3 (127)	11.7 (31)		23.6 (137)	14.5 (51)		4.0 (35)
Hispanic Cln		12.5 (107)	12.5 (128)		11.6 (92)	0.0 (1)	11.6 (30)		16.0 (71)	20.0 (98)		22.2 (141)
Humboldt		3.2 (27)	6.5 (65)		13.7 (114)	3.8 (63)	20.6 (133)		18.0 (97)	29.5 (146)		4.1 (37)
ILH		9.2 (79)	5.9 (58)		6.7 (38)	1.8 (18)	12.4 (39)		23.6 (137)	14.2 (49)		16.0 (129)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ida Grove		18.5 (136)	0.0 (1)		6.4 (30)	4.3 (80)	18.6 (117)		19.4 (118)	15.3 (59)		3.0 (30)
Independen		15.8 (128)	5.3 (53)		11.0 (85)	4.7 (94)	17.8 (103)		19.3 (115)	14.8 (53)		13.0 (110)
Indianola		7.6 (56)	12.1 (125)		16.0 (134)	6.9 (130)	18.4 (114)		18.0 (97)	19.3 (95)		7.2 (62)
Iowa City		15.4 (124)	7.1 (73)		17.1 (138)	8.1 (143)	18.9 (121)		13.4 (34)	19.3 (95)		13.5 (113)
Iowa Falls		0.0 (1)	3.4 (38)		13.9 (118)	9.6 (147)	11.3 (26)		15.1 (54)	9.4 (13)		31.6 (148)
Jamaica		40.0 (146)	0.0 (1)		0.0 (1)	3.7 (60)	14.8 (63)		20.0 (122)	15.0 (55)		0.0 (1)
Jefferson		0.0 (1)	3.8 (41)		9.5 (73)	7.4 (137)	16.8 (97)		11.5 (23)	19.2 (94)		5.9 (48)
KR		17.6 (132)	23.5 (146)		6.5 (36)	3.3 (52)	16.3 (89)		19.6 (120)	10.9 (20)		21.1 (139)
Keokuk		8.8 (72)	4.4 (50)		18.4 (142)	4.5 (86)	18.8 (120)		14.1 (39)	21.2 (111)		14.1 (118)
Keosauqua		11.1 (96)	11.1 (119)		7.7 (53)	4.6 (90)	15.4 (71)		10.3 (16)	20.7 (106)		10.0 (87)
Knoxville		9.5 (82)	11.6 (120)		13.2 (109)	4.0 (69)	20.3 (132)		17.2 (85)	19.1 (93)		9.2 (81)
LCPH		11.1 (96)	6.9 (70)		10.1 (78)	3.7 (60)	13.9 (55)		13.0 (31)	15.6 (63)		18.4 (133)
Lake City		20.0 (139)	0.0 (1)		11.1 (86)	33.3 (151)	11.1 (24)		0.0 (1)	50.0 (151)		0.0 (1)
Lamoni		0.0 (1)	0.0 (1)		14.3 (123)	14.3 (150)	28.6 (150)		10.5 (17)	26.3 (135)		20.0 (136)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Le Mars		8.6 (71)	7.1 (73)		8.8 (65)	4.4 (84)	17.8 (103)		13.5 (35)	14.2 (49)		6.9 (57)
Lenox		12.5 (107)	0.0 (1)		20.8 (144)	2.1 (24)	22.9 (141)		15.4 (60)	11.5 (27)		0.0 (1)
Leon		13.0 (112)	4.3 (47)		8.2 (59)	4.1 (72)	11.3 (26)		28.6 (149)	9.5 (14)		26.7 (145)
Logan		25.9 (144)	7.4 (82)		21.5 (146)	4.1 (72)	15.7 (77)		23.5 (135)	17.6 (78)		22.7 (143)
Manchester		0.0 (1)	15.4 (139)		11.5 (90)	5.1 (113)	17.9 (106)		16.7 (75)	11.9 (28)		5.3 (44)
Manning		50.0 (148)	0.0 (1)		0.0 (1)	0.0 (1)	10.0 (12)		0.0 (1)	11.1 (23)		0.0 (1)
Maquoketa		8.3 (65)	5.6 (55)		7.2 (50)	2.2 (26)	11.7 (31)		18.6 (106)	12.8 (31)		4.5 (40)
Marengo		25.0 (143)	0.0 (1)		1.7 (6)	5.2 (114)	10.3 (15)		13.8 (37)	6.9 (5)		18.2 (131)
Marion		8.1 (60)	3.8 (41)		6.4 (30)	1.4 (12)	10.7 (22)		19.2 (113)	13.1 (38)		8.2 (70)
Marshalltown		9.1 (75)	7.1 (73)		7.0 (46)	4.2 (77)	13.2 (47)		21.1 (127)	15.0 (55)		15.9 (128)
Mason City		8.9 (73)	7.7 (87)		6.8 (42)	3.3 (52)	14.8 (63)		17.1 (84)	18.6 (85)		12.9 (109)
Monticello		0.0 (1)	0.0 (1)		1.9 (7)	7.4 (137)	5.6 (3)		13.0 (31)	13.0 (35)		15.0 (126)
Mt Ayr		11.1 (96)	0.0 (1)		12.7 (105)	6.3 (127)	25.4 (148)		17.9 (94)	25.0 (132)		50.0 (151)
Mt Pleasan		6.3 (49)	10.4 (114)		10.7 (83)	5.2 (114)	16.7 (94)		16.7 (75)	21.1 (109)		10.8 (95)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Mtown Dtown		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)	25.0 (146)		0.0 (1)	33.3 (149)		0.0 (1)
Muscatine		13.9 (116)	8.6 (99)		9.1 (69)	2.6 (35)	13.9 (55)		18.5 (104)	17.7 (79)		8.5 (74)
Nevada		9.1 (75)	0.0 (1)		5.8 (22)	1.9 (21)	19.2 (124)		12.5 (26)	20.8 (108)		8.3 (71)
New Hampto		0.0 (1)	15.0 (138)		12.4 (97)	5.0 (106)	16.5 (91)		15.9 (69)	23.8 (127)		14.3 (119)
Newton		18.0 (134)	6.7 (67)		8.9 (66)	5.9 (123)	19.6 (128)		15.4 (60)	24.3 (131)		3.8 (33)
Northwood		0.0 (1)	0.0 (1)		3.4 (12)	5.2 (114)	10.3 (15)		17.2 (85)	6.9 (5)		7.1 (59)
Norwalk		20.0 (139)	10.0 (113)		16.9 (136)	2.8 (37)	21.1 (135)		14.7 (49)	26.5 (136)		17.6 (130)
Oakridge		0.0 (1)	0.0 (1)		3.0 (9)	1.5 (14)	4.5 (2)		5.0 (7)	5.0 (2)		20.8 (138)
Oelwein		12.5 (107)	4.2 (44)		6.3 (26)	4.7 (94)	21.1 (135)		14.5 (47)	25.5 (133)		0.0 (1)
Onawa		18.2 (135)	6.1 (60)		17.3 (139)	5.0 (106)	19.4 (126)		14.5 (47)	27.4 (140)		12.5 (106)
Orange Cty		9.7 (85)	3.2 (36)		5.7 (19)	3.2 (47)	8.9 (7)		17.6 (91)	5.4 (3)		3.0 (30)
Osage		4.9 (37)	12.2 (126)		5.9 (24)	2.4 (33)	10.0 (12)		11.0 (19)	11.0 (21)		9.3 (83)
Osceola		10.2 (88)	5.1 (52)		14.3 (123)	5.6 (119)	18.7 (118)		17.5 (89)	16.5 (65)		8.5 (74)
Oskaloosa		4.0 (30)	4.0 (43)		9.4 (71)	4.0 (69)	13.0 (46)		15.7 (67)	11.3 (26)		4.9 (43)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ottumwa		8.5 (69)	7.5 (84)		10.3 (79)	5.4 (118)	16.6 (93)		15.9 (69)	18.3 (84)		7.0 (58)
Pella		4.3 (33)	13.0 (130)		6.4 (30)	3.7 (60)	15.6 (75)		22.6 (131)	18.9 (90)		7.4 (65)
Perry		17.9 (133)	3.0 (35)		6.7 (38)	1.9 (21)	11.9 (34)		24.8 (140)	14.8 (53)		2.3 (28)
Pocahontas		0.0 (1)	0.0 (1)		12.5 (103)	7.1 (134)	16.1 (88)		18.2 (101)	15.2 (57)		4.0 (35)
Postville		9.5 (82)	11.9 (121)		18.6 (143)	3.8 (63)	22.4 (140)		18.1 (99)	26.7 (138)		4.8 (42)
Red Oak		13.6 (114)	9.1 (102)		12.4 (97)	5.7 (120)	15.2 (68)		13.5 (35)	10.6 (19)		11.8 (101)
Redfield		0.0 (1)	0.0 (1)		6.7 (38)	3.3 (52)	10.0 (12)		0.0 (1)	11.1 (23)		36.4 (149)
Refugee Cln		0.0 (1)	7.7 (87)		8.3 (61)	0.0 (1)	11.9 (34)		25.6 (142)	9.3 (12)		40.0 (150)
Rock Rapid		9.7 (85)	9.7 (109)		10.7 (83)	4.1 (72)	14.0 (57)		10.7 (18)	17.9 (81)		0.0 (1)
Rock Valle		7.7 (58)	7.7 (87)		8.3 (61)	3.9 (67)	13.3 (49)		12.2 (25)	17.1 (70)		2.5 (29)
Rockwell C		0.0 (1)	6.3 (62)		7.6 (52)	2.2 (26)	18.5 (116)		16.0 (71)	24.0 (130)		0.0 (1)
SDM		8.2 (62)	7.2 (79)		6.4 (30)	3.2 (47)	11.7 (31)		17.7 (92)	12.6 (30)		5.5 (45)
Sac City		57.1 (149)	0.0 (1)		15.9 (133)	1.6 (15)	6.3 (5)		11.8 (24)	8.8 (9)		0.0 (1)
Sheldon		17.1 (131)	7.3 (80)		11.4 (88)	5.0 (106)	18.9 (121)		14.3 (40)	23.1 (123)		8.7 (77)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Shenandoah		10.3 (89)	3.4 (38)		11.7 (93)	6.9 (130)	11.0 (23)		11.3 (22)	12.9 (33)		13.3 (112)
Sibley		14.3 (118)	0.0 (1)		7.9 (56)	3.2 (47)	12.7 (41)		3.3 (6)	13.3 (41)		6.7 (54)
Sidney		7.1 (52)	0.0 (1)		28.3 (149)	7.5 (140)	20.8 (134)		18.5 (104)	29.6 (147)		14.3 (119)
Sigourney		14.3 (118)	23.8 (147)		4.9 (15)	4.9 (99)	11.1 (24)		19.6 (120)	13.0 (35)		0.0 (1)
Sioux Cent		2.9 (25)	8.8 (100)		6.4 (30)	3.2 (47)	16.0 (85)		15.0 (51)	23.6 (125)		7.3 (63)
Sioux City		7.6 (56)	7.6 (86)		9.5 (73)	1.8 (18)	19.8 (129)		18.1 (99)	21.8 (117)		11.9 (103)
Spencer		5.5 (43)	2.7 (33)		8.5 (64)	4.9 (99)	13.7 (53)		18.7 (108)	18.7 (87)		10.2 (93)
Spirit Lak		9.5 (82)	11.9 (121)		5.9 (24)	4.6 (90)	15.7 (77)		19.3 (115)	17.5 (76)		4.3 (39)
Storm Lake		5.0 (40)	2.8 (34)		6.8 (42)	4.2 (77)	14.5 (59)		17.7 (92)	18.8 (89)		8.6 (76)
Stuart		0.0 (1)	28.6 (148)		13.6 (112)	4.5 (86)	29.5 (151)		11.1 (20)	22.2 (120)		9.1 (80)
Tama		15.4 (124)	15.4 (139)		6.7 (38)	3.6 (58)	15.5 (74)		16.9 (82)	21.7 (116)		6.3 (52)
Tama/Settl		0.0 (1)	0.0 (1)		5.0 (17)	5.0 (106)	15.0 (66)		10.0 (13)	20.0 (98)		0.0 (1)
Tipton		8.3 (65)	8.3 (93)		21.4 (145)	7.1 (134)	23.8 (143)		18.2 (101)	31.8 (148)		7.7 (68)
Traer		0.0 (1)	0.0 (1)		8.0 (57)	4.0 (69)	28.0 (149)		16.7 (75)	22.2 (120)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
UIHC		66.7 (150)	0.0 (1)		34.6 (151)	7.7 (142)	15.4 (71)		14.3 (40)	28.6 (144)		14.3 (119)
Vinton		4.2 (32)	12.5 (128)		12.8 (106)	2.0 (23)	18.9 (121)		17.3 (88)	18.7 (87)		11.1 (96)
WDM		8.2 (62)	7.1 (73)		7.7 (53)	3.0 (43)	12.8 (44)		18.8 (110)	13.5 (43)		9.2 (81)
Washington		8.3 (65)	8.3 (93)		14.0 (119)	4.5 (86)	18.0 (108)		17.9 (94)	15.5 (62)		14.6 (124)
Waterloo		11.8 (103)	6.1 (60)		13.4 (111)	5.9 (123)	16.3 (89)		15.8 (68)	19.0 (92)		12.4 (105)
Waukee		8.5 (69)	8.5 (98)		4.9 (15)	1.6 (15)	9.3 (10)		18.6 (106)	9.7 (15)		9.8 (86)
Waukon		16.2 (129)	13.5 (132)		5.7 (19)	3.3 (52)	15.4 (71)		17.0 (83)	21.3 (112)		6.1 (51)
Waverly		18.8 (137)	6.3 (62)		16.8 (135)	4.2 (77)	18.2 (109)		5.5 (8)	17.8 (80)		7.5 (66)
Wbstr Ct		4.5 (35)	9.1 (102)		14.0 (119)	4.4 (84)	18.4 (114)		17.9 (94)	19.7 (97)		6.3 (52)
West Union		5.3 (42)	5.3 (53)		6.4 (30)	0.0 (1)	14.1 (58)		31.0 (150)	17.2 (72)		5.6 (46)
Williamsburg		0.0 (1)	0.0 (1)		0.0 (1)	12.5 (149)	25.0 (146)		16.7 (75)	33.3 (149)		0.0 (1)
Winterset		13.9 (116)	5.6 (55)		13.8 (116)	4.1 (72)	17.9 (106)		18.8 (110)	26.6 (137)		4.2 (38)
Iowa		**10.1**	**7.4**		**10.2**	**3.8**	**15.1**		**17.0**	**16.8**		**11.3**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

National (Prior Year)	**159,266**	**11.0**	**6.0**	**649,753**	**7.7**	**2.9**	**11.2**	**283,555**	**16.1**	**13.8**	**433,852**	**14.5**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)					Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Ackley		100.0 (1)		0.0 (129)		0.0 (110)		0.0 (121)		0.0 (100)		77.8 (112)		0.0 (1)
Adel		56.3 (131)		50.0 (15)		14.3 (74)		25.0 (51)		16.7 (28)		100.0 (1)		0.0 (1)
Albia		50.0 (142)		20.0 (105)		0.0 (110)		0.0 (121)		0.0 (100)		70.6 (136)		3.2 (115)
Algona		80.0 (31)		28.6 (68)		25.0 (24)		19.0 (77)		4.8 (83)		84.7 (66)		2.2 (101)
Allen WHC		74.5 (54)		20.0 (105)		4.8 (108)		17.2 (85)		2.6 (96)		70.4 (137)		1.7 (84)
Allison		73.7 (60)		36.4 (38)		40.0 (9)		25.0 (51)		9.1 (61)		89.1 (31)		0.0 (1)
Altoona		81.5 (27)		35.7 (41)		25.0 (24)		23.1 (62)		14.3 (35)		100.0 (1)		0.0 (1)
Ames		78.7 (36)		37.2 (37)		22.1 (48)		26.2 (47)		14.0 (41)		82.2 (84)		0.7 (53)
Anamosa		76.7 (42)		16.7 (117)		36.4 (14)		25.0 (51)		16.7 (28)		84.9 (65)		2.5 (106)
Ankeny		78.0 (38)		42.9 (28)		27.0 (23)		29.6 (39)		9.5 (60)		66.7 (144)		1.7 (84)
Atlantic		62.2 (110)		13.3 (123)		37.5 (12)		6.7 (118)		6.7 (72)		89.0 (34)		0.0 (1)
Audubon		61.5 (112)		66.7 (8)		0.0 (110)		0.0 (121)		0.0 (100)		98.1 (10)		5.0 (131)
Avoca		14.3 (147)		0.0 (129)		0.0 (110)		0.0 (121)		0.0 (100)		71.4 (131)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Bedford		55.6 (132)		33.3 (45)		0.0 (110)		0.0 (121)		16.7 (28)		80.0 (93)		0.0 (1)
Belle Plaine		85.7 (18)		75.0 (6)		33.3 (16)		66.7 (1)		0.0 (100)		83.3 (73)		0.0 (1)
Bloomfield		71.4 (74)		57.1 (13)		50.0 (6)		62.5 (3)		57.1 (1)		87.0 (44)		0.0 (1)
Boone		73.5 (64)		37.5 (34)		14.3 (74)		50.0 (6)		25.0 (13)		82.5 (80)		1.3 (76)
Brdlwns RP		79.2 (35)		29.2 (67)		17.4 (66)		13.5 (99)		3.4 (92)		90.0 (25)		0.9 (61)
Brooklyn		66.7 (93)		100.0 (1)		25.0 (24)		0.0 (121)		0.0 (100)		83.3 (73)		0.0 (1)
Burlington		51.8 (141)		17.2 (116)		9.6 (95)		13.3 (100)		10.9 (57)		77.9 (111)		3.5 (117)
CHC		66.3 (98)		21.2 (103)		13.3 (84)		10.2 (110)		3.4 (92)		70.2 (139)		2.7 (108)
CR Urban		69.4 (81)		30.4 (63)		20.6 (55)		11.7 (104)		6.5 (76)		84.1 (71)		1.4 (79)
Carlisle		72.0 (72)		28.6 (68)		14.3 (74)		37.5 (20)		14.3 (35)		81.8 (87)		0.0 (1)
Carroll		73.6 (63)		17.6 (114)		7.7 (99)		25.4 (50)		3.9 (89)		81.9 (85)		0.9 (61)
Cedar Falls		72.7 (69)		83.3 (5)		40.0 (9)		50.0 (6)		16.7 (28)		88.1 (37)		0.0 (1)
Centervill		53.7 (139)		7.1 (126)		21.4 (50)		33.3 (26)		0.0 (100)		71.4 (131)		7.5 (140)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Chariton		65.6 (102)		43.8 (25)		0.0 (110)		22.2 (64)		25.0 (13)		81.1 (90)		0.9 (61)
Charles City		67.8 (89)		36.4 (38)		16.7 (67)		19.2 (75)		4.5 (85)		89.1 (31)		1.4 (79)
Cherokee		77.8 (39)		23.5 (96)		13.0 (87)		21.4 (69)		5.9 (79)		85.2 (61)		4.1 (125)
Clarinda		68.4 (84)		40.0 (30)		0.0 (110)		11.1 (105)		20.0 (24)		100.0 (1)		2.0 (96)
Clarion		54.2 (136)		11.8 (124)		21.4 (50)		10.0 (111)		0.0 (100)		76.8 (118)		2.2 (101)
Clinton		68.0 (88)		22.9 (98)		17.6 (64)		14.1 (98)		4.3 (87)		83.6 (72)		1.3 (76)
Colfax		61.5 (112)		25.0 (83)		25.0 (24)		16.7 (86)		0.0 (100)		85.0 (64)		0.0 (1)
Columbus J		75.0 (49)		40.0 (30)		0.0 (110)		25.0 (51)		0.0 (100)		79.3 (99)		1.5 (82)
Corning		42.9 (146)		20.0 (105)		0.0 (110)		37.5 (20)		0.0 (100)		84.2 (69)		2.4 (105)
Corydon		72.0 (72)		33.3 (45)		25.0 (24)		37.5 (20)		33.3 (5)		63.9 (146)		3.7 (118)
Council Blfs		72.5 (70)		19.1 (112)		12.8 (89)		20.4 (71)		8.5 (64)		76.1 (121)		1.8 (92)
Cresco		88.9 (10)		20.0 (105)		16.7 (67)		33.3 (26)		6.7 (72)		86.6 (47)		17.2 (149)
Creston		54.8 (134)		28.6 (68)		0.0 (110)		22.7 (63)		14.3 (35)		86.2 (49)		0.8 (55)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Dallas Ctr		100.0 (1)		100.0 (1)		25.0 (24)		0.0 (121)		0.0 (100)		0.0 (149)		0.0 (1)
Dayton		0.0 (148)		0.0 (129)		0.0 (110)		0.0 (121)		0.0 (100)		0.0 (149)		0.0 (1)
Decorah		86.2 (17)		40.0 (30)		20.0 (56)		41.7 (15)		13.3 (42)		92.0 (18)		11.5 (143)
Denison		67.8 (89)		13.9 (122)		6.5 (106)		12.5 (102)		8.3 (66)		80.2 (92)		1.0 (68)
Dewitt		73.7 (60)		0.0 (129)		0.0 (110)		25.0 (51)		0.0 (100)		75.0 (123)		0.0 (1)
Dubuque		58.5 (127)		24.2 (94)		6.8 (104)		13.1 (101)		5.8 (80)		95.4 (14)		0.4 (48)
Dyersville		57.1 (129)		33.3 (45)		100.0 (1)		0.0 (121)		33.3 (5)		88.9 (35)		0.0 (1)
Eagle Grove		58.6 (126)		30.0 (64)		22.2 (46)		0.0 (121)		0.0 (100)		79.7 (97)		4.8 (128)
Edgerton		66.1 (99)		20.5 (104)		12.5 (90)		15.9 (89)		2.9 (94)		70.2 (139)		1.8 (92)
Eldora		58.8 (125)		33.3 (45)		11.1 (93)		33.3 (26)		0.0 (100)		90.9 (22)		1.7 (84)
Elkader		89.5 (9)		25.0 (83)		20.0 (56)		26.3 (46)		16.7 (28)		84.3 (68)		12.7 (145)
Emmetsburg		74.2 (55)		33.3 (45)		14.3 (74)		11.1 (105)		8.3 (66)		83.3 (73)		11.2 (142)
Esthervill		76.0 (44)		28.6 (68)		18.2 (62)		28.0 (41)		4.8 (83)		85.5 (57)		12.0 (144)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Fairfiel		75.6 (46)		35.7 (41)		0.0 (110)		22.2 (64)		14.3 (35)		69.0 (143)		5.1 (133)
Forest City		59.3 (123)		42.9 (28)		14.3 (74)		25.0 (51)		14.3 (35)		92.4 (17)		1.4 (79)
Fort Dodge		59.1 (124)		25.6 (82)		7.6 (102)		15.6 (91)		9.1 (61)		78.2 (107)		3.7 (118)
Ft Madison		59.7 (122)		28.0 (78)		9.5 (96)		22.2 (64)		4.0 (88)		77.1 (116)		1.7 (84)
Garner		64.3 (104)		25.0 (83)		28.6 (22)		10.0 (111)		12.5 (47)		90.0 (25)		0.0 (1)
Glenwood		73.9 (58)		16.7 (117)		50.0 (6)		11.1 (105)		0.0 (100)		77.8 (112)		1.2 (75)
Grand Junction		0.0 (148)		0.0 (129)		100.0 (1)		0.0 (121)		0.0 (100)		88.9 (35)		0.0 (1)
Greenfield		60.9 (115)		28.6 (68)		33.3 (16)		42.9 (12)		28.6 (12)		74.4 (127)		0.0 (1)
Grinnell		66.7 (93)		22.2 (100)		17.6 (64)		50.0 (6)		11.1 (53)		81.6 (88)		1.7 (84)
Grundy Ctr		54.5 (135)		25.0 (83)		0.0 (110)		33.3 (26)		12.5 (47)		75.0 (123)		0.0 (1)
Guthrie Ct		68.4 (84)		20.0 (105)		0.0 (110)		16.7 (86)		0.0 (100)		78.0 (109)		1.1 (71)
Hampton		69.5 (80)		33.3 (45)		0.0 (110)		31.3 (37)		16.7 (28)		90.8 (23)		0.4 (48)
Harlan		80.5 (30)		17.6 (114)		18.2 (62)		37.5 (20)		0.0 (100)		90.1 (24)		2.5 (106)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Hawarden		88.5 (12)		40.0 (30)		0.0 (110)		25.0 (51)		0.0 (100)		86.4 (48)		1.8 (92)
Hispanic Cln		87.5 (13)		50.0 (15)		25.0 (24)		0.0 (121)		0.0 (100)		89.5 (28)		0.0 (1)
Humboldt		83.9 (23)		60.0 (10)		20.0 (56)		42.9 (12)		20.0 (24)		85.9 (51)		0.8 (55)
ILH		68.6 (83)		28.2 (77)		15.3 (72)		8.9 (115)		2.3 (97)		97.0 (12)		0.2 (47)
Ida Grove		77.8 (39)		0.0 (129)		25.0 (24)		42.9 (12)		0.0 (100)		85.7 (52)		0.0 (1)
Independen		60.5 (116)		18.8 (113)		21.1 (52)		10.0 (111)		13.3 (42)		75.0 (123)		0.6 (51)
Indianola		66.7 (93)		35.7 (41)		13.3 (84)		18.5 (78)		17.9 (27)		74.1 (128)		3.7 (118)
Iowa City		80.8 (28)		44.7 (22)		33.8 (15)		25.9 (49)		12.2 (51)		79.2 (100)		0.9 (61)
Iowa Falls		82.1 (26)		25.0 (83)		7.7 (99)		28.6 (40)		12.5 (47)		84.2 (69)		2.0 (96)
Jamaica		60.0 (117)		0.0 (129)		0.0 (110)		0.0 (121)		0.0 (100)		73.7 (129)		0.0 (1)
Jefferson		80.8 (28)		66.7 (8)		0.0 (110)		0.0 (121)		0.0 (100)		79.6 (98)		4.2 (126)
KR		64.7 (103)		33.3 (45)		12.5 (90)		0.0 (121)		33.3 (5)		80.0 (93)		0.0 (1)
Keokuk		58.0 (128)		5.3 (128)		8.3 (97)		4.5 (120)		5.3 (81)		80.6 (91)		5.3 (134)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Keosauqua		61.1 (114)		25.0 (83)		50.0 (6)		12.5 (102)		25.0 (13)		85.7 (52)		2.0 (96)
Knoxville		67.7 (91)		32.3 (56)		23.1 (44)		26.5 (45)		12.9 (46)		75.9 (122)		2.7 (108)
LCPH		69.1 (82)		26.0 (81)		7.9 (98)		18.5 (78)		6.0 (78)		89.2 (30)		1.7 (84)
Lake City		80.0 (31)		0.0 (129)		0.0 (110)		0.0 (121)		0.0 (100)		100.0 (1)		0.0 (1)
Lamoni		50.0 (142)		0.0 (129)		0.0 (110)		0.0 (121)		0.0 (100)		77.8 (112)		3.4 (116)
Le Mars		90.3 (5)		50.0 (15)		4.5 (109)		25.0 (51)		31.8 (10)		85.3 (58)		0.4 (48)
Lenox		75.0 (49)		28.6 (68)		0.0 (110)		0.0 (121)		0.0 (100)		100.0 (1)		6.5 (139)
Leon		69.6 (79)		0.0 (129)		25.0 (24)		0.0 (121)		0.0 (100)		78.8 (104)		3.9 (124)
Logan		75.0 (49)		58.3 (12)		25.0 (24)		33.3 (26)		50.0 (2)		85.2 (61)		0.0 (1)
Manchester		53.8 (138)		0.0 (129)		0.0 (110)		0.0 (121)		0.0 (100)		97.2 (11)		2.7 (108)
Manning		100.0 (1)		0.0 (129)		0.0 (110)		0.0 (121)		0.0 (100)		66.7 (144)		0.0 (1)
Maquoketa		63.9 (107)		16.7 (117)		25.0 (24)		8.3 (116)		0.0 (100)		82.4 (82)		3.8 (123)
Marengo		71.4 (74)		44.4 (23)		25.0 (24)		33.3 (26)		22.2 (22)		86.2 (49)		2.0 (96)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Marion	211	73.9 (58)	72	25.0 (83)	61	24.6 (42)	75	10.7 (108)	72	2.8 (95)	371	87.1 (43)	953	0.8 (55)
Marshalltown	296	74.0 (57)	113	23.9 (95)	116	12.9 (88)	90	7.8 (117)	112	1.8 (98)	329	79.9 (96)	1,038	0.8 (55)
Mason City	251	65.7 (101)	98	28.6 (68)	78	14.1 (82)	108	19.4 (74)	98	13.3 (42)	512	87.7 (38)	1,051	1.0 (68)
Monticello	8	50.0 (142)	5	20.0 (105)	3	33.3 (16)	3	0.0 (121)	5	0.0 (100)	12	91.7 (20)	38	5.3 (134)
Mt Ayr	18	72.2 (71)	7	28.6 (68)	3	0.0 (110)	9	55.6 (4)	7	14.3 (35)	23	69.6 (141)	47	4.3 (127)
Mt Pleasan	48	75.0 (49)	23	21.7 (102)	19	0.0 (110)	27	33.3 (26)	23	13.0 (45)	119	83.2 (76)	257	1.6 (83)
Mtown Dtown	0	0.0 (148)	1	0.0 (129)	1	0.0 (110)	0	0.0 (121)	1	0.0 (100)	2	100.0 (1)	4	0.0 (1)
Muscatine	210	72.9 (68)	88	28.4 (76)	67	20.9 (53)	88	18.2 (80)	88	1.1 (99)	459	78.2 (107)	1,050	1.9 (95)
Nevada	10	60.0 (117)	1	0.0 (129)	5	20.0 (56)	5	40.0 (16)	1	0.0 (100)	17	82.4 (82)	39	0.0 (1)
New Hampto	20	90.0 (7)	8	37.5 (34)	12	25.0 (24)	8	25.0 (51)	8	0.0 (100)	34	85.3 (58)	105	0.0 (1)
Newton	89	67.4 (92)	29	31.0 (59)	26	15.4 (71)	26	19.2 (75)	29	10.3 (58)	126	82.5 (80)	347	1.7 (84)
Northwood	9	88.9 (10)	4	25.0 (83)	2	0.0 (110)	3	66.7 (1)	4	25.0 (13)	32	87.5 (39)	57	0.0 (1)
Norwalk	10	60.0 (117)	4	75.0 (6)	5	0.0 (110)	5	40.0 (16)	4	50.0 (2)	31	71.0 (134)	43	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Oakridge	3	66.7 (93)	4	25.0 (83)	4	25.0 (24)	2	50.0 (6)	4	25.0 (13)	4	100.0 (1)	61	0.0 (1)
Oelwein	24	54.2 (136)	17	5.9 (127)	7	14.3 (74)	11	0.0 (121)	17	0.0 (100)	61	85.2 (61)	138	28.3 (151)
Onawa	33	75.8 (45)	4	50.0 (15)	5	40.0 (9)	6	16.7 (86)	4	50.0 (2)	65	70.8 (135)	120	0.8 (55)
Orange Cty	31	90.3 (5)	13	30.8 (60)	20	20.0 (56)	7	14.3 (94)	13	7.7 (69)	68	91.2 (21)	149	0.7 (53)
Osage	41	75.6 (46)	10	30.0 (64)	7	14.3 (74)	9	55.6 (4)	10	30.0 (11)	86	91.9 (19)	176	1.7 (84)
Osceola	59	62.7 (109)	21	47.6 (21)	17	23.5 (43)	22	27.3 (42)	21	23.8 (21)	83	81.9 (85)	198	0.0 (1)
Oskaloosa	75	64.0 (106)	26	11.5 (125)	10	0.0 (110)	30	20.0 (72)	26	7.7 (69)	77	79.2 (100)	188	1.1 (71)
Ottumwa	390	68.2 (86)	129	32.6 (55)	96	20.8 (54)	163	20.9 (70)	129	8.5 (64)	428	78.7 (105)	1,093	3.1 (113)
Pella	23	78.3 (37)	10	20.0 (105)	8	0.0 (110)	5	40.0 (16)	10	0.0 (100)	50	80.0 (93)	66	3.0 (112)
Perry	67	73.1 (65)	11	36.4 (38)	32	21.9 (49)	19	15.8 (90)	11	9.1 (61)	4	50.0 (148)	227	0.9 (61)
Pocahontas	8	87.5 (13)	4	25.0 (83)	4	25.0 (24)	6	0.0 (121)	4	25.0 (13)	34	85.3 (58)	43	0.0 (1)
Postville	42	83.3 (24)	16	31.3 (57)	8	62.5 (5)	13	30.8 (38)	16	12.5 (47)	109	87.2 (42)	214	3.7 (118)
Red Oak	44	77.3 (41)	9	33.3 (45)	8	25.0 (24)	13	15.4 (92)	9	11.1 (53)	95	87.4 (41)	183	1.1 (71)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Redfield	8	87.5 (13)	1	0.0 (129)	5	0.0 (110)	0	0.0 (121)	1	0.0 (100)	0	0.0 (149)	19	0.0 (1)
Refugee Cln	14	100.0 (1)	5	100.0 (1)	2	100.0 (1)	5	0.0 (121)	5	0.0 (100)	32	62.5 (147)	62	0.0 (1)
Rock Rapid	31	74.2 (55)	12	50.0 (15)	8	37.5 (12)	15	26.7 (44)	12	25.0 (13)	59	78.0 (109)	117	0.9 (61)
Rock Valle	40	87.5 (13)	20	30.0 (64)	13	7.7 (99)	13	38.5 (19)	20	10.0 (59)	77	87.0 (44)	167	0.6 (51)
Rockwell C	16	75.0 (49)	5	0.0 (129)	8	0.0 (110)	6	33.3 (26)	5	0.0 (100)	54	87.0 (44)	82	4.9 (130)
SDM	411	73.0 (66)	154	30.5 (62)	219	13.2 (86)	129	14.7 (93)	154	3.9 (89)	14	71.4 (131)	788	1.3 (76)
Sac City	7	85.7 (18)	3	33.3 (45)	4	25.0 (24)	4	0.0 (121)	3	33.3 (5)	33	97.0 (12)	54	3.7 (118)
Sheldon	41	85.4 (21)	16	31.3 (57)	14	7.1 (103)	17	17.6 (82)	16	0.0 (100)	101	89.1 (31)	208	15.4 (147)
Shenandoah	29	55.2 (133)	17	35.3 (44)	7	0.0 (110)	16	37.5 (20)	17	11.8 (52)	54	81.5 (89)	120	5.0 (131)
Sibley	14	85.7 (18)	9	44.4 (23)	6	33.3 (16)	7	14.3 (94)	9	22.2 (22)	38	89.5 (28)	72	2.8 (111)
Sidney	14	64.3 (104)	3	0.0 (129)	3	0.0 (110)	2	50.0 (6)	3	0.0 (100)	28	85.7 (52)	51	2.0 (96)
Sigourney	21	57.1 (129)	5	0.0 (129)	1	0.0 (110)	5	20.0 (72)	5	0.0 (100)	30	76.7 (119)	53	0.0 (1)
Sioux Cent	69	79.7 (34)	15	33.3 (45)	29	31.0 (20)	17	17.6 (82)	15	6.7 (72)	131	83.2 (76)	284	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Sioux City	713	69.7 (78)	231	27.3 (80)	191	13.6 (83)	241	10.0 (111)	229	6.6 (75)	1,345	78.9 (102)	2,841	1.0 (68)
Spencer	73	65.8 (100)	26	30.8 (60)	24	29.2 (21)	27	22.2 (64)	26	7.7 (69)	158	89.9 (27)	338	15.4 (147)
Spirit Lak	41	82.9 (25)	12	16.7 (117)	15	6.7 (105)	19	31.6 (36)	12	16.7 (28)	57	94.7 (15)	148	10.8 (141)
Storm Lake	178	73.0 (66)	60	25.0 (83)	37	16.2 (70)	72	18.1 (81)	59	5.1 (82)	369	83.2 (76)	746	6.3 (138)
Stuart	7	71.4 (74)	9	22.2 (100)	6	0.0 (110)	7	14.3 (94)	9	11.1 (53)	13	84.6 (67)	33	0.0 (1)
Tama	38	73.7 (60)	22	22.7 (99)	15	20.0 (56)	23	26.1 (48)	22	4.5 (85)	72	76.4 (120)	180	1.1 (71)
Tama/Settl	5	60.0 (117)	2	0.0 (129)	2	0.0 (110)	2	0.0 (121)	2	0.0 (100)	8	75.0 (123)	18	5.6 (137)
Tipton	10	60.0 (117)	3	0.0 (129)	1	0.0 (110)	4	25.0 (51)	3	0.0 (100)	17	94.1 (16)	32	3.1 (113)
Traer	2	50.0 (142)	2	50.0 (15)	1	0.0 (110)	1	0.0 (121)	2	0.0 (100)	7	100.0 (1)	21	0.0 (1)
UIHC	10	90.0 (7)	2	100.0 (1)	1	100.0 (1)	3	33.3 (26)	2	0.0 (100)	9	77.8 (112)	18	0.0 (1)
Vinton	24	70.8 (77)	9	55.6 (14)	10	10.0 (94)	12	33.3 (26)	9	11.1 (53)	52	76.9 (117)	136	2.2 (101)
WDM	84	79.8 (33)	32	43.8 (25)	39	23.1 (44)	34	17.6 (82)	32	6.3 (77)	14	85.7 (52)	300	0.0 (1)
Washington	36	66.7 (93)	15	60.0 (10)	17	5.9 (107)	11	36.4 (25)	15	33.3 (5)	70	78.6 (106)	125	4.8 (128)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Waterloo	874	75.4 (48)	304	27.6 (79)	238	16.4 (69)	297	22.2 (64)	303	8.3 (66)	959	71.7 (130)	2,975	2.3 (104)
Waukee	46	84.8 (22)	13	15.4 (121)	42	11.9 (92)	15	6.7 (118)	13	0.0 (100)	4	100.0 (1)	201	0.0 (1)
Waukon	37	62.2 (110)	6	0.0 (129)	7	14.3 (74)	14	14.3 (94)	6	0.0 (100)	58	82.8 (79)	136	14.7 (146)
Waverly	17	76.5 (43)	16	37.5 (34)	8	25.0 (24)	11	27.3 (42)	16	18.8 (26)	46	69.6 (141)	115	0.9 (61)
Wbstr Ct	66	68.2 (86)	26	23.1 (97)	20	15.0 (73)	28	10.7 (108)	26	3.8 (91)	123	78.9 (102)	247	5.3 (134)
West Union	19	63.2 (108)	5	0.0 (129)	1	0.0 (110)	4	25.0 (51)	5	0.0 (100)	37	70.3 (138)	80	18.8 (150)
Williamsburg	0	0.0 (148)	0	0.0 (129)	1	0.0 (110)	0	0.0 (121)	0	0.0 (100)	7	85.7 (52)	8	0.0 (1)
Winterset	36	52.8 (140)	16	43.8 (25)	9	22.2 (46)	11	45.5 (11)	16	25.0 (13)	56	87.5 (39)	121	0.8 (55)
Iowa	**13,082**	**71.4**	**4,623**	**28.3**	**4,184**	**16.7**	**4,601**	**18.1**	**4,610**	**7.6**	**17,660**	**81.3**	**46,261**	**2.2**
National (Prior Year)	**168,373**	**71.4**	**88,368**	**31.4**	**91,841**	**23.5**	**60,230**	**17.4**	**71,191**	**8.1**	**285,831**	**82.0**	**655,593**	**3.7**

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ackley	17	5.9 (33)	23.5 (155)	74	5.4 (16)	1.4 (21)	10.8 (16)	32	18.8 (113)	6.3 (6)	25	16.0 (139)
Adel	50	2.0 (11)	12.0 (136)	232	4.7 (13)	3.0 (88)	13.4 (59)	120	23.3 (148)	14.2 (52)	99	6.1 (32)
Albia	71	5.6 (27)	8.5 (94)	310	6.8 (38)	4.8 (140)	13.2 (54)	134	20.1 (133)	19.4 (119)	134	11.2 (107)
Algona	141	7.8 (64)	10.6 (121)	692	6.1 (26)	1.4 (21)	14.6 (84)	313	24.3 (150)	18.8 (113)	328	5.8 (30)
Allen WHC	314	8.0 (67)	5.4 (39)	1,012	7.8 (67)	4.2 (134)	13.9 (71)	343	15.5 (51)	17.8 (104)	346	9.2 (83)
Allison	83	13.3 (139)	10.8 (125)	371	8.9 (83)	2.7 (72)	10.8 (16)	174	16.1 (60)	14.4 (54)	158	8.2 (71)
Altoona	126	7.1 (48)	4.8 (32)	601	6.8 (38)	2.2 (47)	12.5 (41)	267	21.0 (138)	13.9 (48)	249	7.6 (56)
Ames	746	10.1 (102)	8.4 (92)	3,377	9.8 (112)	4.2 (134)	12.5 (41)	1,442	13.9 (33)	12.7 (33)	1,100	14.6 (132)
Anamosa	95	8.4 (76)	5.3 (37)	397	7.6 (60)	2.3 (55)	13.1 (53)	163	16.0 (59)	17.2 (93)	222	14.9 (135)
Ankeny	228	11.4 (122)	8.3 (89)	945	8.0 (70)	1.9 (38)	10.6 (14)	411	12.7 (18)	12.9 (35)	383	7.6 (56)
Atlantic	134	9.0 (86)	9.7 (111)	648	9.3 (99)	1.7 (33)	15.7 (101)	264	18.9 (117)	14.8 (63)	325	9.8 (89)
Audubon	45	6.7 (42)	11.1 (126)	260	5.0 (14)	0.4 (10)	13.5 (64)	130	19.2 (120)	13.1 (36)	122	4.1 (19)
Avoca	22	4.5 (19)	0.0 (1)	112	6.3 (27)	5.4 (145)	10.7 (15)	55	27.3 (154)	12.7 (33)	62	16.1 (140)
Bedford	27	7.4 (52)	7.4 (73)	136	10.3 (120)	1.5 (26)	19.1 (147)	67	13.4 (24)	23.9 (150)	76	13.2 (120)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Belle Plaine	27	7.4 (52)	11.1 (126)	178	3.9 (6)	1.1 (15)	12.4 (39)	91	18.7 (109)	18.7 (111)	76	6.6 (41)
Bloomfield	29	10.3 (107)	6.9 (62)	148	6.8 (38)	4.7 (139)	17.6 (132)	65	18.5 (106)	16.9 (88)	75	2.7 (13)
Boone	113	11.5 (123)	6.2 (43)	557	8.3 (72)	5.6 (148)	9.2 (9)	261	13.4 (24)	9.2 (11)	194	7.7 (58)
Brdlwns RP	4,195	10.4 (109)	6.3 (44)	16,906	7.8 (67)	1.9 (38)	12.8 (47)	7,125	16.5 (67)	14.1 (50)	7,539	7.9 (64)
Brooklyn	15	6.7 (42)	13.3 (150)	103	9.7 (111)	2.9 (81)	17.5 (130)	52	26.9 (153)	13.5 (40)	46	19.6 (153)
Burlington	693	10.1 (102)	8.4 (92)	2,409	6.9 (42)	3.7 (120)	11.4 (25)	870	14.6 (42)	14.0 (49)	645	10.4 (94)
CHC	911	13.1 (138)	7.2 (72)	4,027	8.9 (83)	2.5 (65)	12.6 (44)	1,635	16.6 (71)	16.4 (84)	2,189	16.5 (143)
CR Urban	1,845	10.0 (100)	7.9 (82)	7,881	7.5 (56)	1.8 (35)	12.0 (33)	3,317	16.1 (60)	12.5 (30)	3,786	19.2 (151)
Carlisle	62	12.9 (134)	6.5 (50)	266	11.7 (139)	3.4 (104)	19.2 (148)	127	21.3 (139)	22.8 (148)	100	8.0 (66)
Carroll	283	10.6 (113)	8.1 (84)	1,254	6.5 (30)	1.6 (29)	11.5 (27)	559	15.9 (57)	13.2 (39)	580	6.2 (34)
Cedar Falls	43	4.7 (21)	9.3 (105)	262	7.6 (60)	3.4 (104)	13.4 (59)	123	11.4 (13)	10.6 (16)	120	15.0 (136)
Centervill	118	8.5 (79)	8.5 (94)	432	8.3 (72)	3.2 (98)	16.9 (121)	178	12.9 (20)	19.1 (116)	186	10.8 (102)
Chariton	90	5.6 (27)	12.2 (140)	445	9.0 (89)	5.6 (148)	11.0 (19)	200	18.0 (99)	12.5 (30)	226	15.9 (138)
Charles City	198	9.6 (95)	12.1 (137)	951	5.9 (23)	2.2 (47)	12.3 (35)	439	18.5 (106)	14.6 (60)	472	7.0 (48)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Cherokee	136	8.1 (71)	6.6 (52)	600	7.7 (62)	2.2 (47)	18.3 (137)	274	21.5 (142)	20.1 (128)	282	9.6 (87)
Clarinda	77	6.5 (40)	3.9 (18)	402	10.4 (122)	3.5 (111)	13.7 (67)	176	19.3 (125)	13.1 (36)	203	11.8 (111)
Clarion	157	7.6 (60)	9.6 (109)	683	10.7 (128)	2.6 (69)	18.4 (138)	296	17.9 (96)	17.6 (101)	438	2.5 (10)
Clinton	613	10.1 (102)	7.0 (65)	2,449	9.8 (112)	1.6 (29)	13.8 (68)	915	18.7 (109)	15.7 (75)	1,207	10.5 (96)
Colfax	42	9.5 (94)	11.9 (135)	180	9.4 (104)	1.7 (33)	11.1 (20)	81	12.3 (16)	14.8 (63)	64	14.1 (129)
Columbus J	51	9.8 (97)	3.9 (18)	221	6.8 (38)	2.7 (72)	18.6 (140)	105	17.1 (82)	21.0 (135)	89	9.0 (80)
Corning	41	7.3 (51)	9.8 (112)	168	9.5 (106)	3.0 (88)	14.9 (90)	74	24.3 (150)	13.5 (40)	77	14.3 (130)
Corydon	63	6.3 (36)	14.3 (152)	265	9.8 (112)	5.7 (151)	16.6 (117)	111	19.8 (131)	24.3 (151)	111	13.5 (125)
Council Blfs	1,444	10.5 (112)	7.5 (74)	6,315	9.5 (106)	2.7 (72)	15.1 (92)	2,631	19.2 (120)	17.4 (97)	3,174	16.1 (140)
Cresco	91	11.0 (118)	12.1 (137)	421	7.1 (49)	3.6 (115)	14.3 (77)	181	13.8 (31)	18.8 (113)	193	8.3 (74)
Creston	286	9.1 (87)	6.6 (52)	1,145	11.9 (140)	2.8 (77)	19.0 (146)	448	17.9 (96)	18.8 (113)	567	10.8 (102)
Dallas Ctr	10	20.0 (146)	10.0 (114)	51	21.6 (154)	3.9 (125)	9.8 (11)	20	5.0 (7)	15.0 (70)	23	0.0 (1)
Dayton	1	0.0 (1)	0.0 (1)	3	0.0 (1)	0.0 (1)	0.0 (1)	0	0.0 (1)	0.0 (1)	2	0.0 (1)
Decorah	95	7.4 (52)	13.7 (151)	445	8.5 (77)	2.2 (47)	15.3 (95)	202	15.8 (53)	17.3 (95)	223	6.7 (42)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Denison	373	9.4 (92)	5.4 (39)	1,687	10.0 (116)	2.4 (61)	18.7 (143)	703	17.2 (85)	20.8 (134)	892	14.8 (134)
Dewitt	37	27.0 (153)	8.1 (84)	181	8.3 (72)	1.1 (15)	15.5 (98)	71	28.2 (155)	14.1 (50)	92	8.7 (77)
Dubuque	1,177	8.3 (73)	7.6 (77)	4,998	6.5 (30)	3.4 (104)	8.7 (7)	1,999	16.4 (65)	9.7 (14)	2,747	16.3 (142)
Dyersville	15	20.0 (146)	6.7 (57)	89	5.6 (18)	0.0 (1)	14.6 (84)	46	6.5 (8)	21.7 (141)	55	5.5 (27)
Eagle Grove	87	9.2 (89)	6.9 (62)	429	8.9 (83)	3.3 (100)	17.2 (128)	209	17.2 (85)	19.6 (121)	271	6.3 (37)
Edgerton	1,455	11.0 (118)	8.5 (94)	5,762	10.8 (131)	2.2 (47)	14.0 (72)	2,184	16.8 (75)	16.3 (81)	3,022	12.8 (117)
Eldora	40	15.0 (143)	5.0 (34)	196	7.7 (62)	4.1 (131)	16.8 (119)	94	20.2 (135)	21.3 (138)	48	16.7 (144)
Elkader	125	10.4 (109)	8.8 (100)	559	8.8 (81)	3.2 (98)	11.8 (31)	232	21.6 (143)	10.3 (15)	289	4.8 (23)
Emmetsburg	79	12.7 (133)	6.3 (44)	360	10.3 (120)	3.9 (125)	13.6 (66)	155	16.1 (60)	12.3 (28)	195	7.7 (58)
Esthervill	146	4.8 (22)	10.3 (119)	696	9.5 (106)	3.3 (100)	14.8 (87)	299	20.1 (133)	17.4 (97)	368	8.7 (77)
Fairfiel	94	4.3 (17)	10.6 (121)	415	7.5 (56)	4.1 (131)	14.9 (90)	197	17.3 (89)	17.3 (95)	201	6.5 (39)
Fayette	0	0.0 (1)	0.0 (1)	7	14.3 (150)	0.0 (1)	0.0 (1)	3	0.0 (1)	0.0 (1)	4	0.0 (1)
Forest City	142	5.6 (27)	9.2 (102)	660	6.4 (29)	2.9 (81)	12.3 (35)	291	20.3 (136)	12.0 (26)	312	6.1 (32)
Fort Dodge	708	7.6 (60)	8.1 (84)	2,739	12.4 (142)	2.1 (43)	19.3 (150)	1,066	18.8 (113)	22.0 (143)	1,571	3.0 (15)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ft Madison	215	9.8 (97)	7.9 (82)	905	9.0 (89)	2.2 (47)	14.8 (87)	373	14.5 (39)	17.7 (102)	324	8.6 (75)
Garner	90	6.7 (42)	12.2 (140)	400	4.5 (10)	2.3 (55)	11.3 (24)	187	17.1 (82)	15.0 (70)	170	8.2 (71)
Glenwood	76	11.8 (129)	3.9 (18)	317	12.6 (145)	5.0 (144)	16.1 (107)	129	13.2 (22)	19.4 (119)	168	11.3 (108)
Grand Junction	10	10.0 (100)	0.0 (1)	54	24.1 (155)	0.0 (1)	13.0 (52)	26	15.4 (49)	15.4 (74)	31	3.2 (16)
Greenfield	61	6.6 (41)	13.1 (148)	302	9.3 (99)	4.0 (130)	15.2 (94)	127	13.4 (24)	12.6 (32)	162	17.9 (148)
Grinnell	97	7.2 (50)	8.2 (87)	468	12.4 (142)	3.0 (88)	12.8 (47)	201	22.4 (147)	9.0 (10)	157	18.5 (150)
Grundy Ctr	33	6.1 (35)	12.1 (137)	201	9.0 (89)	2.5 (65)	10.9 (18)	95	16.8 (75)	11.6 (21)	90	13.3 (122)
Guthrie Ct	61	3.3 (14)	6.6 (52)	324	5.2 (15)	1.2 (18)	12.0 (33)	169	19.5 (126)	13.6 (44)	166	3.6 (17)
Hampton	159	5.0 (24)	9.4 (107)	755	6.0 (25)	2.0 (41)	15.6 (100)	339	18.0 (99)	16.5 (86)	349	6.0 (31)
Harlan	120	11.7 (127)	4.2 (23)	532	10.7 (128)	2.1 (43)	16.2 (109)	252	17.1 (82)	16.3 (81)	265	7.9 (64)
Hawarden	71	18.3 (145)	11.3 (128)	356	7.0 (47)	2.8 (77)	16.9 (121)	178	19.1 (119)	18.0 (106)	187	9.1 (82)
Hispanic Cln	24	4.2 (16)	12.5 (142)	141	8.5 (77)	0.7 (11)	17.7 (133)	82	22.0 (144)	22.0 (143)	77	6.5 (39)
Humboldt	89	6.7 (42)	6.7 (57)	421	9.5 (106)	2.4 (61)	19.2 (148)	184	17.4 (90)	25.5 (152)	249	4.4 (21)
ILH	1,293	9.9 (99)	7.0 (65)	5,924	7.2 (54)	1.6 (29)	14.0 (72)	2,552	19.7 (129)	16.0 (79)	2,693	11.7 (110)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ida Grove	89	10.1 (102)	2.2 (15)	389	7.7 (62)	3.6 (115)	17.5 (130)	186	14.5 (39)	18.3 (108)	187	6.4 (38)
Independen	121	11.6 (126)	10.7 (124)	592	9.6 (110)	3.0 (88)	14.4 (79)	282	19.5 (126)	14.9 (67)	230	13.9 (128)
Indianola	208	10.6 (113)	10.6 (121)	1,034	12.8 (146)	3.9 (125)	16.0 (106)	476	17.2 (85)	16.6 (87)	428	10.3 (92)
Iowa City	1,499	13.5 (140)	7.1 (69)	6,275	10.1 (117)	4.9 (143)	13.5 (64)	2,699	14.7 (44)	14.4 (54)	2,940	10.6 (98)
Iowa Falls	67	7.5 (58)	4.5 (28)	315	10.5 (126)	5.4 (145)	11.4 (25)	147	14.3 (36)	12.2 (27)	103	20.4 (154)
Jamaica	16	12.5 (132)	12.5 (142)	97	4.1 (7)	4.1 (131)	14.4 (79)	50	12.0 (15)	20.0 (125)	54	9.3 (86)
Jefferson	70	4.3 (17)	4.3 (25)	349	8.0 (70)	2.9 (81)	18.1 (134)	178	18.0 (99)	19.7 (123)	177	6.2 (34)
KR	61	11.5 (123)	11.5 (132)	347	6.9 (42)	3.5 (111)	12.4 (39)	187	13.4 (24)	12.3 (28)	130	13.1 (119)
Keokuk	191	7.9 (65)	4.2 (23)	802	12.1 (141)	3.1 (94)	14.5 (81)	309	18.1 (102)	17.8 (104)	316	12.0 (114)
Keosauqua	39	5.1 (25)	12.8 (145)	149	5.4 (16)	3.4 (104)	19.5 (151)	72	15.3 (48)	26.4 (153)	73	6.8 (43)
Knoxville	291	9.3 (91)	10.0 (114)	1,292	11.3 (136)	3.6 (115)	15.5 (98)	524	15.6 (52)	17.2 (93)	522	10.5 (96)
LCPH	221	10.4 (109)	6.8 (61)	711	10.1 (117)	3.7 (120)	13.4 (59)	273	14.3 (36)	15.0 (70)	165	19.4 (152)
Lake City	5	20.0 (146)	0.0 (1)	9	11.1 (133)	33.3 (156)	11.1 (20)	2	0.0 (1)	50.0 (155)	0	0.0 (1)
Lamoni	20	0.0 (1)	10.0 (114)	126	11.1 (133)	4.8 (140)	24.6 (154)	65	9.2 (10)	23.1 (149)	70	15.7 (137)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Le Mars	189	7.4 (52)	6.3 (44)	884	8.8 (81)	2.4 (61)	16.9 (121)	421	16.9 (79)	14.5 (58)	415	7.7 (58)
Lenox	32	6.3 (36)	9.4 (107)	177	14.1 (149)	1.1 (15)	16.9 (121)	85	16.5 (67)	11.8 (23)	96	17.7 (147)
Leon	75	8.0 (67)	5.3 (37)	318	10.4 (122)	2.8 (77)	14.2 (75)	130	22.3 (145)	14.6 (60)	158	18.4 (149)
Logan	102	13.7 (142)	7.8 (80)	449	11.1 (133)	2.4 (61)	17.1 (126)	190	18.9 (117)	15.8 (76)	221	16.7 (144)
Manchester	56	5.4 (26)	12.5 (142)	320	6.3 (27)	4.4 (136)	13.4 (59)	161	17.4 (90)	13.7 (45)	187	9.6 (87)
Manning	5	20.0 (146)	0.0 (1)	22	0.0 (1)	0.0 (1)	18.2 (135)	15	6.7 (9)	20.0 (125)	15	0.0 (1)
Maquoketa	106	5.7 (30)	4.7 (31)	561	8.9 (83)	1.8 (35)	12.8 (47)	242	17.8 (94)	14.5 (58)	306	6.9 (46)
Marengo	38	7.9 (65)	13.2 (149)	178	4.5 (10)	3.4 (104)	11.8 (31)	84	17.9 (96)	13.1 (36)	73	4.1 (19)
Marion	583	8.4 (76)	7.0 (65)	2,854	6.7 (35)	1.2 (18)	12.3 (35)	1,317	18.1 (102)	14.9 (67)	1,422	8.2 (71)
Marshalltown	945	8.4 (76)	7.8 (80)	3,902	5.8 (20)	3.7 (120)	13.4 (59)	1,656	17.2 (85)	14.4 (54)	1,021	13.7 (126)
Mason City	744	8.9 (83)	8.2 (87)	3,216	6.9 (42)	2.3 (55)	13.2 (54)	1,344	18.2 (104)	17.0 (89)	1,341	6.8 (43)
Monticello	35	0.0 (1)	2.9 (17)	201	4.5 (10)	3.5 (111)	9.5 (10)	94	14.9 (47)	14.9 (67)	115	10.4 (94)
Mt Ayr	42	4.8 (22)	2.4 (16)	177	12.4 (142)	2.3 (55)	18.6 (140)	80	18.8 (113)	17.5 (100)	84	10.7 (101)
Mt Pleasan	186	7.5 (58)	11.3 (128)	873	9.3 (99)	3.0 (88)	16.4 (114)	381	17.6 (92)	20.2 (129)	354	8.8 (79)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Mtown Dtown	1	0.0 (1)	0.0 (1)	11	9.1 (93)	0.0 (1)	18.2 (135)	5	20.0 (132)	20.0 (125)	2	50.0 (156)
Murray	1	100.0 (156)	0.0 (1)	7	14.3 (150)	0.0 (1)	0.0 (1)	4	0.0 (1)	0.0 (1)	6	0.0 (1)
Muscatine	786	10.2 (106)	7.5 (74)	3,165	7.1 (49)	2.3 (55)	13.8 (68)	1,194	18.7 (109)	18.6 (110)	947	12.1 (115)
Nevada	25	8.0 (67)	4.0 (21)	131	3.8 (5)	6.1 (153)	14.5 (81)	57	15.8 (53)	19.3 (118)	47	17.0 (146)
New Hampto	54	7.4 (52)	9.3 (105)	319	9.1 (93)	3.1 (94)	16.3 (113)	155	16.8 (75)	18.7 (111)	136	11.0 (104)
Newton	271	12.9 (134)	9.6 (109)	1,275	8.4 (76)	4.6 (137)	16.1 (107)	553	15.9 (57)	17.7 (102)	440	11.1 (105)
North Towne	424	9.7 (96)	8.3 (89)	1,974	7.5 (56)	1.5 (26)	11.6 (29)	812	18.5 (106)	9.4 (13)	1,181	13.4 (123)
Northwood	35	5.7 (30)	11.4 (130)	169	6.5 (30)	3.6 (115)	8.9 (8)	73	13.7 (30)	5.5 (5)	84	7.1 (50)
Norwalk	46	23.9 (152)	8.7 (97)	235	14.9 (152)	3.4 (104)	16.6 (117)	103	13.6 (29)	20.4 (132)	102	14.7 (133)
Oakridge	15	0.0 (1)	0.0 (1)	134	2.2 (4)	2.2 (47)	5.2 (5)	71	12.7 (18)	7.0 (8)	67	13.4 (123)
Oelwein	97	10.3 (107)	10.3 (119)	489	5.7 (19)	1.6 (29)	17.4 (129)	215	15.8 (53)	20.5 (133)	250	5.6 (28)
Onawa	93	12.9 (134)	4.3 (25)	416	11.3 (136)	3.1 (94)	15.9 (104)	165	15.8 (53)	21.2 (136)	196	9.2 (83)
Orange Cty	114	7.0 (47)	11.4 (130)	505	6.9 (42)	2.6 (69)	12.5 (41)	235	17.0 (81)	8.5 (9)	237	8.0 (66)
Osage	119	5.9 (33)	6.7 (57)	483	7.5 (56)	2.1 (43)	10.4 (13)	214	14.0 (35)	10.7 (17)	224	9.8 (89)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Osceola	184	10.9 (117)	6.5 (50)	739	9.1 (93)	3.8 (123)	14.7 (86)	304	18.8 (113)	16.4 (84)	355	13.8 (127)
Oskaloosa	176	8.0 (67)	6.3 (44)	812	8.9 (83)	3.6 (115)	13.3 (58)	349	17.8 (94)	13.8 (47)	419	6.9 (46)
Ottumwa	1,300	8.7 (80)	7.5 (74)	5,206	9.1 (93)	3.3 (100)	15.3 (95)	1,981	16.9 (79)	17.4 (97)	2,751	7.0 (48)
Pella	62	12.9 (134)	12.9 (146)	311	7.1 (49)	2.9 (81)	16.4 (114)	142	16.2 (63)	18.3 (108)	144	13.2 (120)
Perry	196	11.2 (121)	6.6 (52)	957	6.9 (42)	1.4 (21)	15.7 (101)	444	22.3 (145)	19.1 (116)	437	4.6 (22)
Pocahontas	32	3.1 (13)	6.3 (44)	196	13.8 (148)	5.6 (148)	16.8 (119)	108	21.3 (139)	14.8 (63)	123	8.1 (69)
Pomeroy	2	0.0 (1)	0.0 (1)	15	6.7 (35)	0.0 (1)	33.3 (156)	6	0.0 (1)	50.0 (155)	10	0.0 (1)
Postville	130	10.8 (115)	4.6 (30)	683	11.6 (138)	2.9 (81)	17.1 (126)	328	14.6 (42)	21.6 (140)	388	7.7 (58)
Red Oak	137	11.7 (127)	10.2 (117)	712	10.5 (126)	3.4 (104)	13.2 (54)	333	13.2 (22)	9.3 (12)	386	10.6 (98)
Redfield	22	9.1 (87)	0.0 (1)	78	7.7 (62)	3.8 (123)	6.4 (6)	30	23.3 (148)	6.7 (7)	40	12.5 (116)
Refugee Cln	26	3.8 (15)	11.5 (132)	191	5.8 (20)	1.0 (14)	11.5 (27)	96	16.7 (73)	13.5 (40)	111	30.6 (155)
Rock Rapid	65	7.7 (62)	9.2 (102)	367	10.4 (122)	1.9 (38)	11.7 (30)	176	10.8 (11)	10.8 (19)	172	5.2 (24)
Rock Valle	121	8.3 (73)	6.6 (52)	545	7.7 (62)	2.0 (41)	15.4 (97)	251	13.9 (33)	14.3 (53)	264	6.8 (43)
Rockwell C	49	12.2 (131)	10.2 (117)	288	9.0 (89)	1.4 (21)	18.8 (144)	151	19.2 (120)	21.2 (136)	154	2.6 (12)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
SDM	1,281	8.9 (83)	8.7 (97)	5,372	6.6 (33)	2.1 (43)	12.7 (45)	2,251	16.7 (73)	14.4 (54)	2,219	7.4 (53)
Sac City	32	21.9 (151)	0.0 (1)	212	9.4 (104)	0.9 (13)	12.7 (45)	112	20.5 (137)	16.1 (80)	118	2.5 (10)
Sheldon	142	12.0 (130)	6.3 (44)	619	9.2 (97)	3.9 (125)	16.2 (109)	253	13.8 (31)	20.2 (129)	323	9.0 (80)
Shenandoah	96	13.5 (140)	5.2 (36)	460	10.7 (128)	5.7 (151)	9.8 (11)	214	12.6 (17)	10.7 (17)	238	13.0 (118)
Sibley	48	6.3 (36)	8.3 (89)	228	6.6 (33)	2.2 (47)	14.0 (72)	102	14.7 (44)	11.8 (23)	122	7.4 (53)
Sidney	34	8.8 (82)	5.9 (42)	196	13.3 (147)	2.6 (69)	14.3 (77)	103	21.4 (141)	14.6 (60)	107	7.5 (55)
Sigourney	45	11.1 (120)	17.8 (154)	219	4.1 (7)	2.7 (72)	12.8 (47)	103	16.5 (67)	11.7 (22)	102	2.9 (14)
Sioux Cent	194	5.7 (30)	7.7 (79)	902	8.9 (83)	1.8 (35)	18.8 (144)	395	19.2 (120)	22.3 (147)	461	7.8 (62)
Sioux City	2,221	7.7 (62)	6.7 (57)	9,917	8.5 (77)	1.3 (20)	18.5 (139)	4,205	18.7 (109)	19.9 (124)	4,414	8.1 (69)
Spencer	235	6.8 (46)	5.5 (41)	1,052	7.4 (55)	2.9 (81)	13.8 (68)	431	14.8 (46)	15.8 (76)	551	8.0 (66)
Spirit Lak	119	9.2 (89)	9.2 (102)	478	7.1 (49)	4.8 (140)	12.3 (35)	193	16.6 (71)	11.9 (25)	239	3.8 (18)
Storm Lake	517	6.4 (39)	5.0 (34)	2,274	6.7 (35)	3.1 (94)	14.8 (87)	972	16.4 (65)	17.1 (92)	1,166	7.8 (62)
Stuart	28	32.1 (154)	7.1 (69)	131	20.6 (153)	1.5 (26)	23.7 (153)	59	11.9 (14)	20.3 (131)	63	11.1 (105)
Tama	138	8.7 (80)	13.0 (147)	671	5.8 (20)	2.8 (77)	14.5 (81)	292	18.2 (104)	18.2 (107)	186	11.8 (111)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Tama/Settl	8	0.0 (1)	0.0 (1)	57	7.0 (47)	3.5 (111)	26.3 (155)	26	19.2 (120)	26.9 (154)	19	5.3 (25)
Tipton	22	4.5 (19)	4.5 (28)	140	9.3 (99)	2.9 (81)	18.6 (140)	73	24.7 (152)	21.9 (142)	81	6.2 (34)
Traer	14	21.4 (150)	7.1 (69)	93	8.6 (80)	5.4 (145)	15.1 (92)	51	19.6 (128)	13.7 (45)	33	0.0 (1)
UIHC	74	81.1 (155)	4.1 (22)	107	41.1 (156)	7.5 (154)	11.2 (23)	27	11.1 (12)	14.8 (63)	38	5.3 (25)
Vinton	76	2.6 (12)	14.5 (153)	463	10.4 (122)	3.0 (88)	15.8 (103)	239	19.7 (129)	16.3 (81)	207	10.6 (98)
WDM	244	9.4 (92)	9.8 (112)	1,217	7.1 (49)	2.5 (65)	12.9 (51)	564	15.4 (49)	13.5 (40)	559	8.6 (75)
Wapello	1	0.0 (1)	0.0 (1)	1	0.0 (1)	0.0 (1)	0.0 (1)	0	0.0 (1)	0.0 (1)	0	0.0 (1)
Washington	126	7.1 (48)	8.7 (97)	597	7.9 (69)	2.3 (55)	13.2 (54)	273	17.6 (92)	15.0 (70)	278	7.2 (52)
Waterloo	2,372	10.8 (115)	7.0 (65)	9,424	9.9 (115)	3.9 (125)	14.2 (75)	3,944	16.5 (67)	17.0 (89)	3,454	14.3 (130)
Waukee	144	8.3 (73)	9.0 (101)	723	5.9 (23)	2.5 (65)	11.1 (20)	305	13.4 (24)	11.5 (20)	319	11.3 (108)
Waukon	85	8.2 (72)	11.8 (134)	365	4.4 (9)	1.4 (21)	16.4 (114)	154	16.2 (63)	22.1 (145)	173	9.2 (83)
Waverly	66	16.7 (144)	7.6 (77)	483	11.0 (132)	2.7 (72)	17.0 (125)	251	13.1 (21)	15.9 (78)	221	11.8 (111)
Wbstr Ct	217	7.4 (52)	6.9 (62)	888	10.1 (117)	3.3 (100)	15.9 (104)	382	16.8 (75)	17.0 (89)	531	5.6 (28)
West Union	61	11.5 (123)	4.9 (33)	247	9.3 (99)	0.8 (12)	16.2 (109)	90	30.0 (156)	22.2 (146)	118	10.2 (91)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Williamsburg	1	0.0 (1)	100.0 (156)	24	8.3 (72)	8.3 (155)	20.8 (152)	14	14.3 (36)	21.4 (139)	14	7.1 (50)
Winterset	90	8.9 (83)	4.4 (27)	414	9.2 (97)	4.6 (137)	16.2 (109)	179	14.5 (39)	19.6 (121)	204	10.3 (92)
Iowa	**40,055**	**9.8**	**7.6**	**173,218**	**8.5**	**2.7**	**14.1**	**73,376**	**17.1**	**16.9**	**79,105**	**10.7**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Ackley	17	82.4 (14)	11	36.4 (23)	3	0.0 (137)	6	16.7 (86)	11	0.0 (130)	22	72.7 (144)	48	2.1 (83)
Adel	49	63.3 (122)	20	35.0 (31)	17	17.6 (66)	19	15.8 (90)	20	10.0 (46)	15	93.3 (11)	94	2.1 (83)
Albia	72	54.2 (146)	23	13.0 (139)	21	9.5 (116)	20	10.0 (130)	23	0.0 (130)	90	77.8 (130)	167	3.6 (119)
Algona	143	76.2 (41)	67	26.9 (69)	37	18.9 (53)	69	20.3 (64)	67	7.5 (73)	333	83.2 (83)	686	2.6 (103)
Allen WHC	318	73.0 (61)	136	26.5 (72)	79	13.9 (91)	127	18.9 (70)	134	9.0 (56)	257	71.6 (147)	828	3.7 (120)
Allison	83	75.9 (42)	35	22.9 (98)	26	30.8 (11)	40	25.0 (44)	35	8.6 (61)	188	90.4 (19)	365	0.5 (21)
Altoona	128	74.2 (50)	79	30.4 (49)	60	16.7 (69)	50	14.0 (111)	79	6.3 (86)	13	92.3 (13)	332	0.9 (39)
Ames	762	78.5 (32)	382	38.7 (19)	393	25.7 (20)	325	28.3 (32)	382	7.9 (71)	971	84.6 (67)	2,443	1.3 (52)
Anamosa	96	71.9 (67)	45	20.0 (113)	29	24.1 (26)	48	20.8 (62)	45	11.1 (38)	144	88.2 (34)	327	3.4 (118)
Ankeny	226	77.9 (37)	101	31.7 (42)	95	22.1 (35)	98	17.3 (83)	101	6.9 (78)	36	80.6 (110)	432	1.4 (59)
Atlantic	136	61.8 (128)	65	16.9 (129)	46	19.6 (48)	68	13.2 (116)	65	3.1 (121)	138	84.1 (71)	323	1.2 (50)
Audubon	45	68.9 (86)	26	23.1 (94)	17	5.9 (132)	22	9.1 (135)	26	0.0 (130)	128	89.1 (28)	216	2.3 (93)
Avoca	22	54.5 (145)	6	33.3 (37)	7	14.3 (84)	10	40.0 (9)	6	16.7 (15)	44	79.5 (122)	80	2.5 (96)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Bedford	27	70.4 (79)	17	23.5 (89)	7	0.0 (137)	17	11.8 (120)	17	17.6 (14)	52	73.1 (143)	96	0.0 (1)
Belle Plaine	28	71.4 (70)	12	41.7 (12)	10	40.0 (4)	10	50.0 (2)	12	8.3 (67)	50	78.0 (129)	145	2.1 (83)
Bloomfield	30	66.7 (99)	17	35.3 (29)	4	25.0 (21)	18	38.9 (10)	17	35.3 (1)	24	87.5 (39)	74	0.0 (1)
Boone	115	67.8 (90)	51	31.4 (44)	66	13.6 (92)	38	42.1 (7)	51	15.7 (20)	188	88.8 (30)	433	2.1 (83)
Brdlwns RP	4,229	78.1 (35)	1,670	29.8 (53)	1,768	19.2 (52)	1,495	11.8 (120)	1,667	3.7 (117)	945	83.9 (75)	10,792	1.3 (52)
Brooklyn	17	58.8 (136)	9	33.3 (37)	11	18.2 (58)	10	0.0 (144)	9	11.1 (38)	39	87.2 (44)	94	1.1 (43)
Burlington	697	55.7 (142)	275	17.1 (127)	192	10.9 (111)	331	14.8 (99)	275	7.3 (74)	846	79.6 (121)	2,337	3.8 (121)
CHC	916	64.3 (114)	421	18.5 (122)	296	13.2 (101)	493	9.9 (133)	421	5.0 (96)	1,042	68.0 (149)	1,977	4.3 (123)
CR Urban	1,915	67.6 (91)	838	29.6 (54)	582	20.4 (42)	852	11.9 (119)	838	4.2 (112)	2,356	82.9 (87)	5,800	1.8 (74)
Carlisle	63	69.8 (83)	32	37.5 (21)	22	22.7 (31)	31	32.3 (18)	32	12.5 (34)	93	89.2 (25)	149	0.7 (28)
Carroll	285	71.6 (69)	132	17.4 (126)	88	11.4 (107)	143	18.9 (70)	132	3.8 (116)	558	82.1 (97)	1,001	1.9 (80)
Cedar Falls	44	86.4 (9)	30	70.0 (2)	14	35.7 (6)	27	55.6 (1)	30	23.3 (3)	85	85.9 (57)	218	0.5 (21)
Centervill	118	58.5 (137)	43	16.3 (133)	29	10.3 (112)	45	20.0 (65)	43	2.3 (125)	82	72.0 (145)	225	7.6 (143)
Chariton	90	66.7 (99)	50	22.0 (103)	23	8.7 (124)	33	27.3 (35)	50	14.0 (29)	128	82.8 (88)	324	1.5 (66)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Charles City	199	66.3 (104)	99	25.3 (78)	58	17.2 (67)	105	20.0 (65)	99	6.1 (88)	464	90.3 (20)	950	1.3 (52)
Cherokee	137	78.1 (35)	55	23.6 (88)	64	14.1 (88)	56	21.4 (60)	55	5.5 (93)	204	80.4 (112)	383	4.4 (127)
Clarinda	77	71.4 (70)	27	44.4 (8)	23	13.0 (103)	35	22.9 (53)	27	22.2 (4)	111	86.5 (50)	256	3.1 (113)
Clarion	157	61.8 (128)	75	25.3 (78)	55	18.2 (58)	67	14.9 (97)	74	4.1 (113)	298	83.6 (80)	607	2.6 (103)
Clinton	614	66.6 (103)	247	21.5 (107)	108	14.8 (83)	238	16.8 (85)	247	8.5 (66)	544	86.2 (51)	1,909	2.2 (91)
Colfax	42	66.7 (99)	13	23.1 (94)	14	7.1 (129)	18	11.1 (124)	13	7.7 (72)	72	86.1 (52)	160	0.0 (1)
Columbus J	51	80.4 (23)	18	44.4 (8)	11	0.0 (137)	20	10.0 (130)	17	0.0 (130)	110	83.6 (80)	226	1.3 (52)
Corning	41	58.5 (137)	19	21.1 (111)	11	18.2 (58)	22	27.3 (35)	19	5.3 (94)	56	76.8 (135)	126	4.8 (133)
Corydon	63	65.1 (110)	27	25.9 (75)	19	15.8 (78)	31	29.0 (29)	27	18.5 (11)	92	77.2 (134)	148	6.1 (140)
Council Blfs	1,473	73.7 (56)	642	21.5 (107)	546	16.1 (75)	607	19.1 (69)	641	9.2 (53)	1,718	79.0 (125)	3,677	1.6 (68)
Cresco	91	82.4 (14)	48	20.8 (112)	38	23.7 (28)	52	28.8 (30)	48	6.3 (86)	178	87.1 (45)	422	15.9 (151)
Creston	285	64.2 (116)	108	23.1 (94)	86	8.1 (126)	123	18.7 (72)	107	10.3 (44)	342	82.7 (90)	858	3.3 (117)
Dallas Ctr	10	90.0 (5)	7	71.4 (1)	9	22.2 (34)	2	0.0 (144)	7	0.0 (130)	0	0.0 (154)	22	0.0 (1)
Dayton	1	0.0 (153)	1	0.0 (147)	0	0.0 (137)	0	0.0 (144)	1	0.0 (130)	0	0.0 (154)	2	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Decorah	96	83.3 (12)	45	44.4 (8)	28	28.6 (14)	40	30.0 (25)	45	17.8 (13)	206	95.1 (9)	452	13.7 (147)
Denison	374	67.6 (91)	149	18.8 (120)	112	8.9 (123)	185	8.6 (136)	149	2.7 (123)	490	79.2 (124)	1,066	0.8 (34)
Dewitt	37	64.9 (111)	17	23.5 (89)	3	0.0 (137)	13	23.1 (49)	17	0.0 (130)	35	85.7 (58)	125	0.8 (34)
Dubuque	1,187	53.3 (147)	486	19.1 (119)	294	11.2 (108)	550	15.6 (92)	486	4.7 (102)	1,911	96.1 (8)	4,510	0.2 (19)
Dyersville	15	53.3 (147)	8	25.0 (81)	5	60.0 (1)	7	14.3 (105)	8	12.5 (34)	41	97.6 (7)	77	0.0 (1)
Eagle Grove	87	64.4 (113)	44	15.9 (136)	27	18.5 (55)	37	5.4 (142)	44	0.0 (130)	214	81.3 (106)	381	3.1 (113)
Edgerton	1,462	66.8 (98)	595	22.2 (100)	400	13.3 (96)	719	14.3 (105)	593	4.6 (105)	959	71.8 (146)	1,813	2.5 (96)
Eldora	40	55.0 (143)	16	18.8 (120)	22	4.5 (133)	15	13.3 (114)	16	0.0 (130)	62	93.5 (10)	147	5.4 (137)
Elkader	126	86.5 (8)	58	25.9 (75)	27	3.7 (135)	74	23.0 (52)	58	8.6 (61)	241	89.2 (25)	570	15.8 (150)
Emmetsburg	79	78.5 (32)	43	27.9 (64)	28	17.9 (64)	38	15.8 (90)	43	9.3 (52)	158	85.4 (60)	355	11.3 (144)
Esthervill	147	67.3 (95)	70	27.1 (68)	45	20.0 (43)	63	25.4 (43)	70	8.6 (61)	326	85.6 (59)	699	13.2 (146)
Fairfiel	94	66.0 (105)	36	27.8 (66)	23	0.0 (137)	37	21.6 (59)	36	13.9 (31)	83	67.5 (150)	212	5.7 (139)
Fayette	0	0.0 (153)	0	0.0 (147)	0	0.0 (137)	0	0.0 (144)	0	0.0 (130)	2	50.0 (153)	5	20.0 (156)
Forest City	143	61.5 (130)	60	31.7 (42)	29	10.3 (112)	79	22.8 (54)	60	8.3 (67)	316	88.6 (31)	662	1.4 (59)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Fort Dodge	713	63.0 (125)	285	20.0 (113)	196	11.7 (106)	293	15.0 (95)	284	6.7 (80)	1,098	82.7 (90)	2,387	3.9 (122)
Ft Madison	217	56.2 (141)	89	18.0 (123)	70	14.3 (84)	110	14.5 (104)	89	4.5 (106)	379	75.7 (140)	908	2.5 (96)
Garner	92	64.1 (117)	36	22.2 (100)	19	15.8 (78)	44	18.2 (76)	36	11.1 (38)	190	87.9 (37)	392	0.8 (34)
Glenwood	76	72.4 (63)	30	20.0 (113)	18	16.7 (69)	34	14.7 (100)	30	3.3 (119)	51	80.4 (112)	126	0.8 (34)
Grand Junction	10	70.0 (81)	6	33.3 (37)	6	33.3 (8)	4	50.0 (2)	6	0.0 (130)	25	88.0 (36)	47	6.4 (141)
Greenfield	61	63.9 (119)	24	29.2 (56)	18	27.8 (17)	29	31.0 (20)	24	16.7 (15)	118	82.2 (95)	238	2.1 (83)
Grinnell	97	68.0 (88)	55	20.0 (113)	57	14.0 (90)	39	17.9 (78)	55	9.1 (54)	157	81.5 (104)	383	2.9 (110)
Grundy Ctr	33	60.6 (134)	23	30.4 (49)	15	13.3 (96)	19	10.5 (127)	23	8.7 (60)	58	77.6 (132)	154	0.6 (26)
Guthrie Ct	61	72.1 (65)	23	21.7 (105)	22	13.6 (92)	28	17.9 (78)	23	0.0 (130)	169	78.7 (128)	272	2.6 (103)
Hampton	163	71.2 (74)	57	35.1 (30)	44	6.8 (130)	66	30.3 (23)	57	14.0 (29)	359	89.4 (24)	738	1.4 (59)
Harlan	121	71.1 (76)	50	16.0 (135)	39	7.7 (128)	58	27.6 (34)	50	4.0 (114)	144	86.8 (48)	290	2.1 (83)
Hawarden	72	83.3 (12)	27	29.6 (54)	38	2.6 (136)	26	30.8 (22)	27	3.7 (117)	135	87.4 (42)	243	1.6 (68)
Hispanic Cln	24	75.0 (43)	10	40.0 (16)	14	35.7 (6)	14	7.1 (140)	10	0.0 (130)	77	89.6 (22)	132	0.0 (1)
Humboldt	89	79.8 (28)	33	36.4 (23)	28	17.9 (64)	36	33.3 (14)	33	21.2 (5)	184	83.7 (78)	377	2.4 (94)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
ILH	1,298	73.1 (60)	604	25.8 (77)	662	16.5 (74)	509	10.2 (128)	602	3.2 (120)	548	86.1 (52)	4,797	1.3 (52)
Ida Grove	89	79.8 (28)	36	11.1 (141)	31	16.1 (75)	31	25.8 (38)	36	8.3 (67)	146	89.0 (29)	280	0.7 (28)
Independen	121	63.6 (120)	64	21.9 (104)	44	20.5 (41)	45	22.2 (56)	63	4.8 (100)	138	84.1 (71)	437	2.5 (96)
Indianola	208	73.6 (57)	89	33.7 (35)	72	20.8 (40)	104	26.9 (37)	89	9.0 (56)	372	83.9 (75)	624	2.9 (110)
Iowa City	1,511	80.6 (22)	612	45.4 (6)	548	31.9 (10)	635	25.8 (38)	607	11.5 (37)	1,195	78.8 (127)	2,724	1.0 (41)
Iowa Falls	66	80.3 (24)	35	42.9 (11)	37	8.1 (126)	28	50.0 (2)	35	14.3 (26)	109	88.1 (35)	264	1.1 (43)
Jamaica	16	81.3 (20)	2	0.0 (147)	11	0.0 (137)	7	0.0 (144)	2	0.0 (130)	46	73.9 (142)	76	1.3 (52)
Jefferson	71	74.6 (46)	28	21.4 (109)	19	21.1 (39)	37	8.1 (138)	28	0.0 (130)	182	81.3 (106)	283	2.1 (83)
KR	64	73.4 (59)	26	30.8 (46)	33	18.2 (58)	24	8.3 (137)	26	15.4 (22)	69	82.6 (92)	264	1.1 (43)
Keokuk	193	52.8 (150)	90	12.2 (140)	49	10.2 (114)	89	13.5 (112)	90	4.4 (108)	304	79.9 (118)	780	5.3 (135)
Keosauqua	39	64.1 (117)	11	36.4 (23)	11	18.2 (58)	21	14.3 (105)	11	18.2 (12)	23	87.0 (46)	73	2.7 (106)
Knoxville	295	67.1 (97)	139	26.6 (70)	85	18.8 (54)	151	23.2 (48)	139	8.6 (61)	362	79.0 (125)	668	4.3 (123)
LCPH	223	69.1 (85)	64	26.6 (70)	46	8.7 (124)	85	17.6 (80)	64	4.7 (102)	259	88.4 (32)	673	1.8 (74)
Lake City	5	80.0 (25)	1	0.0 (147)	1	0.0 (137)	0	0.0 (144)	1	0.0 (130)	2	100.0 (1)	3	0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Lamoni	20	75.0 (43)	9	33.3 (37)	11	0.0 (137)	5	0.0 (144)	9	11.1 (38)	50	86.0 (56)	104	1.9 (80)
Le Mars	194	87.1 (6)	64	39.1 (18)	75	13.3 (96)	76	18.4 (74)	64	15.6 (21)	296	81.4 (105)	574	0.5 (21)
Lenox	32	62.5 (126)	17	17.6 (125)	12	0.0 (137)	14	7.1 (140)	17	5.9 (90)	69	84.1 (71)	138	2.2 (91)
Leon	75	61.3 (131)	31	9.7 (142)	17	41.2 (3)	38	18.4 (74)	31	0.0 (130)	67	80.6 (110)	216	3.2 (115)
Logan	103	77.7 (39)	48	35.4 (28)	33	24.2 (25)	53	35.8 (12)	48	25.0 (2)	89	84.3 (69)	212	0.5 (21)
Manchester	56	41.1 (152)	26	3.8 (146)	25	0.0 (137)	24	12.5 (118)	26	0.0 (130)	155	98.1 (6)	298	0.7 (28)
Manning	5	80.0 (25)	0	0.0 (147)	2	0.0 (137)	2	50.0 (2)	0	0.0 (130)	15	80.0 (116)	19	0.0 (1)
Maquoketa	106	61.3 (131)	31	16.1 (134)	22	9.1 (117)	38	7.9 (139)	31	6.5 (84)	136	83.1 (84)	475	2.5 (96)
Marengo	36	72.2 (64)	20	30.0 (52)	13	23.1 (29)	18	16.7 (86)	20	10.0 (46)	37	89.2 (25)	88	1.1 (43)
Marion	588	69.2 (84)	268	26.5 (72)	211	19.9 (47)	272	14.7 (100)	268	4.9 (99)	1,057	85.2 (62)	2,721	1.6 (68)
Marshalltown	958	73.6 (57)	442	21.3 (110)	412	12.4 (104)	343	12.8 (117)	441	1.8 (128)	1,272	80.0 (116)	2,987	1.8 (74)
Mason City	756	66.0 (105)	326	25.2 (80)	231	15.2 (82)	339	16.5 (88)	326	10.1 (45)	1,425	87.5 (39)	3,119	1.7 (72)
Monticello	35	65.7 (108)	15	20.0 (113)	11	9.1 (117)	10	30.0 (25)	15	6.7 (80)	75	93.3 (11)	156	4.5 (131)
Mt Ayr	42	66.7 (99)	20	35.0 (31)	11	9.1 (117)	25	44.0 (6)	20	10.0 (46)	68	76.5 (136)	130	1.5 (66)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Mt Pleasan	187	70.1 (80)	89	32.6 (41)	72	9.7 (115)	100	31.0 (20)	89	9.0 (56)	385	83.1 (84)	845	2.5 (96)
Mtown Dtown	1	100.0 (1)	2	0.0 (147)	3	0.0 (137)	1	0.0 (144)	2	0.0 (130)	4	75.0 (141)	10	0.0 (1)
Murray	1	0.0 (153)	0	0.0 (147)	0	0.0 (137)	1	0.0 (144)	0	0.0 (130)	3	100.0 (1)	6	16.7 (153)
Muscatine	790	74.2 (50)	354	23.4 (91)	282	15.6 (80)	384	14.1 (109)	354	2.3 (125)	1,363	82.2 (95)	3,349	2.4 (94)
Nevada	24	58.3 (139)	12	8.3 (145)	13	30.8 (11)	12	25.0 (44)	12	0.0 (130)	48	89.6 (22)	107	1.9 (80)
New Hampto	55	81.8 (17)	29	37.9 (20)	28	32.1 (9)	27	33.3 (14)	29	13.8 (32)	82	84.1 (71)	268	0.7 (28)
Newton	273	71.8 (68)	134	23.9 (86)	112	13.4 (95)	114	14.9 (97)	134	5.2 (95)	471	83.0 (86)	1,086	2.1 (83)
North Towne	430	70.9 (77)	199	24.1 (85)	147	14.3 (84)	223	13.5 (112)	199	5.0 (96)	661	88.4 (32)	1,738	1.4 (59)
Northwood	35	74.3 (49)	21	23.8 (87)	14	28.6 (14)	18	33.3 (14)	21	9.5 (51)	80	82.5 (93)	162	0.6 (26)
Norwalk	46	67.4 (94)	20	45.0 (7)	15	6.7 (131)	22	31.8 (19)	20	20.0 (6)	84	77.4 (133)	145	0.7 (28)
Oakridge	15	53.3 (147)	9	22.2 (100)	15	13.3 (96)	7	14.3 (105)	9	11.1 (38)	5	100.0 (1)	107	0.0 (1)
Oelwein	97	68.0 (88)	60	15.0 (137)	26	3.8 (134)	64	15.6 (92)	60	6.7 (80)	218	85.3 (61)	498	22.1 (157)
Onawa	95	63.2 (124)	30	16.7 (130)	19	26.3 (19)	31	25.8 (38)	30	10.0 (46)	83	67.5 (150)	186	1.1 (43)
Orange Cty	114	86.0 (10)	36	36.1 (26)	52	30.8 (11)	36	27.8 (33)	36	8.3 (67)	161	90.7 (17)	314	1.3 (52)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Osage	120	74.2 (50)	43	27.9 (64)	27	18.5 (55)	48	41.7 (8)	43	16.3 (18)	223	91.5 (16)	484	1.7 (72)
Osceola	183	67.2 (96)	63	30.2 (51)	54	16.7 (69)	78	23.1 (49)	63	11.1 (38)	234	81.2 (108)	548	0.7 (28)
Oskaloosa	176	61.9 (127)	88	17.0 (128)	33	9.1 (117)	80	16.3 (89)	88	6.8 (79)	87	79.3 (123)	353	1.4 (59)
Ottumwa	1,305	72.5 (62)	577	27.4 (67)	370	20.0 (43)	620	18.7 (72)	577	7.1 (75)	522	79.9 (118)	2,235	2.9 (110)
Pella	62	74.2 (50)	25	28.0 (62)	21	14.3 (84)	18	22.2 (56)	25	4.0 (114)	98	83.7 (78)	169	1.8 (74)
Perry	197	81.7 (18)	74	21.6 (106)	107	21.5 (37)	69	10.1 (129)	74	2.7 (123)	106	76.4 (138)	620	1.8 (74)
Pocahontas	32	75.0 (43)	18	16.7 (130)	14	21.4 (38)	18	11.1 (124)	18	5.6 (91)	108	86.1 (52)	160	4.4 (127)
Pomeroy	2	100.0 (1)	1	0.0 (147)	1	0.0 (137)	3	0.0 (144)	1	0.0 (130)	4	100.0 (1)	9	0.0 (1)
Postville	130	80.8 (21)	64	40.6 (15)	37	27.0 (18)	62	24.2 (46)	63	19.0 (8)	344	87.5 (39)	690	4.6 (132)
Red Oak	137	74.5 (48)	50	18.0 (123)	36	19.4 (50)	53	9.4 (134)	50	6.0 (89)	158	86.1 (52)	350	0.9 (39)
Redfield	22	63.6 (120)	9	0.0 (147)	6	0.0 (137)	8	0.0 (144)	9	0.0 (130)	2	100.0 (1)	27	0.0 (1)
Refugee Cln	34	100.0 (1)	16	68.8 (3)	16	56.3 (2)	20	5.0 (143)	16	0.0 (130)	82	65.9 (152)	152	0.0 (1)
Rock Rapid	66	71.2 (74)	34	41.2 (14)	30	13.3 (96)	39	20.5 (63)	34	8.8 (59)	136	81.6 (102)	241	0.8 (34)
Rock Valle	124	83.9 (11)	55	29.1 (58)	44	22.7 (31)	53	30.2 (24)	55	16.4 (17)	145	82.1 (97)	299	1.0 (41)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Rockwell C	49	61.2 (133)	23	8.7 (144)	22	9.1 (117)	31	25.8 (38)	23	4.3 (110)	153	87.6 (38)	251	4.4 (127)
SDM	1,327	74.0 (55)	537	29.1 (58)	563	18.5 (55)	530	10.9 (126)	537	2.2 (127)	173	82.1 (97)	1,913	1.8 (74)
Sac City	32	65.6 (109)	13	30.8 (46)	12	25.0 (21)	18	22.2 (56)	13	15.4 (22)	116	92.2 (14)	184	6.5 (142)
Sheldon	142	81.7 (18)	68	25.0 (81)	44	22.7 (31)	64	14.1 (109)	68	4.4 (108)	278	89.9 (21)	619	15.7 (149)
Shenandoah	96	57.3 (140)	43	23.3 (92)	24	0.0 (137)	58	17.2 (84)	43	7.0 (77)	154	83.8 (77)	352	2.8 (107)
Sibley	48	79.2 (31)	25	28.0 (62)	21	28.6 (14)	20	10.0 (130)	25	16.0 (19)	123	87.0 (46)	242	16.5 (152)
Sidney	34	67.6 (91)	11	9.1 (143)	13	38.5 (5)	10	30.0 (25)	11	9.1 (54)	46	80.4 (112)	88	1.1 (43)
Sigourney	45	60.0 (135)	21	14.3 (138)	12	25.0 (21)	17	11.8 (120)	21	4.8 (100)	34	76.5 (136)	94	5.3 (135)
Sioux Cent	197	82.2 (16)	73	28.8 (61)	88	23.9 (27)	70	21.4 (60)	72	5.6 (91)	262	85.1 (64)	537	0.0 (1)
Sioux City	2,328	68.3 (87)	907	22.3 (99)	653	14.1 (88)	999	11.4 (123)	905	4.3 (110)	3,511	79.8 (120)	7,542	1.1 (43)
Spencer	236	74.6 (46)	113	37.2 (22)	78	21.8 (36)	92	29.3 (28)	113	12.4 (36)	478	87.4 (42)	1,054	16.9 (154)
Spirit Lak	118	79.7 (30)	48	29.2 (56)	33	9.1 (117)	52	28.8 (30)	47	14.9 (25)	193	92.2 (14)	465	14.8 (148)
Storm Lake	516	70.0 (81)	239	19.2 (118)	151	13.2 (101)	216	15.3 (94)	238	2.9 (122)	1,055	84.7 (66)	2,282	5.6 (138)
Stuart	28	64.3 (114)	13	30.8 (46)	7	0.0 (137)	11	18.2 (76)	13	15.4 (22)	38	84.2 (70)	93	3.2 (115)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Tama	137	70.8 (78)	74	23.0 (97)	74	13.5 (94)	65	23.1 (49)	74	1.4 (129)	252	80.2 (115)	525	2.5 (96)
Tama/Settl	8	50.0 (151)	10	50.0 (4)	6	16.7 (69)	8	37.5 (11)	10	10.0 (46)	22	81.8 (101)	46	4.3 (123)
Tipton	20	55.0 (143)	16	31.3 (45)	6	16.7 (69)	17	17.6 (80)	16	18.8 (9)	33	84.8 (65)	58	5.2 (134)
Traer	14	71.4 (70)	10	40.0 (16)	10	20.0 (43)	5	20.0 (65)	10	20.0 (6)	32	90.6 (18)	73	1.4 (59)
UIHC	75	86.7 (7)	16	25.0 (81)	5	20.0 (43)	17	17.6 (80)	15	6.7 (80)	13	84.6 (67)	62	1.6 (68)
Vinton	77	71.4 (70)	35	34.3 (34)	25	12.0 (105)	43	25.6 (42)	35	14.3 (26)	127	70.9 (148)	353	2.8 (107)
WDM	248	78.2 (34)	110	33.6 (36)	127	22.8 (30)	116	14.7 (100)	110	6.4 (85)	38	86.8 (48)	597	0.3 (20)
Wapello	1	0.0 (153)	0	0.0 (147)	0	0.0 (137)	0	0.0 (144)	0	0.0 (130)	0	0.0 (154)	1	0.0 (1)
Washington	126	77.8 (38)	53	41.5 (13)	56	16.1 (75)	44	22.7 (55)	53	13.2 (33)	120	83.3 (82)	210	4.3 (123)
Waterloo	2,402	72.0 (66)	961	28.9 (60)	735	17.1 (68)	983	19.7 (68)	958	8.6 (61)	2,609	75.9 (139)	7,521	2.8 (107)
Waukee	145	80.0 (25)	80	25.0 (81)	103	19.4 (50)	80	15.0 (95)	80	5.0 (96)	27	85.2 (62)	439	0.5 (21)
Waukon	85	65.9 (107)	43	23.3 (92)	18	11.1 (109)	43	23.3 (47)	43	4.7 (102)	168	82.1 (97)	381	12.6 (145)
Waverly	68	64.7 (112)	43	34.9 (33)	46	19.6 (48)	37	35.1 (13)	43	18.6 (10)	152	81.6 (102)	339	1.2 (50)
Wbstr Ct	216	74.1 (54)	111	26.1 (74)	65	15.4 (81)	98	13.3 (114)	110	4.5 (106)	377	82.8 (88)	796	4.4 (127)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2018 - 2020 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
West Union	61	77.0 (40)	30	16.7 (130)	9	11.1 (109)	34	14.7 (100)	28	7.1 (75)	105	81.0 (109)	251	18.3 (155)
Williamsburg	1	100.0 (1)	2	50.0 (4)	4	25.0 (21)	1	0.0 (144)	2	0.0 (130)	9	77.8 (130)	12	0.0 (1)
Winterset	90	63.3 (122)	50	36.0 (27)	33	18.2 (58)	43	32.6 (17)	49	14.3 (26)	108	82.4 (94)	286	1.4 (59)
Iowa	**40,556**	**71.0**	**17,363**	**26.5**	**13,983**	**17.4**	**17,532**	**16.9**	**17,328**	**6.5**	**46,513**	**82.7**	**123,520**	**2.7**

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.