WIC PNSS & PedNSS

Table 4A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Ackley	66.7	0.0	33.3	0.0	0.0	0.0	0.0
Adel	84.1	2.3	5.7	0.0	0.0	8.0	0.0
Albia	90.7	1.9	4.7	0.0	0.0	2.8	0.0
Algona	73.3	1.2	21.1	0.8	0.4	3.2	0.0
Allen WHC	42.9	31.5	9.4	0.0	3.7	12.4	0.0
Allison	88.5	1.5	4.6	0.0	0.0	5.3	0.0
Altoona	65.9	18.4	5.1	0.0	2.8	7.8	0.0
Ames	61.3	9.6	15.2	0.3	5.6	8.0	0.0
Anamosa	85.2	3.7	4.4	0.0	0.7	5.9	0.0
Ankeny	66.4	14.9	9.8	0.3	1.8	6.8	0.0
Atlantic	81.7	0.4	12.9	0.0	0.4	4.5	0.0
Audubon	87.1	2.4	5.9	0.0	2.4	2.4	0.0
Avoca	82.1	2.6	5.1	0.0	0.0	10.3	0.0
Bedford	93.9	0.0	4.1	0.0	0.0	2.0	0.0
Belle Plaine	82.9	1.4	8.6	0.0	0.0	7.1	0.0
Bloomfield	91.3	2.2	6.5	0.0	0.0	0.0	0.0
Boone	84.6	4.3	6.4	0.0	0.0	4.8	0.0
Brdlwns RP	31.0	26.7	22.3	0.2	13.0	6.8	0.0
Brooklyn	75.0	0.0	25.0	0.0	0.0	0.0	0.0
Burlington	63.1	19.6	5.8	0.0	0.4	11.2	0.0
CHC	41.3	24.8	17.6	0.1	1.3	14.9	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
CR Urban	38.6	39.5	7.7	0.2	0.9	13.1	0.0
Carlisle	72.4	6.9	8.0	1.1	1.1	10.3	0.0
Carroll	78.9	5.3	12.7	0.2	0.2	2.6	0.0
Cedar Falls	69.6	8.9	9.8	0.0	3.6	8.0	0.0
Centervill	88.4	2.0	8.2	0.0	0.0	1.4	0.0
Chariton	90.6	0.6	6.9	0.0	0.0	1.9	0.0
Charles City	70.9	10.3	10.9	0.3	3.8	3.8	0.0
Cherokee	76.8	1.9	16.6	1.4	0.9	2.4	0.0
Clarinda	93.3	1.3	4.7	0.0	0.0	0.7	0.0
Clarion	30.6	0.0	68.2	0.0	0.0	1.2	0.0
Clinton	65.0	9.2	14.1	0.0	0.2	11.4	0.0
Colfax	81.3	1.6	9.4	0.0	1.6	4.7	1.6
Columbus J	13.0	0.0	45.5	1.3	39.0	1.3	0.0
Corning	96.8	1.6	1.6	0.0	0.0	0.0	0.0
Corydon	93.3	0.0	3.3	0.0	3.3	0.0	0.0
Council Blfs	66.0	3.1	21.3	0.4	0.3	8.8	0.0
Cresco	89.2	0.6	9.5	0.0	0.0	0.6	0.0
Creston	84.5	1.1	10.5	0.3	0.3	3.3	0.0
Dallas Ctr	93.3	0.0	6.7	0.0	0.0	0.0	0.0
Decorah	77.2	0.6	21.5	0.0	0.6	0.0	0.0
Denison	26.8	5.2	61.7	0.5	4.6	1.2	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Dewitt	86.0	0.0	6.0	0.0	0.0	8.0	0.0
Dubuque	64.9	16.8	8.2	0.2	0.9	9.0	0.0
Dyersville	68.6	2.9	25.7	0.0	0.0	2.9	0.0
Eagle Grove	32.9	0.0	64.5	0.0	0.0	2.6	0.0
Edgerton	45.5	23.3	13.4	1.0	1.3	15.3	0.1
Eldora	82.8	0.0	10.3	0.0	0.0	6.9	0.0
Elkader	87.9	0.5	7.1	0.0	0.0	4.4	0.0
Emmetsburg	82.3	4.4	11.5	0.0	0.0	1.8	0.0
Esthervill	57.1	3.3	37.6	0.0	0.0	2.0	0.0
Fairfiel	82.8	3.1	6.3	0.0	1.6	6.3	0.0
Forest City	74.7	1.4	16.3	0.0	0.0	7.7	0.0
Fort Dodge	60.8	7.6	16.6	0.2	0.2	14.2	0.3
Ft Madison	85.2	3.9	6.9	0.0	0.7	3.3	0.0
Garner	65.7	4.4	25.5	0.0	0.0	4.4	0.0
Glenwood	87.6	0.0	10.6	0.0	0.0	1.8	0.0
Grand Junction	76.5	0.0	11.8	0.0	0.0	11.8	0.0
Greenfield	91.2	0.0	5.3	0.0	0.9	2.6	0.0
Grinnell	84.6	3.8	9.6	0.0	0.6	1.3	0.0
Grundy Ctr	94.1	2.4	0.0	0.0	0.0	3.5	0.0
Guthrie Ct	85.2	3.7	9.3	0.0	0.0	1.9	0.0
Hampton	37.1	0.4	60.6	0.4	0.0	1.5	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Harlan	78.2	1.6	15.4	0.0	0.0	4.8	0.0
Hawarden	48.8	1.6	45.7	0.8	0.0	3.1	0.0
Hispanic Cln	2.0	0.0	98.0	0.0	0.0	0.0	0.0
Humboldt	74.1	1.4	24.5	0.0	0.0	0.0	0.0
ILH	31.9	16.3	40.0	0.1	5.6	6.0	0.0
Ida Grove	81.0	0.8	16.7	0.0	0.0	1.6	0.0
Independen	91.6	1.3	3.6	0.0	0.0	3.6	0.0
Indianola	84.5	3.6	6.4	0.0	0.0	5.5	0.0
Iowa City	28.0	46.9	17.9	0.3	1.3	5.6	0.1
Iowa Falls	74.7	4.0	16.2	0.0	0.0	5.1	0.0
Jamaica	42.9	2.9	51.4	0.0	0.0	2.9	0.0
Jefferson	82.1	1.7	12.8	0.9	0.9	1.7	0.0
KR	46.3	31.3	4.8	0.0	2.0	15.6	0.0
Keokuk	80.0	8.3	3.3	0.0	0.0	8.3	0.0
Keosauqua	96.2	0.0	1.9	0.0	0.0	1.9	0.0
Knoxville	89.8	3.1	3.3	0.4	0.4	2.9	0.0
LCPH	50.0	20.7	8.6	0.0	1.7	19.0	0.0
Lamoni	80.4	0.0	6.5	0.0	2.2	10.9	0.0
Le Mars	60.3	4.0	31.0	0.7	0.3	3.7	0.0
Lenox	43.3	0.0	55.2	0.0	0.0	1.5	0.0
Leon	89.0	0.8	8.5	0.0	0.0	1.7	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Logan	82.6	2.4	12.0	0.0	0.0	3.0	0.0
Manchester	85.3	4.3	4.3	0.0	0.0	6.0	0.0
Manning	87.5	0.0	0.0	0.0	0.0	12.5	0.0
Maquoketa	78.2	6.6	4.6	0.5	0.0	10.2	0.0
Marengo	76.2	0.0	15.9	0.0	0.0	7.9	0.0
Marion	70.9	7.1	9.5	0.0	0.8	11.6	0.1
Marshalltown	37.2	3.6	48.5	0.6	8.5	1.7	0.0
Mason City	67.7	6.0	15.1	0.5	0.8	9.9	0.0
Monticello	88.9	4.2	5.6	0.0	0.0	1.4	0.0
Mt Ayr	89.7	0.0	10.3	0.0	0.0	0.0	0.0
Mt Pleasan	78.1	3.8	14.6	0.0	0.3	3.2	0.0
Mtown Dtown	0.0	0.0	57.1	0.0	28.6	14.3	0.0
Muscatine	48.4	8.7	39.6	0.0	1.9	1.5	0.0
Nevada	48.6	0.0	40.5	0.0	0.0	10.8	0.0
New Hampto	74.6	0.0	19.8	0.0	0.0	5.6	0.0
Newton	82.9	5.8	6.7	0.2	1.2	3.2	0.0
North Towne	52.6	18.9	12.1	0.1	4.1	12.1	0.1
Northwood	80.6	3.0	13.4	0.0	1.5	1.5	0.0
Norwalk	82.5	1.3	10.0	0.0	0.0	6.3	0.0
Oakridge	1.4	82.6	4.3	1.4	0.0	10.1	0.0
Oelwein	86.4	0.0	9.2	0.0	0.0	4.3	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Onawa	85.8	0.7	8.5	2.8	0.0	2.1	0.0
Orange Cty	52.5	0.0	46.4	0.0	0.0	1.1	0.0
Osage	88.3	0.6	8.8	0.0	0.0	2.3	0.0
Osceola	58.2	0.4	37.9	0.7	1.1	1.8	0.0
Oskaloosa	80.4	7.2	7.5	0.0	0.0	4.9	0.0
Ottumwa	67.4	8.5	16.8	0.0	1.7	5.8	0.0
Pella	89.2	0.0	7.8	0.0	0.0	2.9	0.0
Perry	38.5	7.4	50.8	0.0	1.9	1.3	0.0
Pocahontas	77.2	12.3	7.0	0.0	0.0	3.5	0.0
Pomeroy	100.0	0.0	0.0	0.0	0.0	0.0	0.0
Postville	32.1	10.4	56.3	0.0	0.0	1.3	0.0
Red Oak	85.8	0.0	11.3	0.4	0.0	2.5	0.0
Redfield	88.9	0.0	0.0	0.0	0.0	11.1	0.0
Refugee Cln	0.0	98.6	1.4	0.0	0.0	0.0	0.0
Rock Rapid	74.8	0.0	14.4	1.8	0.0	9.0	0.0
Rock Valle	40.4	0.0	58.6	0.5	0.0	0.5	0.0
Rockwell C	89.0	0.0	8.3	0.0	0.0	2.8	0.0
SDM	38.8	15.2	31.4	0.2	9.0	5.5	0.0
Sac City	91.8	0.0	8.2	0.0	0.0	0.0	0.0
Sheldon	65.0	3.3	28.5	0.0	0.0	3.3	0.0
Shenandoah	84.4	0.6	13.3	0.0	0.0	1.7	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Sibley	47.7	2.3	43.0	0.0	1.2	5.8	0.0
Sidney	93.5	0.0	6.5	0.0	0.0	0.0	0.0
Sigourney	97.2	0.0	0.0	0.0	0.0	2.8	0.0
Sioux Cent	29.3	0.3	67.0	0.0	0.9	2.5	0.0
Sioux City	31.1	13.0	41.5	3.4	1.5	9.4	0.1
Spencer	78.0	2.0	16.3	0.0	0.9	2.9	0.0
Spirit Lak	83.2	1.8	12.6	0.0	0.0	2.4	0.0
Storm Lake	15.5	5.4	56.9	0.0	11.7	10.4	0.0
Stuart	67.9	14.3	14.3	1.8	0.0	1.8	0.0
Tama	42.9	0.0	42.5	11.6	0.0	3.0	0.0
Tama/Settl	6.7	0.0	13.3	66.7	0.0	13.3	0.0
Tipton	87.3	0.0	7.3	0.0	0.0	5.5	0.0
Traer	90.3	0.0	6.5	0.0	0.0	3.2	0.0
UIHC	64.5	16.1	14.5	0.0	1.6	3.2	0.0
Vinton	89.2	0.0	6.3	0.0	0.0	4.5	0.0
WDM	41.5	23.4	26.5	0.5	3.5	4.7	0.0
Washington	64.2	1.4	31.1	0.0	0.0	3.3	0.0
Waterloo	42.1	26.3	12.3	0.0	4.5	14.7	0.1
Waukee	67.1	9.5	15.5	0.4	2.0	5.6	0.0
Waukon	85.0	0.0	11.5	0.0	0.0	3.5	0.0
Waverly	85.9	3.0	3.0	0.0	1.0	7.0	0.0

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 4A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Racial and Ethnic Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wbstr Ct	52.2	1.6	37.9	0.3	3.8	4.1	0.0
West Union	86.9	0.0	9.5	0.0	0.0	3.6	0.0
Williamsburg	30.0	0.0	70.0	0.0	0.0	0.0	0.0
Winterset	85.9	1.3	8.7	0.0	0.7	3.4	0.0
Iowa	**52.8**	**14.6**	**21.4**	**0.5**	**3.2**	**7.5**	**0.0**
National (Prior Year)	**47.2**	**18.8**	**22.9**	**1.3**	**2.7**	**7.1**	**0.1**

(1) Reporting period is January 1 through December 31
(2) Analyses based on one record per child
(3) Excludes records with unknown data errors.
(4) Prenatal program participation
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions

WIC PNSS & PedNSS

Table 5A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Ackley	38.1	28.6	33.3
Adel	36.4	12.5	51.1
Albia	36.4	17.8	45.8
Algona	39.4	14.7	45.8
Allen WHC	48.4	19.1	32.5
Allison	38.9	16.8	44.3
Altoona	37.3	17.5	45.2
Ames	35.1	22.4	42.5
Anamosa	40.7	17.0	42.2
Ankeny	37.5	20.2	42.3
Atlantic	40.2	17.4	42.4
Audubon	31.8	24.7	43.5
Avoca	33.3	23.1	43.6
Bedford	32.7	14.3	53.1
Belle Plaine	31.4	18.6	50.0
Bloomfield	39.1	13.0	47.8
Boone	28.7	27.1	44.1
Brdlwns RP	36.4	20.5	43.0
Brooklyn	27.8	27.8	44.4
Burlington	47.7	18.3	34.1
CHC	40.6	18.6	40.8

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
CR Urban	41.0	17.6	41.4
Carlisle	33.3	16.1	50.6
Carroll	38.4	15.8	45.8
Cedar Falls	33.0	22.3	44.6
Centervill	45.6	19.7	34.7
Chariton	38.1	18.1	43.8
Charles City	40.6	15.0	44.4
Cherokee	37.4	18.0	44.5
Clarinda	40.9	14.8	44.3
Clarion	38.0	19.4	42.6
Clinton	46.4	15.8	37.8
Colfax	32.8	20.3	46.9
Columbus J	39.0	18.2	42.9
Corning	41.9	21.0	37.1
Corydon	37.8	16.7	45.6
Council Blfs	38.7	20.2	41.1
Cresco	39.2	20.3	40.5
Creston	38.1	19.1	42.8
Dallas Ctr	40.0	33.3	26.7
Decorah	36.7	19.6	43.7
Denison	38.4	18.5	43.0

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Dewitt	44.0	18.0	38.0
Dubuque	42.6	16.9	40.5
Dyersville	25.7	14.3	60.0
Eagle Grove	37.4	11.6	51.0
Edgerton	43.4	18.6	38.0
Eldora	27.6	25.9	46.6
Elkader	38.5	12.6	48.9
Emmetsburg	35.4	21.2	43.4
Esthervill	35.1	22.0	42.9
Fairfiel	35.9	15.6	48.4
Forest City	38.9	19.9	41.2
Fort Dodge	43.8	17.0	39.2
Ft Madison	39.8	20.7	39.5
Garner	41.6	16.1	42.3
Glenwood	42.5	17.7	39.8
Grand Junction	35.3	17.6	47.1
Greenfield	37.7	19.3	43.0
Grinnell	37.2	21.2	41.7
Grundy Ctr	34.1	18.8	47.1
Guthrie Ct	31.5	19.4	49.1
Hampton	37.5	17.0	45.6

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Harlan	41.5	14.9	43.6
Hawarden	35.4	14.2	50.4
Hispanic Cln	22.4	28.6	49.0
Humboldt	33.8	18.7	47.5
ILH	34.2	22.7	43.2
Ida Grove	36.5	16.7	46.8
Independen	35.6	17.8	46.7
Indianola	35.4	19.6	45.0
Iowa City	38.3	19.2	42.5
Iowa Falls	34.3	18.2	47.5
Jamaica	17.1	28.6	54.3
Jefferson	34.2	19.7	46.2
KR	29.9	22.4	47.6
Keokuk	46.0	16.0	38.0
Keosauqua	36.5	15.4	48.1
Knoxville	41.6	18.1	40.3
LCPH	58.6	13.8	27.6
Lamoni	30.4	19.6	50.0
Le Mars	32.0	21.3	46.7
Lenox	29.9	17.9	52.2
Leon	40.7	18.6	40.7

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Logan	43.1	15.0	41.9
Manchester	32.8	19.0	48.3
Manning	37.5	0.0	62.5
Maquoketa	38.1	21.3	40.6
Marengo	41.3	17.5	41.3
Marion	37.1	18.8	44.1
Marshalltown	36.1	21.4	42.5
Mason City	41.8	17.0	41.2
Monticello	36.1	13.9	50.0
Mt Ayr	39.7	13.2	47.1
Mt Pleasan	37.8	19.0	43.2
Mtown Dtown	14.3	57.1	28.6
Muscatine	41.3	20.9	37.8
Nevada	40.5	13.5	45.9
New Hampto	31.7	19.8	48.4
Newton	36.6	20.6	42.8
North Towne	36.7	18.4	44.8
Northwood	41.8	20.9	37.3
Norwalk	40.0	12.5	47.5
Oakridge	27.5	27.5	44.9
Oelwein	41.8	15.8	42.4

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Onawa	43.3	19.1	37.6
Orange Cty	39.2	15.5	45.3
Osage	43.3	15.8	40.9
Osceola	42.6	18.4	39.0
Oskaloosa	38.5	12.8	48.7
Ottumwa	42.5	19.8	37.7
Pella	32.4	15.7	52.0
Perry	32.7	19.4	47.9
Pocahontas	28.1	21.1	50.9
Pomeroy	0.0	75.0	25.0
Postville	32.5	16.7	50.8
Red Oak	33.5	15.9	50.6
Redfield	55.6	7.4	37.0
Refugee Cln	27.1	18.6	54.3
Rock Rapid	26.1	17.1	56.8
Rock Valle	38.9	15.2	46.0
Rockwell C	29.4	14.7	56.0
SDM	40.2	20.2	39.7
Sac City	31.5	12.3	56.2
Sheldon	35.5	22.4	42.1
Shenandoah	39.3	16.2	44.5

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Sibley	32.6	17.4	50.0
Sidney	27.4	16.1	56.5
Sigourney	37.5	19.4	43.1
Sioux Cent	35.8	20.4	43.8
Sioux City	41.0	17.5	41.5
Spencer	38.0	20.9	41.1
Spirit Lak	44.3	16.8	38.9
Storm Lake	36.5	19.6	43.9
Stuart	42.9	14.3	42.9
Tama	33.5	22.7	43.8
Tama/Settl	33.3	20.0	46.7
Tipton	29.1	12.7	58.2
Traer	29.0	29.0	41.9
UIHC	75.8	8.1	16.1
Vinton	28.4	19.9	51.7
WDM	31.1	24.4	44.5
Washington	31.6	23.6	44.8
Waterloo	39.2	19.3	41.6
Waukee	32.1	29.4	38.5
Waukon	41.6	15.9	42.5
Waverly	26.6	24.1	49.2

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 5A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Age Distribution by Clinic

Children Aged < 5 Years (2)

Clinic	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Wbstr Ct	38.9	19.7	41.4
West Union	48.8	11.9	39.3
Williamsburg	20.0	30.0	50.0
Winterset	36.2	26.8	36.9
Iowa	**38.8**	**19.1**	**42.1**
National (Prior Year)	**33.4**	**21.3**	**45.3**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Records rejected if date of birth and date of visit unknown.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ackley		0.0 (1)	33.3 (150)		0.0 (1)	4.8 (138)	14.3 (80)		25.0 (136)	0.0 (1)		12.5 (109)
Adel		0.0 (1)	10.0 (106)		4.7 (20)	3.5 (120)	9.4 (14)		26.2 (137)	11.9 (34)		2.6 (18)
Albia		15.8 (139)	0.0 (1)		2.0 (5)	4.0 (129)	9.1 (12)		27.3 (138)	15.9 (84)		19.0 (143)
Algona		10.7 (107)	16.1 (136)		6.9 (57)	0.8 (34)	14.7 (85)		23.1 (130)	21.2 (129)		5.0 (33)
Allen WHC		6.3 (55)	4.5 (41)		5.2 (28)	3.5 (120)	12.3 (43)		14.0 (36)	19.0 (118)		10.3 (93)
Allison		25.0 (147)	12.5 (121)		10.4 (129)	1.6 (63)	11.2 (25)		17.9 (84)	14.3 (58)		7.0 (52)
Altoona		6.4 (60)	2.1 (32)		5.8 (35)	1.9 (76)	12.6 (49)		22.7 (128)	12.5 (39)		9.1 (74)
Ames		9.7 (96)	9.7 (103)		8.1 (85)	3.9 (128)	11.7 (31)		12.7 (25)	11.6 (32)		11.8 (102)
Anamosa		3.7 (37)	3.7 (37)		3.1 (14)	3.1 (113)	14.8 (86)		16.4 (68)	23.6 (141)		18.3 (140)
Ankeny		10.7 (107)	6.7 (68)		7.0 (59)	1.8 (70)	12.2 (42)		14.0 (36)	13.3 (45)		3.3 (23)
Atlantic		9.5 (93)	14.3 (129)		6.4 (48)	0.5 (22)	13.7 (72)		20.2 (107)	12.4 (37)		8.8 (71)
Audubon		0.0 (1)	0.0 (1)		4.8 (21)	0.0 (1)	13.3 (62)		14.3 (41)	7.1 (12)		1.9 (14)
Avoca		0.0 (1)	0.0 (1)		5.7 (34)	5.7 (145)	11.4 (28)		46.7 (150)	13.3 (45)		13.0 (115)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Bedford		9.1 (87)	9.1 (96)		6.4 (48)	0.0 (1)	14.9 (89)		16.7 (69)	20.8 (125)		18.2 (139)
Belle Plaine		10.0 (97)	30.0 (149)		4.8 (21)	0.0 (1)	16.1 (108)		17.2 (78)	20.7 (124)		8.8 (71)
Bloomfield		0.0 (1)	0.0 (1)		4.8 (21)	4.8 (138)	19.0 (136)		10.0 (12)	20.0 (121)		0.0 (1)
Boone		7.1 (65)	0.0 (1)		5.1 (27)	5.1 (143)	9.0 (11)		11.5 (20)	7.7 (15)		4.1 (28)
Brdlwns RP		10.7 (107)	6.4 (65)		7.9 (80)	1.4 (53)	13.4 (68)		16.9 (72)	14.6 (62)		6.0 (41)
Brooklyn		0.0 (1)	0.0 (1)		5.9 (36)	2.9 (106)	11.8 (34)		35.7 (146)	7.1 (12)		13.3 (119)
Burlington		12.0 (120)	7.7 (80)		5.9 (36)	3.0 (109)	9.9 (15)		14.1 (39)	10.8 (27)		10.8 (98)
CHC		14.9 (137)	8.4 (87)		7.7 (75)	1.7 (66)	10.6 (21)		13.4 (30)	15.7 (80)		15.4 (130)
CR Urban		10.8 (112)	7.5 (77)		7.6 (71)	1.6 (63)	13.0 (53)		17.0 (76)	13.6 (49)		18.3 (140)
Carlisle		5.3 (45)	10.5 (111)		7.1 (61)	2.4 (91)	19.0 (136)		22.5 (126)	22.5 (138)		7.7 (57)
Carroll		10.1 (101)	6.1 (59)		5.9 (36)	0.7 (29)	10.1 (18)		18.3 (90)	12.8 (40)		1.8 (13)
Cedar Falls		5.3 (45)	5.3 (48)		2.0 (5)	5.0 (141)	7.9 (8)		9.5 (10)	4.8 (8)		13.1 (116)
Centervill		7.3 (66)	4.9 (44)		9.8 (119)	1.4 (53)	16.8 (120)		13.7 (31)	17.6 (109)		10.6 (96)
Chariton		8.8 (82)	14.7 (132)		5.2 (28)	5.8 (146)	7.8 (7)		18.3 (90)	8.5 (18)		12.0 (104)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Charles City		2.9 (33)	11.4 (116)		4.9 (25)	1.9 (76)	11.7 (31)		16.7 (69)	14.7 (63)		4.6 (31)
Cherokee		12.5 (125)	0.0 (1)		7.3 (65)	0.5 (22)	16.5 (112)		22.2 (122)	16.7 (93)		12.1 (105)
Clarinda		0.0 (1)	8.6 (89)		6.9 (57)	1.4 (53)	11.8 (34)		19.7 (105)	11.5 (31)		8.8 (71)
Clarion		7.0 (64)	15.8 (134)		11.6 (136)	2.6 (95)	15.9 (102)		22.4 (125)	16.3 (89)		2.6 (18)
Clinton		10.1 (101)	5.1 (47)		9.2 (108)	0.7 (29)	13.4 (68)		21.4 (116)	14.7 (63)		9.4 (80)
Colfax		0.0 (1)	25.0 (145)		9.4 (114)	0.0 (1)	12.5 (47)		19.4 (103)	16.1 (88)		8.3 (63)
Columbus J		9.5 (93)	4.8 (43)		2.7 (8)	1.4 (53)	19.2 (139)		12.1 (22)	21.2 (129)		7.9 (58)
Corning		0.0 (1)	12.5 (121)		8.6 (98)	0.0 (1)	19.0 (136)		30.4 (145)	13.0 (41)		19.4 (144)
Corydon		13.6 (134)	22.7 (144)		7.0 (59)	5.8 (146)	15.1 (92)		10.8 (13)	29.7 (148)		18.0 (138)
Council Blfs		10.7 (107)	8.2 (84)		9.2 (108)	2.1 (83)	13.9 (75)		18.7 (94)	15.3 (75)		13.6 (122)
Cresco		8.3 (77)	11.1 (114)		5.2 (28)	1.3 (45)	14.4 (83)		15.9 (61)	14.3 (58)		9.5 (83)
Creston		9.2 (90)	5.7 (53)		10.0 (124)	2.1 (83)	17.4 (125)		14.2 (40)	22.0 (134)		9.7 (84)
Dallas Ctr		0.0 (1)	0.0 (1)		14.3 (143)	0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)		0.0 (1)
Decorah		14.7 (136)	17.6 (140)		9.9 (120)	0.7 (29)	16.4 (110)		14.9 (49)	20.9 (126)		4.4 (30)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Denison		8.2 (75)	5.5 (51)		7.5 (68)	1.1 (40)	18.1 (133)		18.1 (87)	18.5 (115)		14.2 (125)
Dewitt		50.0 (149)	0.0 (1)		8.3 (91)	4.2 (133)	16.7 (117)		30.0 (142)	15.0 (71)		10.0 (86)
Dubuque		8.1 (74)	5.8 (55)		6.3 (43)	3.3 (116)	7.6 (6)		16.9 (72)	8.1 (17)		15.6 (132)
Dyersville		0.0 (1)	16.7 (137)		3.0 (12)	0.0 (1)	9.1 (12)		5.6 (5)	16.7 (93)		10.0 (86)
Eagle Grove		12.1 (124)	9.1 (96)		7.5 (68)	4.8 (138)	10.9 (24)		13.3 (27)	16.0 (85)		4.0 (27)
Edgerton		10.8 (112)	9.5 (100)		8.3 (91)	2.1 (83)	11.7 (31)		18.0 (86)	13.9 (55)		12.7 (113)
Eldora		22.2 (145)	0.0 (1)		12.3 (140)	3.5 (120)	21.1 (146)		14.8 (45)	25.9 (146)		13.3 (119)
Elkader		12.9 (131)	6.5 (66)		10.2 (127)	1.1 (40)	10.7 (22)		15.2 (52)	6.3 (11)		4.3 (29)
Emmetsburg		15.0 (138)	0.0 (1)		8.3 (91)	1.9 (76)	14.8 (86)		11.4 (19)	13.6 (49)		11.0 (99)
Esthervill		4.7 (42)	9.3 (99)		7.6 (71)	1.7 (66)	11.8 (34)		17.9 (84)	13.7 (53)		10.3 (93)
Fairfiel		4.5 (41)	13.6 (126)		6.6 (50)	4.1 (131)	16.5 (112)		8.6 (9)	17.2 (102)		4.8 (32)
Forest City		4.2 (38)	12.5 (121)		8.5 (97)	1.4 (53)	14.2 (79)		21.8 (119)	10.3 (23)		5.4 (36)
Fort Dodge		9.2 (90)	8.8 (92)		9.3 (112)	1.8 (70)	16.0 (104)		21.1 (113)	18.5 (115)		3.6 (25)
Ft Madison		10.1 (101)	7.2 (74)		7.7 (75)	1.7 (66)	16.0 (104)		15.4 (55)	17.9 (111)		6.7 (51)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Garner		0.0 (1)	20.8 (142)		4.6 (18)	3.8 (127)	13.1 (56)		14.0 (36)	15.8 (82)		6.3 (45)
Glenwood		12.0 (120)	0.0 (1)		6.3 (43)	4.5 (137)	18.0 (132)		15.2 (52)	17.4 (107)		8.3 (63)
Grand Junction		20.0 (143)	0.0 (1)		29.4 (150)	0.0 (1)	17.6 (130)		0.0 (1)	25.0 (143)		0.0 (1)
Greenfield		10.0 (97)	10.0 (106)		10.2 (127)	2.8 (103)	19.4 (140)		15.9 (61)	13.6 (49)		18.6 (142)
Grinnell		0.0 (1)	6.7 (68)		9.9 (120)	2.6 (95)	13.2 (60)		24.2 (135)	10.6 (24)		20.4 (147)
Grundy Ctr		10.0 (97)	0.0 (1)		7.9 (80)	1.3 (45)	10.5 (20)		21.2 (114)	12.1 (35)		7.9 (58)
Guthrie Ct		5.9 (51)	11.8 (119)		3.0 (12)	1.0 (37)	9.9 (15)		22.2 (122)	14.8 (65)		0.0 (1)
Hampton		3.6 (36)	12.5 (121)		4.8 (21)	1.2 (42)	15.3 (95)		21.6 (117)	18.0 (113)		7.2 (55)
Harlan		2.4 (32)	4.9 (44)		8.7 (102)	0.5 (22)	15.8 (100)		20.5 (109)	13.6 (49)		6.0 (41)
Hawarden		23.1 (146)	11.5 (117)		7.6 (71)	0.0 (1)	16.0 (104)		16.1 (67)	17.9 (111)		10.0 (86)
Hispanic Cln		0.0 (1)	22.2 (143)		8.3 (91)	0.0 (1)	20.8 (144)		28.0 (140)	20.0 (121)		3.4 (24)
Humboldt		13.0 (132)	4.3 (40)		9.5 (117)	0.7 (29)	16.8 (120)		21.0 (112)	22.6 (139)		1.0 (12)
ILH		10.6 (106)	6.6 (67)		8.1 (85)	1.3 (45)	14.5 (84)		19.3 (102)	16.9 (101)		10.4 (95)
Ida Grove		6.3 (55)	3.1 (35)		8.9 (104)	3.3 (116)	17.9 (131)		6.9 (6)	22.4 (137)		6.3 (45)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Independen		12.8 (130)	15.4 (133)		9.4 (114)	1.9 (76)	14.1 (77)		16.0 (64)	16.0 (85)		15.3 (129)
Indianola		14.3 (135)	8.6 (89)		12.0 (139)	3.4 (119)	12.0 (41)		22.0 (121)	13.4 (47)		13.5 (121)
Iowa City		12.0 (120)	7.9 (82)		6.7 (53)	4.4 (136)	11.9 (39)		15.0 (50)	12.4 (37)		10.6 (96)
Iowa Falls		12.5 (125)	6.3 (62)		9.5 (117)	2.1 (83)	12.6 (49)		13.3 (27)	15.6 (78)		14.6 (127)
Jamaica		0.0 (1)	40.0 (151)		0.0 (1)	3.0 (109)	21.2 (147)		12.5 (23)	37.5 (150)		8.7 (70)
Jefferson		13.3 (133)	13.3 (125)		9.2 (108)	1.8 (70)	16.5 (112)		14.8 (45)	16.7 (93)		5.6 (38)
KR		6.9 (62)	10.3 (109)		6.6 (50)	1.5 (59)	13.1 (56)		10.9 (15)	15.6 (78)		11.1 (100)
Keokuk		10.8 (112)	4.6 (42)		10.6 (132)	2.8 (103)	13.5 (70)		19.2 (101)	17.3 (105)		12.8 (114)
Keosauqua		0.0 (1)	27.3 (147)		2.0 (5)	4.1 (131)	24.5 (150)		16.0 (64)	36.0 (149)		3.1 (20)
Knoxville		10.1 (101)	8.4 (87)		11.9 (138)	4.3 (134)	12.3 (43)		14.4 (43)	16.7 (93)		16.2 (136)
LCPH		6.3 (55)	6.3 (62)		10.5 (130)	3.5 (120)	7.0 (4)		45.5 (149)	0.0 (1)		27.8 (149)
Lamoni		0.0 (1)	28.6 (148)		11.4 (134)	0.0 (1)	20.5 (143)		17.4 (80)	21.7 (132)		16.1 (135)
Le Mars		8.3 (77)	5.0 (46)		9.4 (114)	1.0 (37)	17.0 (122)		15.7 (59)	16.4 (90)		8.6 (67)
Lenox		0.0 (1)	14.3 (129)		13.1 (142)	1.6 (63)	16.4 (110)		14.3 (41)	10.7 (25)		15.4 (130)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Leon		4.2 (38)	0.0 (1)		10.9 (133)	1.8 (70)	15.5 (98)		17.4 (80)	13.0 (41)		19.7 (145)
Logan		5.6 (48)	11.1 (114)		8.0 (84)	1.2 (42)	16.7 (117)		15.9 (61)	17.4 (107)		12.4 (107)
Manchester		15.8 (139)	5.3 (48)		5.4 (32)	6.3 (148)	10.8 (23)		18.5 (93)	11.1 (29)		8.6 (67)
Manning		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)	12.5 (47)		20.0 (106)	20.0 (121)		0.0 (1)
Maquoketa		3.0 (34)	6.1 (59)		7.2 (64)	1.5 (59)	13.3 (62)		15.2 (52)	16.5 (91)		5.8 (40)
Marengo		0.0 (1)	14.3 (129)		8.3 (91)	5.0 (141)	13.3 (62)		14.8 (45)	14.8 (65)		3.2 (21)
Marion		7.8 (71)	8.8 (92)		7.9 (80)	0.7 (29)	13.3 (62)		18.8 (96)	14.2 (57)		7.3 (56)
Marshalltown		7.9 (72)	8.2 (84)		4.2 (16)	3.6 (125)	14.1 (77)		13.7 (31)	13.2 (44)		11.8 (102)
Mason City		9.4 (92)	6.8 (71)		6.7 (53)	1.7 (66)	11.9 (39)		18.2 (88)	14.9 (70)		6.2 (44)
Monticello		0.0 (1)	5.9 (56)		4.2 (16)	0.0 (1)	7.0 (4)		18.2 (88)	12.1 (35)		10.0 (86)
Mt Ayr		0.0 (1)	0.0 (1)		14.3 (143)	0.0 (1)	14.3 (80)		24.1 (133)	17.2 (102)		9.1 (74)
Mt Pleasan		4.7 (42)	7.8 (81)		8.6 (98)	1.3 (45)	16.6 (116)		15.1 (51)	19.0 (118)		11.1 (100)
Mtown Dtown		0.0 (1)	0.0 (1)		14.3 (143)	0.0 (1)	14.3 (80)		50.0 (151)	0.0 (1)		50.0 (151)
Muscatine		8.9 (83)	8.9 (94)		7.7 (75)	1.3 (45)	15.1 (92)		19.6 (104)	19.1 (120)		14.7 (128)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Nevada		16.7 (141)	16.7 (137)		2.9 (10)	8.6 (150)	11.4 (28)		38.5 (147)	7.7 (15)		25.0 (148)
New Hampto		11.1 (115)	5.6 (52)		8.2 (89)	1.8 (70)	13.6 (71)		16.0 (64)	16.0 (85)		13.6 (122)
Newton		5.7 (49)	11.5 (117)		8.4 (96)	2.9 (106)	15.8 (100)		18.7 (94)	14.8 (65)		12.4 (107)
North Towne		8.2 (75)	6.8 (71)		7.9 (80)	1.5 (59)	13.8 (74)		17.4 (80)	11.3 (30)		15.6 (132)
Northwood		0.0 (1)	18.2 (141)		6.3 (43)	3.2 (114)	7.9 (8)		13.0 (26)	0.0 (1)		9.3 (77)
Norwalk		9.1 (87)	9.1 (96)		11.8 (137)	5.3 (144)	11.8 (34)		13.9 (35)	16.7 (93)		20.0 (146)
Oakridge		0.0 (1)	0.0 (1)		1.5 (4)	2.9 (106)	5.9 (3)		22.6 (127)	9.7 (20)		9.3 (77)
Oelwein		4.9 (44)	9.8 (104)		6.3 (43)	0.6 (27)	16.5 (112)		16.9 (72)	22.1 (135)		1.9 (14)
Onawa		11.5 (118)	3.8 (38)		7.5 (68)	2.2 (87)	12.7 (51)		22.2 (122)	13.0 (41)		8.4 (65)
Orange Cty		6.8 (61)	13.6 (126)		8.7 (102)	3.5 (120)	16.3 (109)		9.5 (10)	9.5 (19)		10.0 (86)
Osage		8.3 (77)	2.8 (34)		8.1 (85)	1.9 (76)	13.1 (56)		15.4 (55)	15.4 (76)		12.5 (109)
Osceola		9.0 (86)	6.0 (58)		6.0 (40)	2.6 (95)	12.4 (46)		18.9 (98)	15.1 (73)		13.1 (116)
Oskaloosa		5.9 (51)	9.8 (104)		9.9 (120)	3.2 (114)	13.1 (56)		20.2 (107)	15.8 (82)		7.1 (54)
Ottumwa		8.9 (83)	7.6 (79)		8.1 (85)	2.4 (91)	13.3 (62)		17.0 (76)	15.0 (71)		6.1 (43)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Pella		25.0 (147)	16.7 (137)		9.0 (105)	2.0 (82)	16.0 (104)		13.7 (31)	15.7 (80)		17.2 (137)
Perry		10.5 (105)	7.0 (73)		8.2 (89)	1.0 (37)	18.6 (135)		23.2 (131)	21.1 (128)		3.8 (26)
Pocahontas		9.1 (87)	0.0 (1)		14.3 (143)	1.8 (70)	21.4 (148)		24.1 (133)	24.1 (142)		5.4 (36)
Pomeroy		0.0 (1)	0.0 (1)		25.0 (148)	0.0 (1)	50.0 (151)		0.0 (1)	100.0 (151)		0.0 (1)
Postville		7.5 (68)	2.5 (33)		9.0 (105)	1.3 (45)	15.0 (90)		13.3 (27)	16.8 (100)		6.5 (50)
Red Oak		5.7 (49)	8.6 (89)		9.3 (112)	3.0 (109)	12.3 (43)		11.0 (16)	7.6 (14)		12.5 (109)
Redfield		11.1 (115)	0.0 (1)		7.4 (66)	3.7 (126)	3.7 (2)		30.0 (142)	0.0 (1)		0.0 (1)
Refugee Cln		12.5 (125)	25.0 (145)		2.9 (10)	1.5 (59)	13.2 (60)		8.3 (7)	16.7 (93)		28.8 (150)
Rock Rapid		6.3 (55)	0.0 (1)		12.8 (141)	0.9 (35)	8.3 (10)		8.5 (8)	5.1 (9)		6.3 (45)
Rock Valle		7.3 (66)	7.3 (76)		6.8 (56)	0.5 (22)	17.4 (125)		15.5 (58)	10.7 (25)		8.5 (66)
Rockwell C		5.9 (51)	5.9 (56)		10.5 (130)	1.9 (76)	20.0 (141)		21.8 (119)	21.8 (133)		0.0 (1)
SDM		9.6 (95)	9.6 (101)		7.1 (61)	1.4 (53)	12.8 (52)		17.4 (80)	14.1 (56)		6.3 (45)
Sac City		8.3 (77)	0.0 (1)		8.6 (98)	0.0 (1)	17.1 (123)		21.6 (117)	21.6 (131)		2.4 (17)
Sheldon		12.5 (125)	6.3 (62)		7.8 (79)	2.4 (91)	15.6 (99)		17.3 (79)	21.0 (127)		12.6 (112)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Shenandoah		11.4 (117)	5.7 (53)		11.4 (134)	3.0 (109)	10.2 (19)		10.8 (13)	10.8 (27)		15.9 (134)
Sibley		0.0 (1)	15.8 (134)		6.3 (43)	1.3 (45)	17.5 (129)		18.9 (98)	13.5 (48)		13.2 (118)
Sidney		20.0 (143)	10.0 (106)		7.4 (66)	0.0 (1)	13.0 (53)		30.0 (142)	10.0 (22)		8.1 (60)
Sigourney		16.7 (141)	0.0 (1)		2.8 (9)	2.8 (103)	11.3 (26)		20.7 (110)	13.8 (54)		2.1 (16)
Sioux Cent		6.3 (55)	7.9 (82)		10.0 (124)	0.6 (27)	20.3 (142)		20.9 (111)	25.4 (144)		9.2 (76)
Sioux City		6.9 (62)	6.2 (61)		7.6 (71)	1.3 (45)	17.1 (123)		18.3 (90)	18.1 (114)		8.1 (60)
Spencer		7.5 (68)	10.4 (110)		6.2 (41)	2.7 (101)	15.3 (95)		13.7 (31)	16.5 (91)		10.0 (86)
Spirit Lak		8.9 (83)	8.9 (94)		5.6 (33)	4.3 (134)	13.0 (53)		14.8 (45)	14.8 (65)		3.2 (21)
Storm Lake		8.0 (73)	7.2 (74)		6.2 (41)	2.6 (95)	14.8 (86)		15.7 (59)	17.6 (109)		9.9 (85)
Stuart		50.0 (149)	0.0 (1)		27.8 (149)	0.0 (1)	18.5 (134)		12.5 (23)	16.7 (93)		10.0 (86)
Tama		4.3 (40)	10.9 (113)		5.3 (31)	2.7 (101)	15.0 (90)		18.8 (96)	18.8 (117)		9.4 (80)
Tama/Settl		0.0 (1)	0.0 (1)		7.1 (61)	0.0 (1)	21.4 (148)		28.6 (141)	28.6 (147)		0.0 (1)
Tipton		0.0 (1)	0.0 (1)		5.9 (36)	0.0 (1)	13.7 (72)		27.6 (139)	17.2 (102)		8.1 (60)
Traer		0.0 (1)	0.0 (1)		6.7 (53)	3.3 (116)	13.3 (62)		23.5 (132)	5.9 (10)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
UIHC		87.1 (151)	3.2 (36)		50.0 (151)	7.5 (149)	10.0 (17)		11.1 (17)	22.2 (136)		6.3 (45)
Vinton		3.4 (35)	13.8 (128)		9.0 (105)	2.6 (95)	15.4 (97)		22.8 (129)	15.2 (74)		12.1 (105)
WDM		12.0 (120)	9.6 (101)		7.7 (75)	2.4 (91)	11.8 (34)		16.9 (72)	11.6 (32)		5.3 (34)
Washington		8.3 (77)	8.3 (86)		4.0 (15)	0.5 (22)	11.5 (30)		15.4 (55)	15.4 (76)		5.6 (38)
Waterloo		11.6 (119)	7.5 (77)		8.6 (98)	2.3 (90)	14.0 (76)		16.7 (69)	17.3 (105)		14.2 (125)
Waukee		5.9 (51)	11.8 (119)		4.9 (25)	4.0 (129)	11.3 (26)		12.0 (21)	9.8 (21)		9.4 (80)
Waukon		5.3 (45)	10.5 (111)		4.6 (18)	0.9 (35)	17.4 (125)		21.3 (115)	25.5 (145)		7.0 (52)
Waverly		12.5 (125)	4.2 (39)		6.6 (50)	2.2 (87)	15.9 (102)		14.4 (43)	14.4 (61)		13.6 (122)
Wbstr Ct		10.7 (107)	5.3 (48)		9.2 (108)	2.6 (95)	15.2 (94)		19.0 (100)	14.3 (58)		5.3 (34)
West Union		7.7 (70)	0.0 (1)		9.9 (120)	1.2 (42)	21.0 (145)		38.7 (148)	22.6 (139)		9.3 (77)
Williamsburg		0.0 (1)	0.0 (1)		16.7 (147)	16.7 (151)	16.7 (117)		0.0 (1)	0.0 (1)		0.0 (1)
Winterset		10.0 (97)	6.7 (68)		10.1 (126)	2.2 (87)	17.4 (125)		11.1 (17)	14.8 (65)		8.6 (67)
Iowa		**9.7**	**7.7**		**7.7**	**2.1**	**13.6**		**17.2**	**15.1**		**10.1**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

National (Prior Year)	**128,450**	**10.6**	**6.2**	**541,788**	**7.9**	**3.2**	**10.3**	**226,640**	**15.9**	**12.9**	**372,706**	**14.6**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Ackley		33.3 (148)		50.0 (10)		0.0 (115)		0.0 (131)		0.0 (106)		60.0 (142)		0.0 (1)
Adel		75.0 (50)		22.2 (88)		25.0 (28)		18.2 (68)		11.1 (42)		85.7 (60)		0.0 (1)
Albia		50.0 (138)		7.7 (140)		0.0 (115)		18.2 (68)		0.0 (106)		73.7 (129)		12.9 (138)
Algona		82.8 (21)		33.3 (35)		20.0 (46)		18.2 (68)		13.3 (30)		85.0 (68)		2.9 (94)
Allen WHC		66.4 (102)		26.2 (66)		17.1 (71)		19.3 (62)		8.2 (57)		67.6 (137)		4.6 (118)
Allison		75.0 (50)		21.4 (97)		30.0 (18)		16.7 (76)		14.3 (25)		89.7 (35)		0.8 (42)
Altoona		70.8 (73)		22.2 (88)		17.4 (70)		17.6 (72)		3.7 (92)		100.0 (1)		2.7 (92)
Ames		80.4 (30)		38.4 (30)		29.0 (21)		31.1 (24)		6.8 (70)		81.1 (97)		1.3 (52)
Anamosa		59.3 (124)		24.0 (79)		18.2 (61)		28.0 (34)		16.0 (19)		85.7 (60)		7.6 (132)
Ankeny		74.7 (58)		27.5 (61)		13.9 (93)		11.1 (108)		5.9 (73)		91.7 (24)		0.0 (1)
Atlantic		48.8 (144)		15.6 (124)		10.0 (108)		15.8 (85)		3.1 (96)		78.4 (108)		3.2 (97)
Audubon		70.0 (79)		15.4 (126)		12.5 (97)		11.1 (108)		0.0 (106)		90.2 (32)		1.4 (54)
Avoca		60.0 (122)		100.0 (1)		25.0 (28)		66.7 (3)		50.0 (2)		91.7 (24)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Bedford		81.8 (24)		12.5 (132)		0.0 (115)		28.6 (31)		12.5 (34)		77.3 (117)		0.0 (1)
Belle Plaine		80.0 (31)		20.0 (103)		60.0 (3)		50.0 (6)		0.0 (106)		72.2 (132)		2.0 (74)
Bloomfield		70.0 (79)		50.0 (10)		0.0 (115)		66.7 (3)		50.0 (2)		100.0 (1)		0.0 (1)
Boone		64.3 (113)		36.4 (33)		18.5 (59)		40.0 (12)		18.2 (12)		88.9 (41)		3.9 (112)
Brdlwns RP		76.4 (45)		27.7 (60)		19.3 (56)		11.1 (108)		4.3 (85)		80.7 (98)		1.4 (54)
Brooklyn		44.4 (145)		50.0 (10)		25.0 (28)		0.0 (131)		0.0 (106)		66.7 (138)		0.0 (1)
Burlington		57.5 (125)		19.8 (112)		17.8 (66)		16.1 (83)		5.4 (77)		78.2 (109)		3.5 (108)
CHC		65.2 (106)		17.4 (119)		14.2 (92)		8.4 (123)		5.6 (75)		68.5 (135)		6.1 (126)
CR Urban		70.2 (78)		29.7 (50)		19.5 (54)		12.9 (104)		2.8 (98)		82.6 (86)		1.8 (69)
Carlisle		63.2 (116)		60.0 (5)		30.0 (18)		46.2 (10)		13.3 (30)		92.0 (22)		2.6 (89)
Carroll		71.0 (72)		15.6 (124)		12.1 (103)		20.8 (55)		2.2 (101)		84.0 (77)		2.1 (80)
Cedar Falls		94.7 (5)		58.3 (6)		37.5 (10)		50.0 (6)		8.3 (55)		81.3 (96)		1.1 (46)
Centervill		61.0 (121)		21.7 (96)		0.0 (115)		21.7 (53)		4.3 (85)		100.0 (1)		10.4 (135)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Chariton		70.6 (75)		5.3 (142)		12.5 (97)		15.4 (89)		5.3 (79)		84.6 (73)		0.0 (1)
Charles City		65.7 (104)		22.2 (88)		16.7 (72)		22.0 (52)		4.4 (84)		93.1 (18)		1.2 (48)
Cherokee		81.3 (26)		26.1 (67)		26.3 (27)		28.0 (34)		4.3 (85)		75.7 (120)		5.1 (121)
Clarinda		74.3 (59)		55.6 (8)		18.2 (61)		21.1 (54)		33.3 (4)		78.0 (110)		2.9 (94)
Clarion		71.9 (68)		30.8 (45)		27.8 (22)		19.4 (60)		7.7 (61)		87.1 (54)		2.4 (87)
Clinton		66.5 (101)		23.8 (81)		17.5 (69)		19.4 (60)		8.9 (54)		86.8 (56)		2.6 (89)
Colfax		75.0 (50)		22.2 (88)		0.0 (115)		14.3 (95)		11.1 (42)		86.4 (57)		0.0 (1)
Columbus J		85.7 (10)		40.0 (25)		0.0 (115)		11.1 (108)		0.0 (106)		77.1 (118)		0.0 (1)
Corning		75.0 (50)		30.0 (47)		16.7 (72)		30.0 (27)		10.0 (46)		82.4 (87)		7.0 (129)
Corydon		68.2 (93)		41.7 (22)		0.0 (115)		33.3 (18)		25.0 (8)		87.9 (50)		10.4 (135)
Council Blfs		72.3 (67)		24.0 (79)		18.1 (64)		20.5 (56)		9.9 (51)		81.5 (93)		1.4 (54)
Cresco		83.3 (19)		22.2 (88)		26.7 (26)		23.1 (48)		5.6 (75)		85.0 (68)		17.4 (144)
Creston		78.2 (37)		25.6 (68)		12.9 (96)		18.9 (65)		7.9 (59)		81.5 (93)		5.4 (124)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Dallas Ctr		80.0 (31)		50.0 (10)		25.0 (28)		0.0 (131)		0.0 (106)		0.0 (148)		0.0 (1)
Decorah		79.4 (35)		52.6 (9)		35.7 (12)		18.8 (66)		15.8 (20)		94.1 (16)		14.6 (142)
Denison		67.3 (96)		21.4 (97)		11.4 (104)		4.8 (128)		0.0 (106)		77.9 (111)		0.6 (40)
Dewitt		33.3 (148)		42.9 (21)		0.0 (115)		20.0 (57)		0.0 (106)		93.3 (17)		0.0 (1)
Dubuque		50.2 (137)		16.8 (120)		10.7 (106)		16.0 (84)		2.2 (101)		97.4 (14)		0.1 (39)
Dyersville		50.0 (138)		0.0 (143)		0.0 (115)		33.3 (18)		0.0 (106)		100.0 (1)		0.0 (1)
Eagle Grove		69.7 (85)		9.5 (138)		0.0 (115)		0.0 (131)		0.0 (106)		74.0 (126)		3.3 (100)
Edgerton		69.9 (84)		19.5 (113)		14.5 (87)		13.7 (100)		3.0 (97)		74.0 (126)		4.4 (116)
Eldora		55.6 (129)		14.3 (128)		0.0 (115)		0.0 (131)		0.0 (106)		94.4 (15)		4.9 (119)
Elkader		78.1 (38)		20.8 (102)		0.0 (115)		27.6 (37)		8.3 (55)		92.8 (20)		18.4 (147)
Emmetsburg		85.0 (13)		28.6 (52)		16.7 (72)		23.1 (48)		7.1 (65)		88.9 (41)		15.1 (143)
Esthervill		67.4 (95)		28.0 (58)		15.0 (84)		22.2 (51)		4.0 (89)		84.2 (75)		14.2 (141)
Fairfiel		54.5 (132)		28.6 (52)		0.0 (115)		14.3 (95)		14.3 (25)		50.0 (145)		3.3 (100)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Forest City		64.6 (112)		33.3 (35)		0.0 (115)		24.3 (46)		7.4 (64)		89.6 (36)		1.4 (54)
Fort Dodge		64.9 (108)		16.8 (120)		15.5 (81)		15.5 (88)		4.7 (83)		82.0 (89)		3.1 (96)
Ft Madison		50.0 (138)		11.1 (135)		17.9 (65)		5.0 (127)		3.7 (92)		79.1 (107)		1.4 (54)
Garner		62.5 (119)		20.0 (103)		12.5 (97)		19.0 (64)		15.0 (24)		87.9 (50)		1.5 (61)
Glenwood		68.0 (94)		26.7 (64)		20.0 (46)		0.0 (131)		6.7 (71)		88.9 (41)		0.0 (1)
Grand Junction		80.0 (31)		33.3 (35)		0.0 (115)		50.0 (6)		0.0 (106)		100.0 (1)		8.3 (134)
Greenfield		70.0 (79)		33.3 (35)		30.8 (16)		25.0 (43)		13.3 (30)		89.4 (39)		3.3 (100)
Grinnell		70.0 (79)		28.0 (58)		15.8 (80)		14.3 (95)		12.0 (37)		85.1 (66)		4.2 (114)
Grundy Ctr		70.0 (79)		33.3 (35)		10.0 (108)		10.0 (120)		11.1 (42)		77.8 (112)		1.4 (54)
Guthrie Ct		64.7 (109)		20.0 (103)		20.0 (46)		16.7 (76)		0.0 (106)		77.8 (112)		1.2 (48)
Hampton		65.5 (105)		40.0 (25)		15.4 (82)		23.3 (47)		10.0 (46)		88.0 (49)		1.6 (64)
Harlan		70.7 (74)		10.0 (136)		0.0 (115)		26.9 (41)		5.0 (80)		85.2 (63)		1.7 (66)
Hawarden		84.6 (14)		10.0 (136)		14.3 (89)		27.3 (38)		0.0 (106)		83.3 (82)		2.0 (74)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Hispanic Cln		66.7 (98)		20.0 (103)		50.0 (5)		0.0 (131)		0.0 (106)		91.7 (24)		0.0 (1)
Humboldt		78.3 (36)		17.6 (117)		23.1 (39)		26.7 (42)		11.8 (38)		85.2 (63)		3.4 (105)
ILH		76.2 (46)		24.2 (78)		13.8 (94)		11.9 (106)		4.2 (88)		73.9 (128)		1.9 (73)
Ida Grove		84.4 (16)		25.0 (71)		22.2 (41)		33.3 (18)		12.5 (34)		92.9 (19)		1.8 (69)
Independen		69.2 (86)		25.0 (71)		18.8 (57)		30.4 (26)		0.0 (106)		88.2 (48)		3.8 (111)
Indianola		75.7 (49)		31.0 (44)		34.6 (13)		34.1 (17)		3.4 (94)		85.7 (60)		3.6 (109)
Iowa City		79.7 (34)		45.5 (20)		29.5 (20)		29.0 (30)		11.4 (41)		77.8 (112)		0.7 (41)
Iowa Falls		68.8 (90)		56.3 (7)		16.7 (72)		71.4 (2)		25.0 (8)		90.0 (34)		1.2 (48)
Jamaica		100.0 (1)		0.0 (143)		0.0 (115)		0.0 (131)		0.0 (106)		66.7 (138)		3.4 (105)
Jefferson		81.3 (26)		0.0 (143)		27.3 (23)		5.9 (125)		0.0 (106)		80.7 (98)		2.1 (80)
KR		80.6 (29)		41.7 (22)		18.2 (61)		16.7 (76)		16.7 (16)		68.4 (136)		0.0 (1)
Keokuk		60.0 (122)		16.7 (122)		12.5 (97)		22.5 (50)		4.8 (82)		81.4 (95)		4.1 (113)
Keosauqua		54.5 (132)		50.0 (10)		0.0 (115)		33.3 (18)		16.7 (16)		0.0 (148)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Knoxville		66.4 (102)		25.4 (70)		17.6 (68)		19.2 (63)		9.5 (53)		77.8 (112)		6.1 (126)
LCPH		68.8 (90)		28.6 (52)		12.5 (97)		15.0 (92)		0.0 (106)		70.0 (134)		2.8 (93)
Lamoni		85.7 (10)		33.3 (35)		0.0 (115)		0.0 (131)		0.0 (106)		100.0 (1)		0.0 (1)
Le Mars		87.1 (8)		36.8 (32)		21.4 (44)		13.0 (103)		15.8 (20)		74.8 (124)		0.0 (1)
Lenox		57.1 (126)		0.0 (143)		0.0 (115)		20.0 (57)		0.0 (106)		88.9 (41)		2.2 (83)
Leon		62.5 (119)		17.6 (117)		62.5 (2)		10.5 (117)		0.0 (106)		83.9 (78)		2.1 (80)
Logan		77.8 (39)		18.2 (115)		57.1 (4)		30.8 (25)		18.2 (12)		85.0 (68)		0.0 (1)
Manchester		36.8 (147)		9.1 (139)		0.0 (115)		15.4 (89)		0.0 (106)		98.1 (13)		0.0 (1)
Manning		50.0 (138)		0.0 (143)		0.0 (115)		100.0 (1)		0.0 (106)		100.0 (1)		0.0 (1)
Maquoketa		51.5 (136)		15.0 (127)		0.0 (115)		5.6 (126)		5.0 (80)		82.0 (89)		2.3 (85)
Marengo		71.4 (69)		28.6 (52)		50.0 (5)		0.0 (131)		0.0 (106)		100.0 (1)		0.0 (1)
Marion		65.0 (107)		24.6 (77)		14.3 (89)		15.7 (87)		5.9 (73)		83.7 (79)		1.4 (54)
Marshalltown		73.9 (62)		22.2 (88)		10.7 (106)		17.9 (71)		1.9 (104)		80.7 (98)		2.0 (74)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Mason City		63.0 (118)		22.6 (87)		20.7 (45)		15.4 (89)		9.7 (52)		91.1 (28)		2.2 (83)
Monticello		76.5 (44)		28.6 (52)		0.0 (115)		50.0 (6)		14.3 (25)		92.6 (21)		6.4 (128)
Mt Ayr		56.3 (128)		25.0 (71)		20.0 (46)		27.3 (38)		0.0 (106)		84.6 (73)		0.0 (1)
Mt Pleasan		69.2 (86)		40.0 (25)		16.1 (79)		29.7 (28)		10.0 (46)		86.3 (59)		3.2 (97)
Mtown Dtown		100.0 (1)		0.0 (143)		0.0 (115)		0.0 (131)		0.0 (106)		50.0 (145)		0.0 (1)
Muscatine		76.1 (47)		20.0 (103)		13.2 (95)		14.0 (99)		1.9 (104)		82.0 (89)		2.6 (89)
Nevada		66.7 (98)		16.7 (122)		50.0 (5)		0.0 (131)		0.0 (106)		90.9 (29)		0.0 (1)
New Hampto		77.8 (39)		35.7 (34)		45.5 (9)		16.7 (76)		14.3 (25)		75.0 (122)		0.0 (1)
Newton		75.0 (50)		26.9 (62)		14.6 (86)		13.6 (101)		3.8 (91)		83.6 (81)		1.5 (61)
North Towne		68.9 (89)		21.1 (101)		16.7 (72)		9.0 (122)		7.8 (60)		89.5 (38)		1.0 (43)
Northwood		54.5 (132)		28.6 (52)		22.2 (41)		11.1 (108)		7.1 (65)		87.5 (52)		1.7 (66)
Norwalk		54.5 (132)		30.8 (45)		25.0 (28)		33.3 (18)		7.7 (61)		73.1 (131)		0.0 (1)
Oakridge		50.0 (138)		20.0 (103)		9.1 (111)		0.0 (131)		0.0 (106)		100.0 (1)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Oelwein		68.3 (92)		23.1 (84)		0.0 (115)		17.6 (72)		7.7 (61)		81.9 (92)		19.7 (149)
Onawa		64.3 (113)		6.7 (141)		33.3 (14)		35.3 (15)		0.0 (106)		46.7 (147)		2.0 (74)
Orange Cty		86.4 (9)		38.5 (29)		37.5 (10)		28.6 (31)		15.4 (22)		91.8 (23)		1.7 (66)
Osage		70.3 (77)		26.9 (62)		16.7 (72)		42.3 (11)		11.5 (40)		91.4 (27)		1.2 (48)
Osceola		72.7 (66)		25.0 (71)		8.7 (112)		16.7 (76)		7.1 (65)		80.0 (103)		1.5 (61)
Oskaloosa		54.9 (131)		21.9 (95)		27.3 (23)		18.5 (67)		6.3 (72)		100.0 (1)		2.4 (87)
Ottumwa		71.2 (70)		23.5 (82)		19.4 (55)		17.4 (74)		5.4 (77)		83.3 (82)		1.6 (64)
Pella		75.0 (50)		20.0 (103)		16.7 (72)		0.0 (131)		0.0 (106)		86.4 (57)		0.0 (1)
Perry		77.6 (41)		12.9 (131)		25.0 (28)		4.3 (129)		0.0 (106)		80.0 (103)		3.3 (100)
Pocahontas		63.6 (115)		14.3 (128)		20.0 (46)		0.0 (131)		0.0 (106)		77.8 (112)		5.0 (120)
Pomeroy		0.0 (150)		0.0 (143)		0.0 (115)		0.0 (131)		0.0 (106)		100.0 (1)		0.0 (1)
Postville		75.0 (50)		48.0 (18)		6.7 (114)		39.1 (14)		29.2 (7)		85.2 (63)		4.3 (115)
Red Oak		74.3 (59)		12.0 (134)		18.8 (57)		16.7 (76)		4.0 (89)		88.4 (46)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Redfield		55.6 (129)		0.0 (143)		0.0 (115)		0.0 (131)		0.0 (106)		100.0 (1)		0.0 (1)
Refugee Cln		100.0 (1)		50.0 (10)		80.0 (1)		11.1 (108)		0.0 (106)		54.8 (143)		0.0 (1)
Rock Rapid		64.7 (109)		50.0 (10)		12.5 (97)		11.1 (108)		0.0 (106)		85.0 (68)		1.0 (43)
Rock Valle		81.4 (25)		18.2 (115)		25.0 (28)		25.0 (43)		13.6 (29)		75.4 (121)		1.8 (69)
Rockwell C		64.7 (109)		14.3 (128)		0.0 (115)		29.4 (29)		7.1 (65)		89.3 (40)		5.2 (123)
SDM		73.9 (62)		29.9 (49)		22.8 (40)		9.4 (121)		2.4 (99)		73.2 (130)		1.8 (69)
Sac City		83.3 (19)		50.0 (10)		33.3 (14)		27.3 (38)		16.7 (16)		90.2 (32)		13.1 (139)
Sheldon		85.4 (12)		29.6 (51)		25.0 (28)		14.8 (94)		11.1 (42)		88.9 (41)		18.0 (146)
Shenandoah		74.3 (59)		12.5 (132)		0.0 (115)		4.0 (130)		0.0 (106)		80.4 (101)		2.3 (85)
Sibley		84.2 (18)		0.0 (143)		0.0 (115)		0.0 (131)		0.0 (106)		90.9 (29)		25.6 (151)
Sidney		90.0 (7)		33.3 (35)		20.0 (46)		40.0 (12)		33.3 (4)		61.5 (141)		0.0 (1)
Sigourney		50.0 (138)		20.0 (103)		25.0 (28)		25.0 (43)		0.0 (106)		66.7 (138)		17.4 (144)
Sioux Cent		84.4 (16)		32.3 (42)		25.0 (28)		28.6 (31)		10.0 (46)		87.2 (53)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Sioux City		67.3 (96)		21.3 (99)		17.8 (66)		10.5 (117)		3.4 (94)		79.5 (106)		1.0 (43)
Spencer		82.1 (22)		46.8 (19)		14.3 (89)		35.1 (16)		17.0 (15)		89.6 (36)		19.6 (148)
Spirit Lak		71.1 (71)		38.9 (28)		11.1 (105)		31.6 (23)		17.6 (14)		88.3 (47)		13.9 (140)
Storm Lake		73.9 (62)		21.3 (99)		14.5 (87)		10.3 (119)		2.2 (101)		83.7 (79)		5.5 (125)
Stuart		57.1 (126)		100.0 (1)		0.0 (115)		0.0 (131)		100.0 (1)		83.3 (82)		5.1 (121)
Tama		76.1 (47)		20.0 (103)		15.4 (82)		15.0 (92)		0.0 (106)		85.1 (66)		3.3 (100)
Tama/Settl		100.0 (1)		100.0 (1)		0.0 (115)		66.7 (3)		33.3 (4)		71.4 (133)		7.7 (133)
Tipton		42.9 (146)		37.5 (31)		25.0 (28)		14.3 (95)		25.0 (8)		75.0 (122)		7.1 (130)
Traer		75.0 (50)		25.0 (71)		0.0 (115)		0.0 (131)		25.0 (8)		90.9 (29)		0.0 (1)
UIHC		93.5 (6)		22.2 (88)		0.0 (115)		11.1 (108)		12.5 (34)		0.0 (148)		4.5 (117)
Vinton		69.0 (88)		26.7 (64)		10.0 (108)		15.8 (85)		13.3 (30)		52.6 (144)		3.4 (105)
WDM		82.1 (22)		25.6 (68)		26.9 (25)		11.4 (107)		7.0 (69)		82.4 (87)		1.3 (52)
Washington		81.3 (26)		41.2 (24)		30.4 (17)		16.7 (76)		11.8 (38)		84.2 (75)		7.3 (131)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Waterloo		70.4 (76)		31.2 (43)		18.5 (59)		17.3 (75)		8.2 (57)		76.4 (119)		3.2 (97)
Waukee		76.9 (43)		25.0 (71)		22.2 (41)		11.1 (108)		0.0 (106)		83.3 (82)		1.1 (46)
Waukon		63.2 (116)		30.0 (47)		0.0 (115)		20.0 (57)		10.0 (46)		79.6 (105)		12.7 (137)
Waverly		66.7 (98)		18.8 (114)		8.7 (112)		13.3 (102)		0.0 (106)		87.0 (55)		2.0 (74)
Wbstr Ct		77.3 (42)		22.7 (86)		14.8 (85)		7.9 (124)		2.3 (100)		80.3 (102)		3.7 (110)
West Union		84.6 (14)		23.1 (84)		20.0 (46)		12.5 (105)		15.4 (22)		84.8 (72)		23.5 (150)
Williamsburg		0.0 (150)		100.0 (1)		50.0 (5)		0.0 (131)		0.0 (106)		0.0 (148)		0.0 (1)
Winterset		73.3 (65)		23.5 (82)		20.0 (46)		27.8 (36)		0.0 (106)		74.3 (125)		2.0 (74)
Iowa		**71.0**		**25.8**		**18.1**		**16.6**		**6.0**		**82.9**		**2.9**
National (Prior Year)		**73.5**		**36.7**		**28.0**		**18.1**		**7.9**		**81.4**		**3.3**

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* National includes previous year's data for all participating States and ITOs
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Ackley		0.0 (1)	13.3 (144)		4.3 (17)	2.9 (116)	7.2 (7)		17.2 (81)	0.0 (1)		20.8 (154)
Adel		5.6 (34)	9.3 (106)		4.1 (14)	2.6 (106)	12.0 (44)		19.0 (104)	12.4 (37)		7.4 (51)
Albia		6.8 (58)	11.0 (132)		4.3 (17)	3.4 (129)	11.3 (36)		19.9 (119)	18.4 (122)		11.1 (99)
Algona		7.7 (72)	13.3 (144)		6.8 (60)	1.5 (44)	15.3 (116)		22.0 (140)	20.3 (139)		6.0 (29)
Allen WHC		5.9 (37)	5.6 (39)		6.1 (40)	3.6 (135)	11.1 (29)		15.6 (54)	15.3 (81)		7.6 (55)
Allison		14.9 (152)	8.0 (79)		8.0 (100)	2.1 (73)	11.0 (26)		20.5 (127)	15.2 (78)		6.6 (37)
Altoona		5.9 (37)	6.7 (53)		5.4 (28)	2.1 (73)	13.0 (65)		17.7 (87)	14.9 (74)		10.0 (90)
Ames		11.7 (138)	8.3 (85)		7.3 (74)	3.6 (135)	10.7 (19)		14.1 (24)	11.2 (31)		17.9 (145)
Anamosa		10.3 (123)	5.7 (40)		5.4 (28)	1.7 (51)	12.4 (50)		13.6 (20)	19.3 (130)		16.8 (143)
Ankeny		8.9 (99)	9.4 (109)		7.5 (80)	1.6 (48)	11.1 (29)		14.1 (24)	12.0 (35)		6.5 (35)
Atlantic		10.3 (123)	11.5 (135)		7.4 (78)	0.7 (18)	15.3 (116)		19.9 (119)	15.4 (83)		9.2 (74)
Audubon		6.5 (54)	4.3 (24)		5.6 (35)	0.0 (1)	13.7 (79)		21.2 (133)	14.2 (63)		4.1 (13)
Avoca		8.7 (94)	0.0 (1)		3.4 (9)	2.6 (106)	13.8 (83)		26.3 (150)	15.8 (89)		19.2 (152)
Bedford		3.6 (22)	17.9 (156)		9.4 (123)	2.0 (69)	18.8 (142)		16.2 (63)	20.6 (141)		12.8 (116)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Belle Plaine		10.0 (121)	16.7 (153)		4.4 (19)	0.5 (14)	11.2 (32)		23.7 (146)	16.5 (108)		5.5 (24)
Bloomfield		9.1 (105)	0.0 (1)		6.6 (54)	3.7 (139)	13.2 (69)		23.5 (145)	10.3 (25)		4.2 (15)
Boone		9.2 (108)	9.2 (105)		6.1 (40)	4.4 (153)	8.4 (10)		14.9 (43)	10.2 (23)		13.1 (121)
Brdlwns RP		9.9 (119)	7.1 (62)		7.6 (84)	1.9 (62)	12.9 (60)		16.9 (76)	14.5 (69)		9.6 (82)
Brooklyn		5.9 (37)	5.9 (41)		9.8 (128)	1.6 (48)	18.9 (144)		25.0 (148)	18.3 (120)		14.5 (134)
Burlington		10.3 (123)	8.7 (95)		6.2 (42)	2.7 (111)	10.5 (17)		14.6 (37)	12.8 (42)		12.2 (112)
CHC		12.3 (143)	6.7 (53)		7.7 (86)	2.1 (73)	11.0 (26)		15.3 (48)	15.2 (78)		18.1 (146)
CR Urban		10.0 (121)	8.1 (83)		7.3 (74)	1.7 (51)	12.1 (45)		17.0 (78)	13.4 (48)		19.0 (150)
Carlisle		6.7 (56)	6.7 (53)		10.3 (135)	1.0 (28)	16.1 (126)		22.4 (142)	19.4 (132)		7.9 (61)
Carroll		9.0 (100)	7.1 (62)		6.3 (48)	0.8 (22)	13.0 (65)		18.7 (99)	17.5 (113)		7.8 (57)
Cedar Falls		2.2 (20)	6.5 (50)		3.6 (11)	4.3 (151)	11.1 (29)		8.6 (7)	10.2 (23)		13.9 (131)
Centervill		11.3 (134)	5.2 (36)		10.3 (135)	2.5 (101)	17.5 (136)		14.8 (40)	21.3 (146)		9.9 (87)
Chariton		6.1 (43)	11.2 (133)		5.7 (37)	5.1 (156)	9.7 (11)		17.1 (79)	10.4 (26)		15.2 (136)
Charles City		8.5 (86)	10.8 (126)		5.1 (24)	2.4 (91)	11.6 (41)		20.6 (128)	15.2 (78)		7.2 (47)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Cherokee		9.2 (108)	4.9 (33)		6.3 (48)	0.9 (25)	17.6 (137)		20.8 (130)	18.7 (125)		8.0 (62)
Clarinda		5.9 (37)	5.9 (41)		7.9 (94)	2.2 (80)	13.2 (69)		16.3 (66)	14.0 (60)		12.9 (118)
Clarion		6.3 (49)	10.8 (126)		11.6 (147)	2.0 (69)	19.1 (146)		19.3 (110)	16.4 (105)		3.9 (12)
Clinton		11.4 (136)	7.5 (67)		9.1 (118)	1.1 (31)	14.2 (92)		20.8 (130)	16.0 (96)		10.9 (96)
Colfax		4.3 (25)	10.9 (128)		9.3 (121)	2.4 (91)	15.6 (121)		14.4 (33)	20.0 (134)		11.7 (107)
Columbus J		7.8 (74)	5.9 (41)		4.0 (13)	2.4 (91)	16.6 (130)		16.8 (74)	21.2 (144)		6.7 (39)
Corning		5.3 (33)	10.5 (122)		9.9 (130)	2.7 (111)	13.7 (79)		19.1 (107)	9.0 (12)		15.2 (136)
Correctionvl		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)	33.3 (157)		28.6 (151)	42.9 (155)		20.0 (153)
Corydon		6.7 (56)	16.7 (153)		6.2 (42)	3.5 (133)	15.2 (114)		14.1 (24)	18.8 (127)		12.2 (112)
Council Blfs		9.7 (117)	7.6 (69)		9.2 (119)	1.8 (56)	14.2 (92)		18.3 (94)	16.2 (103)		16.2 (139)
Cresco		10.9 (132)	12.0 (139)		4.8 (22)	2.0 (69)	12.8 (59)		11.3 (13)	15.1 (76)		8.6 (69)
Creston		9.3 (112)	4.7 (29)		10.6 (140)	2.2 (80)	15.6 (121)		18.2 (92)	15.6 (87)		11.1 (99)
Dallas Ctr		0.0 (1)	9.1 (99)		16.2 (154)	0.0 (1)	5.4 (2)		0.0 (1)	8.3 (10)		0.0 (1)
Dayton		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)	50.0 (158)		0.0 (1)	100.0 (157)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Decorah		8.6 (92)	14.3 (148)		6.9 (65)	0.9 (25)	13.7 (79)		18.4 (95)	16.9 (110)		7.5 (53)
Denison		8.3 (82)	6.0 (44)		7.8 (89)	1.5 (44)	18.3 (140)		18.4 (95)	20.7 (142)		13.5 (129)
Dewitt		27.3 (155)	6.1 (46)		7.8 (89)	1.0 (28)	14.7 (108)		25.9 (149)	9.9 (21)		9.4 (78)
Dubuque		8.0 (78)	7.7 (70)		6.5 (52)	3.0 (119)	8.1 (9)		15.5 (51)	9.7 (17)		18.1 (146)
Dyersville		0.0 (1)	9.1 (99)		2.2 (7)	1.1 (31)	13.2 (69)		4.2 (6)	22.9 (150)		13.4 (127)
Eagle Grove		9.2 (108)	6.9 (57)		7.9 (94)	2.5 (101)	18.1 (139)		16.4 (68)	18.7 (125)		4.8 (17)
Edgerton		10.3 (123)	7.3 (64)		8.9 (114)	1.6 (48)	11.3 (36)		17.4 (84)	13.6 (55)		12.1 (110)
Eldora		11.8 (139)	8.8 (96)		7.0 (66)	2.1 (73)	14.4 (102)		17.8 (88)	18.9 (128)		10.6 (94)
Elkader		10.8 (131)	7.7 (70)		7.0 (66)	1.2 (36)	11.2 (32)		20.4 (125)	10.6 (27)		5.4 (22)
Emmetsburg		14.0 (148)	4.7 (29)		12.0 (149)	2.9 (116)	13.1 (68)		14.1 (24)	13.4 (48)		5.6 (26)
Esthervill		6.8 (58)	9.5 (110)		8.5 (111)	3.1 (122)	12.7 (56)		19.0 (104)	16.4 (105)		7.1 (44)
Fairfiel		6.4 (53)	10.3 (120)		8.9 (114)	2.2 (80)	17.2 (132)		14.8 (40)	18.4 (122)		7.0 (42)
Fayette		0.0 (1)	20.0 (157)		12.0 (149)	3.6 (135)	13.3 (73)		21.4 (134)	14.3 (66)		8.7 (70)
Forest City		6.9 (61)	9.9 (114)		7.8 (89)	1.2 (36)	11.2 (32)		18.7 (99)	9.5 (16)		5.9 (27)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Fort Dodge		7.7 (72)	7.5 (67)		10.0 (131)	1.9 (62)	17.2 (132)		19.8 (117)	18.2 (118)		4.6 (16)
Ft Madison		10.6 (129)	7.8 (74)		8.7 (112)	1.8 (56)	13.4 (77)		12.5 (15)	14.9 (74)		7.8 (57)
Garner		4.7 (27)	11.6 (136)		5.2 (25)	2.3 (86)	11.4 (38)		15.7 (56)	14.0 (60)		8.4 (68)
Glenwood		7.3 (67)	3.7 (20)		9.4 (123)	2.6 (106)	12.3 (47)		14.9 (43)	13.5 (53)		12.1 (110)
Grand Junction		8.3 (82)	0.0 (1)		25.5 (156)	0.0 (1)	14.9 (111)		11.1 (12)	16.7 (109)		3.3 (11)
Greenfield		7.3 (67)	9.1 (99)		6.8 (60)	3.4 (129)	14.3 (97)		10.9 (11)	9.4 (14)		17.0 (144)
Grinnell		7.1 (64)	4.4 (26)		11.0 (146)	4.0 (145)	11.5 (39)		18.7 (99)	9.3 (13)		21.3 (155)
Grundy Ctr		3.3 (21)	13.3 (144)		6.8 (60)	2.4 (91)	11.2 (32)		17.8 (88)	10.0 (22)		11.9 (109)
Guthrie Ct		5.7 (36)	8.6 (94)		6.4 (51)	0.3 (13)	10.8 (21)		20.9 (132)	12.0 (35)		6.5 (35)
Hampton		5.2 (32)	13.5 (147)		5.6 (35)	1.4 (42)	15.5 (120)		22.6 (143)	18.1 (117)		8.3 (66)
Harlan		8.5 (86)	4.2 (23)		7.4 (78)	1.3 (40)	14.3 (97)		17.4 (84)	13.4 (48)		8.3 (66)
Hawarden		13.2 (147)	11.8 (137)		8.3 (107)	0.7 (18)	19.4 (147)		17.1 (79)	20.0 (134)		9.3 (76)
Hispanic Cln		5.0 (29)	15.0 (151)		6.2 (42)	1.4 (42)	19.9 (148)		28.7 (152)	21.8 (147)		8.9 (71)
Humboldt		7.9 (75)	7.9 (77)		7.9 (94)	1.8 (56)	19.0 (145)		19.5 (113)	22.1 (148)		4.8 (17)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
ILH		10.9 (132)	7.7 (70)		7.6 (84)	1.5 (44)	13.9 (86)		18.0 (90)	16.1 (99)		11.1 (99)
Ida Grove		6.3 (49)	2.1 (15)		7.7 (86)	3.7 (139)	16.2 (127)		10.8 (10)	16.3 (104)		7.0 (42)
Independen		9.9 (119)	10.7 (124)		8.4 (109)	3.4 (129)	12.9 (60)		20.3 (124)	14.5 (69)		11.7 (107)
Indianola		10.4 (128)	12.2 (140)		10.9 (144)	2.3 (86)	15.2 (114)		16.9 (76)	17.5 (113)		11.4 (105)
Iowa City		12.0 (141)	6.6 (52)		6.8 (60)	3.2 (125)	11.0 (26)		15.7 (56)	12.5 (39)		9.9 (87)
Iowa Falls		8.5 (86)	5.1 (35)		8.4 (109)	2.4 (91)	10.8 (21)		17.2 (81)	12.6 (40)		19.1 (151)
Jamaica		0.0 (1)	9.1 (99)		5.4 (28)	3.2 (125)	12.9 (60)		10.3 (9)	23.1 (151)		13.1 (121)
Jefferson		8.0 (78)	9.3 (106)		6.2 (42)	1.2 (36)	18.6 (141)		20.0 (121)	20.0 (134)		5.9 (27)
KR		12.5 (144)	10.9 (128)		6.5 (52)	3.3 (127)	10.3 (15)		14.4 (33)	11.2 (31)		10.6 (94)
Keokuk		8.5 (86)	4.0 (22)		10.7 (141)	2.3 (86)	13.3 (73)		20.7 (129)	16.1 (99)		14.1 (132)
Keosauqua		0.0 (1)	16.0 (152)		3.5 (10)	1.8 (56)	24.6 (152)		15.5 (51)	31.0 (153)		5.5 (24)
Knoxville		9.6 (115)	9.9 (114)		9.2 (119)	3.1 (122)	13.3 (73)		14.1 (24)	16.0 (96)		9.2 (74)
LCPH		6.3 (49)	6.3 (48)		10.5 (138)	3.5 (133)	7.0 (5)		45.5 (157)	0.0 (1)		27.8 (156)
Lamoni		0.0 (1)	8.0 (79)		13.0 (151)	0.6 (17)	18.8 (142)		14.1 (24)	16.9 (110)		14.7 (135)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Le Mars		8.7 (94)	6.9 (57)		9.6 (127)	1.7 (51)	15.6 (121)		19.8 (117)	13.6 (55)		7.1 (44)
Lenox		10.3 (123)	10.3 (120)		10.3 (135)	0.5 (14)	14.4 (102)		22.1 (141)	14.0 (60)		18.3 (148)
Leon		6.1 (43)	6.1 (46)		10.2 (134)	2.5 (101)	14.4 (102)		18.2 (92)	16.1 (99)		16.2 (139)
Logan		8.8 (96)	9.6 (113)		7.8 (89)	2.8 (115)	17.4 (135)		16.5 (70)	13.5 (53)		16.2 (139)
Luana		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)		0.0 (1)
Manchester		9.1 (105)	10.6 (123)		4.2 (15)	3.4 (129)	12.9 (60)		18.5 (97)	15.8 (89)		13.0 (120)
Manning		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)	25.0 (153)		16.7 (72)	33.3 (154)		0.0 (1)
Maquoketa		4.1 (24)	8.3 (85)		10.5 (138)	1.7 (51)	13.0 (65)		20.2 (122)	13.0 (44)		6.1 (30)
Marengo		6.3 (49)	12.5 (141)		7.5 (80)	2.2 (80)	12.9 (60)		23.8 (147)	15.5 (84)		1.0 (7)
Marion		8.5 (86)	9.5 (110)		7.1 (68)	1.1 (31)	12.5 (53)		16.3 (66)	14.5 (69)		8.9 (71)
Marshalltown		7.9 (75)	8.4 (88)		5.4 (28)	3.0 (119)	13.5 (78)		16.8 (74)	15.1 (76)		15.3 (138)
Mason City		7.5 (70)	8.4 (88)		7.1 (68)	2.3 (86)	12.3 (47)		19.7 (116)	16.1 (99)		7.2 (47)
Monticello		0.0 (1)	7.7 (70)		4.9 (23)	2.0 (69)	10.2 (14)		13.8 (22)	13.8 (57)		11.2 (102)
Mt Ayr		6.1 (43)	3.0 (16)		10.9 (144)	1.1 (31)	14.7 (108)		15.9 (60)	13.4 (48)		9.8 (86)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Mt Pleasan		9.0 (100)	8.5 (91)		9.3 (121)	2.1 (73)	16.0 (125)		15.6 (54)	19.3 (130)		9.4 (78)
Mtown Dtown		0.0 (1)	0.0 (1)		14.3 (153)	0.0 (1)	14.3 (97)		50.0 (158)	0.0 (1)		50.0 (158)
Murray		50.0 (157)	0.0 (1)		28.6 (157)	0.0 (1)	7.1 (6)		14.3 (31)	0.0 (1)		10.0 (90)
Muscatine		8.6 (92)	7.8 (74)		5.7 (37)	2.6 (106)	12.7 (56)		18.7 (99)	18.3 (120)		12.6 (115)
Nevada		5.0 (29)	10.0 (117)		4.2 (15)	8.4 (158)	10.9 (24)		15.8 (58)	15.8 (89)		18.5 (149)
New Hampto		8.8 (96)	8.8 (96)		7.2 (72)	2.2 (80)	15.0 (112)		18.1 (91)	13.9 (59)		8.2 (64)
Newton		9.6 (115)	9.9 (114)		7.7 (86)	3.9 (143)	13.3 (73)		14.1 (24)	14.2 (63)		13.5 (129)
North Towne		8.5 (86)	8.8 (96)		7.3 (74)	1.8 (56)	11.5 (39)		19.1 (107)	9.8 (19)		13.4 (127)
Northwood		5.9 (37)	11.8 (137)		7.1 (68)	1.9 (62)	7.8 (8)		16.2 (63)	7.4 (7)		6.1 (30)
Norwalk		20.0 (154)	10.9 (128)		13.2 (152)	4.0 (145)	14.0 (90)		13.2 (16)	15.8 (89)		16.4 (142)
Oakridge		0.0 (1)	0.0 (1)		1.5 (6)	2.9 (116)	5.9 (3)		22.6 (143)	9.7 (17)		9.3 (76)
Oelwein		6.8 (58)	10.2 (119)		5.3 (26)	0.7 (18)	13.9 (86)		15.4 (50)	15.8 (89)		7.2 (47)
Onawa		6.1 (43)	3.1 (18)		7.3 (74)	2.1 (73)	15.4 (119)		16.4 (68)	18.2 (118)		10.9 (96)
Orange Cty		6.1 (43)	14.9 (150)		6.8 (60)	2.5 (101)	13.8 (83)		15.5 (51)	10.9 (28)		9.6 (82)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Osage		8.3 (82)	6.5 (50)		8.0 (100)	2.4 (91)	10.9 (24)		14.6 (37)	9.4 (14)		7.8 (57)
Osceola		12.7 (146)	7.8 (74)		6.6 (54)	3.0 (119)	13.7 (79)		18.8 (103)	15.5 (84)		14.1 (132)
Oskaloosa		11.4 (136)	4.3 (24)		8.1 (105)	4.3 (151)	12.4 (50)		16.5 (70)	12.9 (43)		8.0 (62)
Ottumwa		9.0 (100)	8.0 (79)		8.7 (112)	2.6 (106)	14.6 (107)		17.3 (83)	15.9 (95)		7.7 (56)
Pella		12.5 (144)	10.9 (128)		8.9 (114)	2.5 (101)	14.2 (92)		15.8 (58)	15.8 (89)		13.2 (125)
Perry		7.9 (75)	8.4 (88)		6.6 (54)	1.0 (28)	16.3 (128)		21.6 (135)	21.2 (144)		6.7 (39)
Pocahontas		7.1 (64)	4.8 (31)		11.6 (147)	4.4 (153)	13.8 (83)		20.2 (122)	13.4 (48)		9.9 (87)
Pomeroy		0.0 (1)	0.0 (1)		5.3 (26)	0.0 (1)	31.6 (156)		0.0 (1)	50.0 (156)		0.0 (1)
Postville		9.0 (100)	3.0 (16)		8.3 (107)	2.4 (91)	15.8 (124)		15.3 (48)	19.5 (133)		9.4 (78)
Red Oak		10.7 (130)	10.7 (124)		9.4 (123)	2.1 (73)	12.3 (47)		14.6 (37)	9.8 (19)		13.1 (121)
Redfield		14.3 (149)	0.0 (1)		8.0 (100)	4.0 (145)	6.0 (4)		36.4 (155)	4.5 (6)		3.2 (10)
Refugee Cln		5.9 (37)	17.6 (155)		3.3 (8)	2.4 (91)	12.2 (46)		10.2 (8)	16.9 (110)		29.0 (157)
Rock Rapid		7.4 (69)	7.4 (65)		9.5 (126)	0.5 (14)	10.3 (15)		13.7 (21)	7.7 (8)		5.0 (20)
Rock Valle		6.1 (43)	7.9 (77)		7.1 (68)	0.9 (25)	16.5 (129)		14.5 (36)	13.3 (45)		7.4 (51)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Rockwell C		14.6 (151)	14.6 (149)		7.9 (94)	0.7 (18)	16.8 (131)		19.3 (110)	17.9 (116)		2.5 (9)
SDM		9.2 (108)	9.5 (110)		6.6 (54)	1.5 (44)	13.2 (69)		16.2 (63)	15.5 (84)		7.8 (57)
Sac City		7.0 (62)	7.0 (61)		5.5 (33)	0.8 (22)	15.1 (113)		21.9 (139)	21.1 (143)		4.1 (13)
Sheldon		9.0 (100)	5.4 (37)		7.5 (80)	3.8 (141)	14.2 (92)		15.2 (47)	16.4 (105)		8.2 (64)
Shenandoah		11.8 (139)	3.9 (21)		9.0 (117)	3.8 (141)	10.0 (12)		14.3 (31)	11.2 (31)		12.9 (118)
Sibley		3.8 (23)	11.3 (134)		6.2 (42)	1.9 (62)	14.3 (97)		19.5 (113)	11.5 (34)		6.2 (33)
Sidney		4.9 (28)	4.9 (33)		10.1 (133)	1.7 (51)	12.7 (56)		20.4 (125)	8.3 (10)		9.7 (85)
Sigourney		8.3 (82)	8.3 (85)		6.2 (42)	2.4 (91)	14.3 (97)		14.4 (33)	13.3 (45)		5.0 (20)
Sioux Cent		8.0 (78)	6.4 (49)		10.8 (143)	1.1 (31)	20.2 (150)		21.7 (136)	22.5 (149)		7.1 (44)
Sioux City		7.1 (64)	6.9 (57)		7.9 (94)	0.8 (22)	17.8 (138)		19.6 (115)	18.9 (128)		9.0 (73)
Southridge		9.1 (105)	9.3 (106)		5.9 (39)	2.2 (80)	11.7 (42)		19.4 (112)	14.4 (68)		13.1 (121)
Spencer		7.5 (70)	6.7 (53)		7.8 (89)	1.9 (62)	12.6 (54)		13.5 (19)	12.6 (40)		6.6 (37)
Spirit Lak		9.4 (113)	8.5 (91)		7.2 (72)	3.6 (135)	10.6 (18)		16.7 (72)	10.9 (28)		5.4 (22)
Storm Lake		7.0 (62)	6.9 (57)		6.3 (48)	2.7 (111)	13.9 (86)		14.9 (43)	14.7 (73)		7.5 (53)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese		≥ 2 Yrs Overweight and Obese			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Stuart		35.7 (156)	3.6 (19)		18.2 (155)	0.0 (1)	21.5 (151)		14.8 (40)	20.4 (140)		12.2 (112)
Tama		8.0 (78)	12.6 (142)		4.7 (21)	3.3 (127)	14.2 (92)		18.6 (98)	18.6 (124)		11.3 (103)
Tama/Settl		0.0 (1)	0.0 (1)		5.5 (33)	4.1 (150)	26.0 (155)		40.0 (156)	20.0 (134)		13.2 (125)
Tipton		0.0 (1)	4.5 (27)		3.8 (12)	1.9 (62)	14.5 (106)		32.5 (154)	15.6 (87)		6.9 (41)
Traer		18.2 (153)	9.1 (99)		10.0 (131)	4.0 (145)	10.0 (12)		21.7 (136)	10.9 (28)		2.3 (8)
UIHC		83.9 (158)	5.4 (37)		45.8 (158)	6.7 (157)	10.8 (21)		16.0 (61)	8.0 (9)		6.4 (34)
Vinton		5.1 (31)	12.8 (143)		9.8 (128)	3.9 (143)	13.9 (86)		21.7 (136)	14.2 (63)		10.4 (93)
WDM		11.3 (134)	10.1 (118)		6.7 (58)	2.3 (86)	12.4 (50)		14.0 (23)	13.8 (57)		9.6 (82)
Wapello		0.0 (1)	0.0 (1)		0.0 (1)	0.0 (1)	25.0 (153)		0.0 (1)	100.0 (157)		0.0 (1)
Washington		6.6 (55)	8.1 (83)		5.4 (28)	1.8 (56)	10.7 (19)		19.2 (109)	14.3 (66)		4.9 (19)
Waterloo		9.8 (118)	7.4 (65)		7.9 (94)	3.1 (122)	12.6 (54)		16.0 (61)	15.3 (81)		12.8 (116)
Waukee		9.5 (114)	8.0 (79)		6.7 (58)	2.7 (111)	11.7 (42)		11.5 (14)	13.3 (45)		10.2 (92)
Waukon		4.5 (26)	9.1 (99)		4.5 (20)	1.3 (40)	14.1 (91)		13.3 (17)	17.7 (115)		7.2 (47)
Waverly		12.2 (142)	8.5 (91)		8.0 (100)	1.9 (62)	14.8 (110)		19.0 (104)	12.4 (37)		11.3 (103)

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 6A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Growth and Anemia Indicators by Clinic

Children Aged < 5 Years (2)

Combined 3 Years

Comparison of Growth and Anemia Indicators (3,4)

Clinic	Birth Weight (5)			Height and Weight (6,7)							Anemia Low Hb/Hct (8)	
		Low	High		Short Stature	Under-Weight	Obese	≥ 2 Yrs Overweight and Obese				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% (Rank)	% (Rank)	% (Rank)	Number	% Overweight (Rank)	% Obese (Rank)	Number	% (Rank)
Wbstr Ct		8.8 (96)	6.0 (44)		8.2 (106)	2.4 (91)	15.3 (116)		17.5 (86)	16.0 (96)		6.1 (30)
West Union		14.3 (149)	4.8 (31)		10.7 (141)	1.2 (36)	17.2 (132)		29.5 (153)	24.6 (152)		11.0 (98)
Williamsburg		0.0 (1)	100.0 (158)		8.0 (100)	4.0 (145)	20.0 (149)		13.3 (17)	20.0 (134)		9.5 (81)
Winterset		5.6 (34)	4.5 (27)		7.5 (80)	4.9 (155)	14.4 (102)		15.1 (46)	14.5 (69)		11.4 (105)
Iowa		**9.4**	**7.8**		**7.6**	**2.2**	**13.3**		**17.3**	**16.0**		**11.2**

(1) Reporting period is January 1 through December 31.
(2) Analyses based on one record per child.
(3) Excludes records with unknown data or errors.
(4) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(5) Infants born during the reporting period included in the analysis.
(6) Based on 2006 WHO growth chart percentiles for children under 2 years of age; short stature is defined as length-for-age <= 2.3rd percentile, underweight is defined as weight-for-length <= 2.3rd percentile, and high weight-for-length (labeled as "Obese") is defined as >= 97.7th percentile.
(7) Based on 2000 CDC growth chart percentiles for children 2 years of age and older; short stature is defined as height-for-age < 5th percentile, underweight is defined as BMI-for-age < 5th percentile, overweight is defined as BMI-for-age >= 85th to < 95th percentile, and obesity is defined as >= 95th percentile.
(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Ackley		73.3 (58)		38.5 (11)		0.0 (136)		25.0 (28)		0.0 (131)		63.6 (154)		2.2 (67)
Adel		70.4 (76)		14.8 (135)		11.8 (107)		12.1 (118)		3.7 (105)		91.5 (26)		5.4 (129)
Albia		55.4 (143)		7.5 (148)		9.1 (122)		13.2 (114)		2.5 (117)		79.0 (129)		3.9 (113)
Algona		68.7 (91)		26.3 (53)		17.6 (52)		18.3 (71)		8.8 (37)		83.8 (87)		2.3 (73)
Allen WHC		70.8 (72)		29.9 (30)		15.7 (69)		21.5 (49)		12.8 (13)		73.9 (145)		4.8 (126)
Allison		73.9 (53)		10.8 (143)		15.6 (70)		23.4 (38)		5.4 (72)		87.5 (51)		1.6 (46)
Altoona		71.4 (62)		27.8 (45)		18.9 (43)		13.8 (106)		3.8 (102)		95.8 (14)		2.2 (67)
Ames		78.7 (28)		41.8 (8)		27.4 (14)		29.8 (15)		6.0 (63)		84.1 (85)		1.6 (46)
Anamosa		68.2 (94)		16.7 (122)		22.7 (23)		13.5 (111)		8.3 (42)		92.1 (22)		2.8 (88)
Ankeny		78.1 (31)		27.8 (45)		19.8 (37)		16.5 (86)		8.7 (39)		77.5 (138)		0.8 (26)
Atlantic		61.6 (131)		21.2 (88)		17.4 (56)		17.6 (76)		5.9 (66)		77.9 (136)		3.5 (107)
Audubon		76.1 (42)		21.2 (88)		8.3 (127)		16.0 (92)		0.0 (131)		85.0 (75)		2.5 (83)
Avoca		73.9 (53)		25.0 (65)		25.0 (20)		33.3 (10)		8.3 (42)		83.3 (95)		2.3 (73)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Bedford		60.7 (134)		10.5 (145)		22.2 (25)		16.7 (83)		10.5 (26)		68.8 (150)		0.0 (1)
Belle Plaine		71.0 (69)		35.7 (18)		27.8 (13)		26.7 (24)		7.1 (55)		78.5 (133)		2.0 (62)
Bloomfield		60.9 (133)		18.8 (111)		12.5 (94)		11.8 (121)		12.5 (14)		100.0 (1)		0.0 (1)
Boone		69.7 (83)		21.5 (86)		8.7 (125)		30.2 (14)		7.7 (50)		89.2 (37)		3.4 (105)
Brdlwns RP		78.2 (30)		29.8 (31)		20.4 (32)		9.5 (135)		3.0 (110)		79.6 (125)		1.7 (50)
Brooklyn		52.6 (148)		17.6 (118)		6.3 (131)		0.0 (148)		5.9 (66)		88.7 (39)		3.7 (110)
Burlington		58.3 (139)		15.8 (129)		12.2 (102)		13.6 (107)		4.9 (83)		78.9 (130)		4.4 (121)
CHC		61.6 (131)		17.3 (120)		11.3 (112)		9.8 (133)		4.7 (85)		67.9 (152)		5.8 (133)
CR Urban		66.8 (104)		26.5 (49)		17.8 (51)		11.2 (125)		2.4 (118)		82.0 (111)		1.9 (56)
Carlisle		68.3 (93)		38.6 (10)		28.0 (12)		25.6 (27)		11.4 (21)		92.3 (21)		1.0 (29)
Carroll		71.3 (65)		15.4 (133)		9.7 (119)		15.9 (93)		2.7 (115)		84.0 (86)		2.7 (87)
Cedar Falls		87.5 (6)		64.3 (2)		41.2 (3)		51.7 (2)		28.6 (2)		82.4 (105)		0.5 (20)
Centervill		58.8 (137)		20.4 (93)		0.0 (136)		16.1 (91)		4.1 (97)		94.1 (19)		5.9 (135)
Chariton		66.3 (112)		15.8 (129)		16.7 (60)		24.5 (33)		5.3 (73)		85.4 (71)		3.1 (96)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Charles City		66.0 (113)		22.7 (79)		18.6 (46)		21.1 (52)		5.3 (73)		89.4 (36)		1.6 (46)
Cherokee		76.8 (39)		19.4 (105)		12.3 (100)		14.8 (100)		4.5 (90)		79.2 (127)		4.3 (120)
Clarinda		68.2 (94)		24.2 (73)		10.0 (117)		26.1 (25)		12.1 (18)		75.0 (142)		3.8 (111)
Clarion		70.9 (70)		20.4 (93)		15.5 (72)		10.4 (130)		2.9 (111)		88.1 (45)		2.4 (79)
Clinton		65.5 (118)		19.1 (108)		13.8 (84)		15.0 (98)		8.9 (36)		89.6 (34)		3.2 (101)
Colfax		66.7 (105)		16.7 (122)		0.0 (136)		12.5 (117)		4.2 (93)		88.4 (43)		2.2 (67)
Columbus J		75.0 (47)		28.0 (41)		0.0 (136)		3.8 (147)		0.0 (131)		85.2 (73)		2.8 (88)
Corning		68.4 (92)		18.2 (116)		14.3 (78)		20.0 (56)		9.1 (33)		73.9 (145)		5.7 (131)
Correctionvl		100.0 (1)		33.3 (21)		0.0 (136)		25.0 (28)		33.3 (1)		100.0 (1)		0.0 (1)
Corydon		58.3 (139)		35.0 (19)		21.4 (30)		33.3 (10)		17.5 (6)		84.7 (77)		9.4 (142)
Council Blfs		73.2 (59)		22.5 (80)		17.3 (57)		17.3 (77)		8.1 (48)		81.4 (114)		1.5 (44)
Cresco		79.3 (21)		26.5 (49)		22.2 (25)		27.4 (21)		4.1 (97)		84.8 (76)		15.4 (151)
Creston		66.6 (111)		21.6 (85)		13.5 (85)		16.5 (86)		6.3 (60)		79.5 (126)		4.4 (121)
Dallas Ctr		90.9 (5)		66.7 (1)		20.0 (33)		0.0 (148)		0.0 (131)		100.0 (1)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Dayton		0.0 (155)		0.0 (150)		0.0 (136)		0.0 (148)		0.0 (131)		100.0 (1)		0.0 (1)
Decorah		80.0 (17)		44.2 (6)		27.3 (15)		29.6 (16)		19.2 (5)		95.7 (16)		14.2 (149)
Denison		65.9 (115)		16.3 (127)		9.5 (121)		6.4 (143)		2.0 (125)		75.6 (141)		0.6 (22)
Dewitt		66.7 (105)		29.6 (32)		18.2 (47)		16.7 (83)		3.7 (105)		95.2 (17)		1.7 (50)
Dubuque		54.4 (145)		15.7 (132)		11.9 (106)		15.4 (94)		2.9 (111)		95.8 (14)		0.1 (19)
Dyersville		54.5 (144)		20.0 (98)		33.3 (6)		22.2 (46)		0.0 (131)		97.9 (12)		0.0 (1)
Eagle Grove		71.3 (65)		20.8 (91)		13.2 (88)		9.5 (135)		0.0 (131)		86.4 (64)		2.3 (73)
Edgerton		65.2 (119)		20.3 (95)		12.0 (104)		12.1 (118)		4.4 (91)		76.8 (139)		3.9 (113)
Eldora		52.9 (147)		13.0 (139)		0.0 (136)		4.8 (145)		0.0 (131)		88.6 (40)		5.3 (127)
Elkader		83.2 (12)		20.2 (96)		4.8 (134)		18.0 (74)		4.8 (84)		91.7 (24)		18.5 (155)
Emmetsburg		79.1 (23)		23.1 (78)		14.3 (78)		17.0 (81)		5.8 (68)		87.3 (54)		11.2 (145)
Esthervill		65.8 (117)		23.8 (74)		16.0 (68)		22.5 (45)		9.5 (30)		86.8 (58)		12.4 (147)
Fairfiel		56.4 (142)		19.5 (104)		9.7 (119)		17.1 (80)		12.2 (17)		68.8 (150)		5.3 (127)
Fayette		66.7 (105)		33.3 (21)		20.0 (33)		23.1 (41)		6.7 (57)		73.1 (147)		24.4 (158)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Forest City		62.9 (127)		17.5 (119)		9.1 (122)		17.0 (81)		3.8 (102)		86.0 (66)		1.8 (53)
Fort Dodge		67.3 (100)		19.7 (103)		12.8 (92)		16.7 (83)		5.5 (71)		86.8 (58)		4.0 (116)
Ft Madison		57.1 (141)		13.4 (138)		12.1 (103)		11.5 (124)		4.7 (85)		78.9 (130)		3.3 (104)
Garner		67.8 (98)		23.8 (74)		8.3 (127)		22.0 (48)		11.9 (19)		85.7 (67)		1.6 (46)
Glenwood		71.1 (67)		16.3 (127)		5.0 (133)		20.0 (56)		4.7 (85)		84.6 (80)		0.0 (1)
Grand Junction		66.7 (105)		33.3 (21)		0.0 (136)		37.5 (7)		0.0 (131)		88.2 (44)		7.5 (139)
Greenfield		67.3 (100)		28.1 (40)		22.7 (23)		28.2 (18)		6.3 (60)		80.4 (121)		3.6 (108)
Grinnell		72.6 (60)		25.3 (64)		16.7 (60)		21.3 (50)		6.0 (63)		83.8 (87)		3.8 (111)
Grundy Ctr		70.0 (78)		28.0 (41)		8.0 (129)		13.6 (107)		4.0 (99)		82.3 (107)		1.9 (56)
Guthrie Ct		74.3 (52)		25.6 (60)		18.2 (47)		18.6 (69)		5.1 (79)		81.3 (115)		3.2 (101)
Hampton		70.8 (72)		30.8 (28)		14.0 (83)		22.1 (47)		9.0 (35)		88.9 (38)		1.8 (53)
Harlan		67.2 (102)		22.4 (82)		12.3 (100)		26.0 (26)		7.5 (52)		83.5 (93)		1.2 (36)
Hawarden		78.9 (27)		22.5 (80)		2.3 (135)		19.0 (67)		5.0 (81)		90.5 (28)		1.1 (33)
Hispanic Cln		70.0 (78)		37.5 (15)		40.0 (4)		7.1 (142)		0.0 (131)		89.8 (33)		0.0 (1)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Humboldt		80.9 (15)		26.5 (49)		13.2 (88)		22.8 (44)		14.3 (10)		86.3 (65)		2.9 (92)
ILH		76.2 (41)		24.3 (72)		16.7 (60)		8.5 (140)		2.4 (118)		82.2 (108)		2.2 (67)
Ida Grove		75.8 (45)		19.1 (108)		11.4 (111)		18.2 (72)		8.5 (41)		93.0 (20)		1.0 (29)
Independen		64.8 (120)		17.1 (121)		11.7 (109)		20.9 (54)		1.2 (130)		83.3 (95)		4.0 (116)
Indianola		76.1 (42)		26.5 (49)		24.3 (21)		25.0 (28)		5.1 (79)		88.1 (45)		3.1 (96)
Iowa City		80.2 (16)		45.6 (5)		31.1 (8)		23.1 (41)		9.3 (32)		78.1 (134)		1.2 (36)
Iowa Falls		78.0 (33)		38.3 (13)		19.6 (38)		37.8 (6)		10.6 (25)		83.8 (87)		0.7 (23)
Jamaica		77.3 (36)		0.0 (150)		0.0 (136)		7.7 (141)		0.0 (131)		80.6 (118)		1.4 (42)
Jefferson		71.1 (67)		21.3 (87)		12.5 (94)		10.7 (129)		4.3 (92)		84.4 (84)		2.9 (92)
KR		67.2 (102)		29.6 (32)		25.6 (18)		12.9 (115)		7.4 (53)		83.5 (93)		1.0 (29)
Keokuk		51.7 (149)		11.9 (142)		9.8 (118)		11.6 (123)		2.4 (118)		79.7 (124)		5.7 (131)
Keosauqua		64.0 (123)		37.5 (15)		0.0 (136)		20.0 (56)		12.5 (14)		100.0 (1)		4.0 (116)
Knoxville		69.8 (82)		25.7 (59)		17.5 (55)		21.3 (50)		6.0 (63)		82.6 (104)		5.6 (130)
LCPH		68.8 (89)		28.6 (37)		12.5 (94)		15.0 (98)		0.0 (131)		70.0 (149)		2.8 (88)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Lamoni		68.0 (96)		25.0 (65)		0.0 (136)		0.0 (148)		8.3 (42)		90.2 (30)		1.0 (29)
Le Mars		84.7 (8)		28.0 (41)		17.9 (49)		24.4 (34)		6.7 (57)		82.1 (110)		0.5 (20)
Lenox		59.0 (136)		14.3 (137)		0.0 (136)		9.5 (135)		10.0 (28)		81.0 (116)		0.8 (26)
Leon		63.4 (126)		16.7 (122)		35.3 (5)		27.1 (22)		2.1 (123)		82.9 (103)		4.4 (121)
Logan		78.1 (31)		28.3 (39)		18.9 (43)		27.5 (20)		13.3 (12)		84.5 (81)		1.4 (42)
Luana		0.0 (155)		0.0 (150)		0.0 (136)		0.0 (148)		0.0 (131)		0.0 (158)		0.0 (1)
Manchester		42.4 (154)		10.0 (146)		6.7 (130)		16.3 (89)		0.0 (131)		96.0 (13)		0.0 (1)
Manning		66.7 (105)		0.0 (150)		0.0 (136)		100.0 (1)		0.0 (131)		100.0 (1)		0.0 (1)
Maquoketa		62.8 (128)		10.7 (144)		8.6 (126)		10.9 (128)		3.6 (108)		86.6 (61)		1.9 (56)
Marengo		66.0 (113)		15.8 (129)		11.1 (113)		10.3 (131)		0.0 (131)		100.0 (1)		1.9 (56)
Marion		67.7 (99)		25.5 (61)		17.6 (52)		17.9 (75)		4.6 (89)		83.0 (99)		2.3 (73)
Marshalltown		70.9 (70)		20.2 (96)		13.5 (85)		15.1 (97)		1.8 (126)		79.1 (128)		2.4 (79)
Mason City		66.7 (105)		20.8 (91)		14.6 (76)		14.2 (104)		6.9 (56)		87.0 (57)		2.1 (64)
Monticello		74.4 (51)		16.7 (122)		0.0 (136)		21.1 (52)		5.6 (70)		94.9 (18)		3.1 (96)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Mt Ayr		63.6 (125)		25.9 (55)		11.1 (113)		30.8 (13)		11.1 (23)		81.8 (112)		0.9 (28)
Mt Pleasan		70.0 (78)		25.8 (57)		11.7 (109)		23.1 (41)		6.7 (57)		84.5 (81)		2.9 (92)
Mtown Dtown		100.0 (1)		0.0 (150)		0.0 (136)		0.0 (148)		0.0 (131)		50.0 (156)		0.0 (1)
Murray		0.0 (155)		0.0 (150)		0.0 (136)		0.0 (148)		0.0 (131)		83.3 (95)		8.3 (141)
Muscatine		74.8 (48)		18.8 (111)		13.1 (91)		12.7 (116)		2.8 (114)		87.1 (55)		2.9 (92)
Nevada		50.0 (150)		7.1 (149)		33.3 (6)		9.1 (139)		0.0 (131)		87.7 (49)		2.8 (88)
New Hampto		75.9 (44)		25.0 (65)		26.7 (16)		20.0 (56)		9.1 (33)		85.6 (70)		1.9 (56)
Newton		70.5 (75)		21.1 (90)		15.0 (75)		17.2 (78)		2.3 (122)		83.6 (92)		2.5 (83)
North Towne		68.0 (96)		23.3 (77)		15.5 (72)		13.5 (111)		4.2 (93)		88.1 (45)		1.3 (39)
Northwood		73.5 (56)		26.3 (53)		26.7 (16)		28.6 (17)		5.3 (73)		84.7 (77)		0.7 (23)
Norwalk		74.5 (49)		25.0 (65)		12.0 (104)		24.1 (36)		8.3 (42)		83.0 (99)		2.0 (62)
Oakridge		50.0 (150)		20.0 (98)		9.1 (122)		0.0 (148)		0.0 (131)		100.0 (1)		0.0 (1)
Oelwein		71.4 (62)		22.1 (83)		5.7 (132)		24.4 (34)		5.2 (77)		84.7 (77)		20.1 (157)
Onawa		58.4 (138)		14.6 (136)		12.5 (94)		19.6 (64)		2.4 (118)		60.6 (155)		2.2 (67)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Orange Cty		84.2 (10)		36.5 (17)		28.6 (11)		35.1 (9)		5.8 (68)		90.2 (30)		1.5 (44)
Osage		71.8 (61)		24.5 (71)		16.7 (60)		31.1 (12)		8.2 (47)		92.0 (23)		1.1 (33)
Osceola		69.1 (86)		19.8 (102)		15.1 (74)		14.7 (102)		3.7 (105)		78.8 (132)		1.8 (53)
Oskaloosa		61.7 (130)		18.9 (110)		13.3 (87)		12.1 (118)		4.7 (85)		90.9 (27)		4.2 (119)
Ottumwa		73.5 (56)		23.7 (76)		17.6 (52)		18.1 (73)		5.2 (77)		85.2 (73)		3.2 (101)
Pella		75.4 (46)		30.0 (29)		15.6 (70)		25.0 (28)		10.0 (28)		82.2 (108)		2.4 (79)
Perry		84.3 (9)		17.7 (117)		16.7 (60)		9.3 (138)		1.8 (126)		83.3 (95)		2.4 (79)
Pocahontas		78.6 (29)		16.7 (122)		16.7 (60)		13.6 (107)		0.0 (131)		88.5 (42)		13.9 (148)
Pomeroy		80.0 (17)		0.0 (150)		0.0 (136)		0.0 (148)		0.0 (131)		100.0 (1)		0.0 (1)
Postville		79.1 (23)		42.3 (7)		20.0 (33)		20.9 (54)		15.6 (9)		87.6 (50)		5.8 (133)
Red Oak		69.3 (85)		18.3 (114)		17.0 (58)		14.8 (100)		4.2 (93)		80.3 (122)		1.9 (56)
Redfield		50.0 (150)		0.0 (150)		0.0 (136)		0.0 (148)		0.0 (131)		75.0 (142)		10.0 (143)
Refugee Cln		100.0 (1)		58.8 (3)		52.9 (1)		4.5 (146)		0.0 (131)		65.5 (153)		0.0 (1)
Rock Rapid		70.4 (76)		33.3 (21)		17.9 (49)		19.1 (66)		4.2 (93)		87.1 (55)		0.7 (23)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Rock Valle		81.9 (13)		29.2 (34)		30.6 (9)		23.5 (37)		13.8 (11)		83.0 (99)		2.3 (73)
Rockwell C		60.4 (135)		9.4 (147)		14.3 (78)		16.3 (89)		6.3 (60)		87.5 (51)		4.5 (124)
SDM		74.5 (49)		28.5 (38)		21.8 (29)		9.7 (134)		1.6 (129)		83.0 (99)		2.1 (64)
Sac City		53.5 (146)		25.0 (65)		11.8 (107)		20.0 (56)		5.0 (81)		90.1 (32)		6.4 (138)
Sheldon		79.0 (26)		25.5 (61)		19.0 (41)		19.4 (65)		3.9 (100)		91.7 (24)		14.3 (150)
Shenandoah		61.8 (129)		12.5 (140)		0.0 (136)		10.0 (132)		1.8 (126)		80.6 (118)		2.3 (73)
Sibley		81.1 (14)		15.4 (133)		19.0 (41)		11.1 (127)		7.7 (50)		87.4 (53)		19.3 (156)
Sidney		65.9 (115)		12.5 (140)		30.0 (10)		26.9 (23)		12.5 (14)		76.0 (140)		1.7 (50)
Sigourney		63.9 (124)		19.2 (107)		20.0 (33)		20.0 (56)		11.5 (20)		85.7 (67)		10.4 (144)
Sioux Cent		79.3 (21)		30.9 (27)		18.8 (45)		23.2 (39)		9.4 (31)		86.6 (61)		0.0 (1)
Sioux City		69.1 (86)		18.3 (114)		12.5 (94)		11.8 (121)		2.9 (111)		80.6 (118)		1.3 (39)
Southridge		77.0 (38)		18.4 (113)		13.2 (88)		4.9 (144)		0.0 (131)		81.5 (113)		3.1 (96)
Spencer		76.6 (40)		33.3 (21)		19.5 (39)		27.9 (19)		11.4 (21)		87.9 (48)		15.4 (151)
Spirit Lak		77.4 (34)		25.9 (55)		10.8 (115)		23.2 (39)		8.8 (37)		90.3 (29)		16.3 (153)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)						Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Storm Lake		64.8 (120)		19.4 (105)		14.4 (77)		14.0 (105)		2.1 (123)		84.5 (81)		4.5 (124)
Stuart		64.3 (122)		33.3 (21)		0.0 (136)		14.3 (103)		16.7 (7)		78.1 (134)		7.8 (140)
Tama		69.1 (86)		21.7 (84)		10.7 (116)		20.0 (56)		0.0 (131)		80.9 (117)		2.6 (85)
Tama/Settl		71.4 (62)		41.7 (9)		22.2 (25)		37.5 (7)		8.3 (42)		85.7 (67)		5.9 (135)
Tipton		50.0 (150)		29.2 (34)		16.7 (60)		13.6 (107)		16.7 (7)		77.8 (137)		6.3 (137)
Traer		77.3 (36)		38.5 (11)		16.7 (60)		44.4 (4)		23.1 (3)		89.5 (35)		1.3 (39)
UIHC		86.0 (7)		20.0 (98)		0.0 (136)		20.0 (56)		10.5 (26)		100.0 (1)		3.9 (113)
Vinton		68.8 (89)		28.9 (36)		12.8 (92)		18.9 (68)		11.1 (23)		74.2 (144)		3.6 (108)
WDM		77.4 (34)		33.8 (20)		25.5 (19)		11.2 (125)		3.8 (102)		82.4 (105)		2.2 (67)
Wapello		0.0 (155)		0.0 (150)		0.0 (136)		0.0 (148)		0.0 (131)		100.0 (1)		0.0 (1)
Washington		79.6 (19)		28.0 (41)		14.3 (78)		15.2 (96)		2.7 (115)		86.5 (63)		2.6 (85)
Waterloo		69.5 (84)		25.8 (57)		17.0 (58)		17.2 (78)		7.2 (54)		79.9 (123)		3.1 (96)
Waukee		79.1 (23)		27.4 (47)		23.5 (22)		15.4 (94)		5.3 (73)		83.8 (87)		1.2 (36)
Waukon		73.9 (53)		24.6 (70)		14.3 (78)		24.6 (32)		3.3 (109)		83.7 (91)		12.3 (146)

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

WIC PNSS & PedNSS

Table 7A

2017 - 2019 Pediatric Nutrition Surveillance (1)

Iowa

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic (2)

Children Aged < 5 Years (3)

Combined 3 Years

Comparison of Breastfeeding, TV Viewing, and Smoking in Household (4,5)

Clinic	Breastfeeding (6)					Exclusive Breastfeeding (7)				≤ 2 Hours TV Viewing/Day (8)		Smoking in Household	
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)	Number	% Breastfed At Least 3 Months (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% (Rank)	Number	% (Rank)
Waverly		69.9 (81)		38.2 (14)		21.9 (28)		44.0 (5)		21.8 (4)		88.6 (40)		1.1 (33)
Wbstr Ct		79.5 (20)		27.3 (48)		20.5 (31)		16.5 (86)		3.9 (100)		85.4 (71)		3.4 (105)
West Union		83.3 (11)		20.0 (98)		12.5 (94)		13.3 (113)		8.7 (39)		86.8 (58)		18.1 (154)
Williamsburg		100.0 (1)		50.0 (4)		50.0 (2)		50.0 (3)		0.0 (131)		50.0 (156)		0.0 (1)
Winterset		70.8 (72)		25.4 (63)		19.5 (39)		18.5 (70)		8.1 (48)		72.4 (148)		2.1 (64)
Iowa		**70.8**		**24.4**		**17.0**		**15.5**		**5.0**		**83.7**		**3.1**

(1) Reporting period is January 1 through December 31.
(2) TV Viewing and Smoking in Household data are not reported by all contributors.
(3) Analyses based on one record per child.
(4) Excludes records with unknown data or errors.
(5) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.
(7) The analysis includes infants who were ever breastfed and who turned that age during the reporting period by/on their date of visit.
(8) Children aged 2 years and older included in the analyses.
* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.