

# WIC PNSS & PedNSS

## Table 3F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Racial and Ethnic Distribution by Local Agency

Local Agency	White, Not Hispanic	Black, Not Hispanic	Hispanic	American Indian/Alaska Native	Asian/Pacific Islander	Multiple Races	All Other/Unknown
	%	%	%	%	%	%	%
Agency 31	43.1	20.3	25.2	0.3	7.2	4.0	0.0
Agency 33	87.1	3.5	8.2	0.0	1.2	0.0	0.0
Agency 34	59.3	24.7	8.5	0.0	2.6	5.0	0.0
Agency 35	76.1	11.1	7.3	0.5	3.0	2.1	0.0
Agency 37	83.2	1.6	14.4	0.0	0.8	0.0	0.0
Agency 38	62.5	3.8	24.6	1.0	6.5	1.5	0.0
Agency 39	57.9	1.2	38.2	0.8	1.2	0.8	0.0
Agency 41	77.3	3.8	14.9	0.2	2.2	1.6	0.0
Agency 42	77.6	3.1	18.4	0.0	1.0	0.0	0.0
Agency 43	62.7	20.3	8.0	0.6	5.0	3.5	0.0
Agency 45	69.1	8.8	18.7	0.0	2.1	1.2	0.0
Agency 46	77.8	5.9	11.7	0.0	3.6	1.0	0.0
Agency 47	56.0	3.5	33.2	0.3	6.7	0.3	0.0
Agency 48	71.7	2.8	23.7	0.3	1.2	0.3	0.0
Agency 49	44.4	10.0	36.8	3.8	2.5	2.5	0.0
Agency 50	70.0	5.3	22.1	0.3	0.3	2.0	0.0
Agency 51	40.3	34.8	20.5	0.2	2.0	2.2	0.0
Agency 52	79.3	2.3	14.3	1.0	0.8	2.3	0.0
Agency 53	90.0	2.4	5.2	0.0	0.9	1.4	0.0
Agency 54	58.7	23.2	11.3	0.5	1.1	5.3	0.0
<b>Iowa</b>	<b>61.3</b>	<b>13.2</b>	<b>18.7</b>	<b>0.5</b>	<b>3.7</b>	<b>2.6</b>	<b>0.0</b>
<b>National (Prior Year)</b>	<b>53.3</b>	<b>21.1</b>	<b>16.9</b>	<b>2.2</b>	<b>2.9</b>	<b>3.1</b>	<b>0.5</b>

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Age Distribution by Local Agency

Local Agency	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Agency 31	0.0	2.5	6.7	57.6	30.1	3.1	0.0
Agency 33	0.0	1.2	3.5	65.5	27.5	2.3	0.0
Agency 34	0.0	1.0	6.3	61.1	29.1	2.5	0.0
Agency 35	0.0	2.5	7.8	62.1	26.5	1.0	0.2
Agency 37	0.0	2.4	8.0	63.2	24.4	2.0	0.0
Agency 38	0.0	1.7	5.2	60.7	31.2	1.2	0.0
Agency 39	0.0	3.9	7.9	55.1	29.1	3.5	0.4
Agency 41	0.0	0.8	4.0	64.0	29.8	1.4	0.0
Agency 42	0.0	2.0	5.1	64.3	27.0	1.5	0.0
Agency 43	0.2	1.7	4.2	63.9	28.7	1.4	0.0
Agency 45	0.0	4.0	8.2	59.4	26.8	1.7	0.0
Agency 46	0.0	1.9	7.5	65.4	23.5	1.5	0.2
Agency 47	0.0	2.9	5.2	62.1	27.1	2.6	0.0
Agency 48	0.0	2.5	4.4	62.9	29.3	0.6	0.3
Agency 49	0.2	2.2	6.5	57.1	32.4	1.7	0.0
Agency 50	0.3	3.9	7.0	59.7	27.2	2.0	0.0
Agency 51	0.0	0.7	2.9	56.2	37.4	2.9	0.0
Agency 52	0.0	3.1	5.7	64.1	26.2	1.0	0.0
Agency 53	0.0	2.8	5.2	61.6	29.4	0.9	0.0
Agency 54	0.3	3.2	7.8	61.6	25.6	1.5	0.0
<b>Iowa</b>	<b>0.1</b>	<b>2.3</b>	<b>6.1</b>	<b>60.7</b>	<b>28.8</b>	<b>2.0</b>	<b>0.0</b>
<b>National (Prior Year)</b>	<b>0.1</b>	<b>2.9</b>	<b>8.1</b>	<b>62.2</b>	<b>24.7</b>	<b>1.8</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31

(2) Excludes data with errors

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Maternal Health Indicators by Local Agency

<b>Comparison of Maternal Health Indicators (2,3)</b>														
Local Agency	Prepregnancy BMI			Weight Gain			Anemia (Low Hb/Hct)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	(4)			(5)			(6)							
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agency 31		3.8 (13)	57.0 (1)	23.3 (13)	43.6 (6)		2.0 (2)		3.7 (2)		100.0 (1)		*	
Agency 33		2.2 (3)	71.1 (20)	16.9 (1)	53.8 (20)		*		*		*		*	
Agency 34		2.4 (4)	62.2 (12)	22.3 (10)	48.3 (15)		31.4 (7)		38.7 (5)		*		*	
Agency 35		4.7 (18)	64.6 (18)	21.2 (6)	51.8 (19)		45.3 (11)		*		*		*	
Agency 37		4.6 (17)	60.1 (4)	27.5 (17)	47.2 (13)		*		*		*		*	
Agency 38		3.4 (10)	61.4 (9)	22.9 (12)	42.4 (3)		17.7 (6)		21.3 (3)		*		*	
Agency 39		4.5 (16)	59.7 (2)	31.8 (20)	42.7 (4)		*		*		*		*	
Agency 41		3.8 (13)	66.4 (19)	18.6 (4)	48.1 (14)		*		*		*		*	
Agency 42		1.9 (2)	62.6 (14)	20.3 (5)	44.6 (9)		*		*		*		*	
Agency 43		3.6 (12)	61.7 (10)	22.4 (11)	45.3 (11)		8.6 (5)		*		*		*	
Agency 45		3.2 (8)	61.1 (7)	18.5 (3)	49.1 (18)		3.3 (3)		*		*		*	
Agency 46		3.3 (9)	62.1 (11)	24.0 (15)	46.1 (12)		33.8 (8)		*		*		*	
Agency 47		3.1 (7)	61.2 (8)	23.8 (14)	42.7 (4)		*		*		*		*	
Agency 48		4.3 (15)	63.6 (16)	28.0 (19)	42.3 (2)		*		*		*		*	
Agency 49		3.5 (11)	63.2 (15)	21.6 (7)	41.9 (1)		1.7 (1)		3.5 (1)		*		*	
Agency 50		1.7 (1)	60.8 (6)	25.5 (16)	45.0 (10)		*		*		*		*	
Agency 51		2.8 (5)	59.7 (2)	22.2 (9)	44.0 (7)		35.6 (9)		26.1 (4)		*		*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Maternal Health Indicators by Local Agency

<b>Comparison of Maternal Health Indicators (2,3)</b>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agency 52	5.1 (20)	60.6 (5)		21.7 (8)	48.7 (16)		7.3 (4)		*		*		*	
Agency 53	4.7 (18)	64.2 (17)		27.5 (17)	44.4 (8)		*		*		*		*	
Agency 54	3.0 (6)	62.2 (12)		18.4 (2)	48.8 (17)		36.1 (10)		*		*		*	
<b>Iowa</b>	<b>3.5</b>	<b>61.2</b>		<b>22.5</b>	<b>45.7</b>		<b>14.4</b>		<b>14.4</b>		<b>100.0</b>		<b>100.0</b>	
<b>National (Prior Year)</b>	<b>120,741</b>	<b>4.0</b>	<b>59.6</b>	<b>97,554</b>	<b>22.4</b>	<b>47.7</b>	<b>26,239</b>	<b>27.4</b>	<b>53,856</b>	<b>17.5</b>	<b>104,317</b>	<b>9.9</b>	<b>103,483</b>	<b>9.3</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Maternal Behavioral Indicators by Local Agency

<b>Comparison of Maternal Behavioral Indicators (2,3)</b>									
<b>Local Agency</b>	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy (4)		During Pregnancy (5)	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31		30.6 (11)	11.6 (11)		1.3 (11)		12.1 (13)		86.6 (12)
Agency 33		30.6 (11)	6.5 (2)		0.0 (19)		*		*
Agency 34		25.7 (16)	10.9 (10)		0.8 (14)		16.2 (10)		89.4 (6)
Agency 35		36.5 (4)	12.9 (12)		2.8 (2)		15.6 (11)		88.9 (7)
Agency 37		36.7 (3)	7.5 (6)		3.1 (1)		*		*
Agency 38		37.7 (2)	10.8 (9)		1.8 (8)		18.4 (8)		87.3 (9)
Agency 39		27.7 (14)	10.4 (8)		2.7 (3)		*		*
Agency 41		38.6 (1)	19.4 (16)		1.2 (12)		22.3 (3)		86.8 (11)
Agency 42		25.7 (16)	14.5 (13)		0.0 (19)		26.0 (2)		89.7 (4)
Agency 43		31.7 (9)	24.3 (20)		2.3 (6)		22.0 (4)		86.5 (13)
Agency 45		32.9 (7)	7.3 (5)		1.5 (10)		18.8 (6)		91.3 (3)
Agency 46		24.9 (19)	20.9 (18)		2.6 (4)		*		*
Agency 47		25.2 (18)	7.2 (4)		0.3 (17)		49.8 (1)		93.3 (2)
Agency 48		31.2 (10)	5.1 (1)		0.3 (17)		*		*
Agency 49		28.5 (13)	9.8 (7)		0.7 (16)		*		93.4 (1)
Agency 50		32.8 (8)	20.0 (17)		2.5 (5)		16.7 (9)		82.4 (14)
Agency 51		16.0 (20)	21.0 (19)		1.7 (9)		18.6 (7)		87.4 (8)
Agency 52		33.4 (6)	7.1 (3)		0.8 (14)		20.5 (5)		87.0 (10)
Agency 53		25.9 (15)	15.6 (15)		1.2 (12)		13.2 (12)		89.6 (5)
Agency 54		35.2 (5)	14.5 (13)		1.9 (7)		*		*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Maternal Behavioral Indicators by Local Agency

<b>Iowa</b>		<b>30.9</b>	<b>13.1</b>		<b>1.5</b>		<b>19.6</b>		<b>88.2</b>
<b>National (Prior Year)</b>	<b>109,069</b>	<b>69.2</b>	<b>7.4</b>	<b>136,111</b>	<b>23.4</b>	<b>99,278</b>	<b>22.2</b>	<b>98,734</b>	<b>88.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Smoking Indicators by Local Agency

Comparison of Smoking Indicators (2,3)										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31		22.4 (3)		13.3 (2)		23.9 (4)		5.2 (6)		1.6 (9)
Agency 33		39.1 (20)		20.9 (17)		*		6.3 (9)		1.9 (10)
Agency 34		27.3 (8)		14.5 (5)		40.6 (1)		6.4 (10)		1.4 (8)
Agency 35		24.1 (6)		15.4 (7)		21.1 (7)		1.6 (1)		0.8 (4)
Agency 37		36.0 (18)		26.1 (20)		*		10.0 (16)		0.7 (3)
Agency 38		22.7 (4)		16.0 (9)		22.3 (6)		7.8 (12)		2.1 (12)
Agency 39		21.7 (2)		15.9 (8)		*		3.3 (2)		0.8 (4)
Agency 41		35.6 (17)		23.1 (19)		23.1 (5)		6.9 (11)		1.3 (7)
Agency 42		31.1 (12)		17.3 (10)		*		13.8 (20)		12.7 (20)
Agency 43		34.1 (15)		20.3 (14)		18.0 (10)		9.7 (15)		3.1 (15)
Agency 45		32.5 (13)		17.3 (10)		34.0 (3)		13.3 (19)		3.1 (15)
Agency 46		29.7 (9)		14.3 (4)		*		11.8 (17)		3.3 (18)
Agency 47		23.6 (5)		13.9 (3)		*		8.1 (13)		7.8 (19)
Agency 48		30.2 (10)		18.2 (12)		*		3.6 (3)		0.0 (1)
Agency 49		25.3 (7)		14.5 (5)		*		6.1 (8)		2.9 (14)
Agency 50		36.4 (19)		20.4 (15)		38.9 (2)		4.9 (5)		1.9 (10)
Agency 51		16.8 (1)		7.8 (1)		*		3.7 (4)		0.0 (1)
Agency 52		32.7 (14)		22.8 (18)		20.4 (9)		5.4 (7)		0.9 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Smoking Indicators by Local Agency

<b>Comparison of Smoking Indicators (2,3)</b>										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
Agency 53		34.9 (16)		20.7 (16)		*		9.3 (14)		2.7 (13)
Agency 54		30.7 (11)		18.4 (13)		20.7 (8)		13.0 (18)		3.2 (17)
<b>Iowa</b>		<b>28.1</b>		<b>16.7</b>		<b>26.3</b>		<b>7.3</b>		<b>2.3</b>
<b>National (Prior Year)</b>	<b>115,645</b>	<b>28.4</b>	<b>127,715</b>	<b>14.9</b>	<b>28,227</b>	<b>36.2</b>	<b>104,358</b>	<b>10.5</b>	<b>128,247</b>	<b>3.4</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Infant Health Indicators by Local Agency

<b>Comparison of Infant Health Indicators (2,3)</b>									
Local Agency	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31	7.3 (11)	8.0 (12)		9.7 (9)		2.6 (7)		79.2 (3)	
Agency 33	11.0 (19)	6.1 (1)		13.3 (19)		3.6 (16)		68.3 (14)	
Agency 34	6.4 (6)	9.3 (19)		8.9 (4)		2.9 (11)		68.5 (13)	
Agency 35	8.8 (17)	8.3 (14)		8.2 (2)		2.8 (9)		59.8 (20)	
Agency 37	6.5 (7)	7.3 (9)		11.3 (15)		1.5 (3)		66.8 (15)	
Agency 38	8.2 (15)	9.1 (18)		11.2 (13)		3.2 (13)		73.7 (7)	
Agency 39	6.3 (5)	6.3 (2)		12.4 (17)		3.7 (18)		80.3 (2)	
Agency 41	5.7 (3)	9.5 (20)		10.8 (12)		2.6 (7)		66.7 (16)	
Agency 42	3.2 (1)	6.5 (3)		6.8 (1)		1.3 (1)		78.4 (4)	
Agency 43	7.7 (13)	7.6 (11)		9.4 (8)		3.6 (16)		69.8 (10)	
Agency 45	7.2 (9)	8.7 (16)		9.2 (6)		3.5 (15)		65.9 (17)	
Agency 46	7.2 (9)	8.1 (13)		9.9 (10)		2.8 (9)		73.5 (8)	
Agency 47	6.5 (7)	7.1 (6)		9.2 (6)		2.3 (6)		69.8 (10)	
Agency 48	8.1 (14)	7.5 (10)		13.2 (18)		1.5 (3)		63.8 (19)	
Agency 49	8.3 (16)	8.3 (14)		10.1 (11)		3.0 (12)		71.6 (9)	
Agency 50	5.4 (2)	7.2 (8)		9.0 (5)		2.1 (5)		74.3 (6)	
Agency 51	12.0 (20)	7.1 (6)		11.3 (15)		4.2 (19)		85.5 (1)	
Agency 52	7.5 (12)	6.9 (5)		13.3 (19)		1.3 (1)		75.8 (5)	
Agency 53	6.0 (4)	9.0 (17)		8.3 (3)		3.2 (13)		69.7 (12)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8F

### 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Infant Health Indicators by Local Agency

Comparison of Infant Health Indicators (2,3)									
Local Agency	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 54		9.8 (18)	6.7 (4)		11.2 (13)		5.4 (20)		64.0 (18)
<b>Iowa</b>		<b>7.6</b>	<b>8.0</b>		<b>10.2</b>		<b>2.9</b>		<b>72.0</b>
<b>National (Prior Year)</b>	<b>125,347</b>	<b>9.3</b>	<b>6.9</b>	<b>125,751</b>	<b>11.6</b>	<b>106,512</b>	<b>3.9</b>	<b>126,356</b>	<b>53.2</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* National includes previous year's data for all participating States and ITOs.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

#### Comparison of Racial and Ethnic Distribution by Local Agency Combined 3 Years

Local Agency	White, Not Hispanic  %	Black, Not Hispanic  %	Hispanic  %	American Indian/Alaska Native  %	Asian/Pacific Islander  %	Multiple Races  %	All Other/Unknown  %
Agency 31	45.5	17.6	24.3	0.3	7.7	4.6	0.0
Agency 33	79.2	3.5	15.7	0.0	1.2	0.4	0.0
Agency 34	61.3	22.9	7.9	0.2	2.2	4.3	1.1
Agency 35	77.2	10.9	7.0	0.2	2.5	2.2	0.0
Agency 36	*	*	*	*	*	*	*
Agency 37	85.8	0.8	12.4	0.2	0.6	0.2	0.0
Agency 38	61.2	4.0	25.0	1.6	6.8	1.4	0.1
Agency 39	63.0	0.7	34.1	0.7	0.8	0.9	0.0
Agency 41	79.3	4.1	12.9	0.2	1.9	1.5	0.0
Agency 42	78.6	4.1	15.4	0.0	0.6	1.3	0.0
Agency 43	63.8	19.1	8.6	0.3	4.2	3.7	0.3
Agency 45	70.8	7.9	17.8	0.0	2.3	1.1	0.0
Agency 46	76.4	6.5	13.2	0.1	2.4	1.4	0.1
Agency 47	54.6	2.9	33.5	0.2	7.8	1.1	0.0
Agency 48	70.5	1.5	25.1	0.3	2.3	0.3	0.0
Agency 49	44.3	10.3	36.6	3.2	3.2	2.4	0.0
Agency 50	70.0	4.2	23.9	0.2	0.3	1.4	0.0
Agency 51	41.6	34.9	18.6	0.2	2.8	1.9	0.0
Agency 52	78.1	3.3	15.0	0.5	0.5	2.5	0.0
Agency 53	90.8	2.3	5.2	0.3	0.8	0.6	0.1
Agency 54	60.9	22.8	10.6	0.3	1.2	4.3	0.0
<b>Iowa</b>	<b>62.3</b>	<b>12.3</b>	<b>18.4</b>	<b>0.5</b>	<b>3.7</b>	<b>2.6</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

### Comparison of Age Distribution by Local Agency Combined 3 Years

Local Agency	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Agency 31	0.1	2.7	6.7	58.2	29.6	2.7	0.0
Agency 33	0.0	1.7	5.5	62.6	28.1	2.1	0.0
Agency 34	0.1	1.6	6.3	62.9	27.4	1.6	0.0
Agency 35	0.1	2.9	8.2	64.0	23.5	1.0	0.2
Agency 36	*	*	*	*	*	*	*
Agency 37	0.0	2.3	7.1	65.1	23.5	2.0	0.1
Agency 38	0.0	2.1	6.3	61.5	28.2	1.8	0.1
Agency 39	0.1	2.4	8.2	58.3	28.6	2.1	0.3
Agency 41	0.1	1.3	6.7	64.6	25.7	1.6	0.0
Agency 42	0.0	1.7	5.8	65.0	26.3	1.1	0.0
Agency 43	0.3	1.8	5.1	64.5	27.2	1.2	0.0
Agency 45	0.0	3.8	9.0	63.6	22.1	1.5	0.0
Agency 46	0.1	2.6	7.6	64.5	24.0	1.2	0.1
Agency 47	0.0	2.1	6.3	65.8	24.1	1.8	0.0
Agency 48	0.0	2.2	7.9	61.9	26.7	1.3	0.1
Agency 49	0.0	2.1	6.7	61.8	27.5	1.9	0.0
Agency 50	0.3	3.9	8.1	61.2	24.9	1.7	0.0
Agency 51	0.0	0.9	4.7	54.8	36.4	3.1	0.1
Agency 52	0.1	3.2	7.2	66.3	22.1	1.2	0.0
Agency 53	0.0	2.3	4.9	65.7	26.1	1.0	0.0
Agency 54	0.1	3.5	8.3	63.4	23.7	1.0	0.0
<b>Iowa</b>	<b>0.1</b>	<b>2.4</b>	<b>6.8</b>	<b>62.1</b>	<b>26.8</b>	<b>1.8</b>	<b>0.0</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

#### Comparison of Maternal Health Indicators by Local Agency

#### Combined 3 Years

<b>Comparison of Maternal Health Indicators</b> <sup>(2,3)</sup>														
Local Agency	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
Agency 31	4.1 (14)	57.1 (2)	25.1 (19)	42.1 (1)	2.6 (3)	3.4 (2)	77.4 (18)	62.5 (16)						
Agency 33	3.4 (7)	64.1 (20)	21.4 (7)	47.5 (13)	27.8 (14)	22.0 (14)	43.8 (10)	42.2 (14)						
Agency 34	3.3 (6)	61.4 (10)	21.0 (5)	49.1 (18)	34.5 (17)	30.9 (20)	57.6 (15)	40.1 (12)						
Agency 35	4.1 (14)	59.3 (3)	20.4 (2)	50.5 (20)	39.7 (20)	29.6 (19)	47.0 (12)	22.9 (3)						
Agency 36	*	*	*	*	*	*	*	*						
Agency 37	3.9 (12)	62.3 (14)	22.5 (11)	49.0 (17)	30.6 (15)	17.2 (11)	31.2 (5)	23.0 (4)						
Agency 38	4.3 (16)	59.6 (4)	23.3 (14)	43.1 (3)	19.6 (12)	19.6 (12)	70.1 (17)	28.0 (7)						
Agency 39	2.8 (4)	60.4 (9)	27.2 (20)	44.3 (5)	0.9 (1)	0.9 (1)	22.4 (1)	19.7 (1)						
Agency 41	3.4 (7)	63.7 (19)	21.0 (5)	48.2 (14)	4.9 (5)	9.1 (9)	44.7 (11)	29.6 (9)						
Agency 42	2.0 (1)	62.5 (15)	24.4 (18)	45.0 (7)	32.7 (16)	24.7 (17)	41.5 (9)	*						
Agency 43	3.0 (5)	62.5 (15)	22.7 (12)	45.9 (9)	10.5 (10)	6.6 (6)	39.7 (8)	33.5 (11)						
Agency 45	4.5 (17)	59.6 (4)	20.2 (1)	48.2 (14)	4.1 (4)	4.5 (4)	54.4 (14)	40.8 (13)						
Agency 46	3.9 (12)	61.9 (13)	23.9 (15)	46.5 (11)	22.4 (13)	22.9 (16)	*	*						
Agency 47	2.4 (2)	62.5 (15)	21.9 (10)	45.7 (8)	2.1 (2)	4.1 (3)	33.2 (6)	20.6 (2)						
Agency 48	4.5 (17)	61.4 (10)	23.1 (13)	45.9 (9)	6.3 (8)	9.5 (10)	25.7 (3)	26.2 (5)						
Agency 49	3.5 (9)	60.3 (8)	21.4 (7)	44.3 (5)	5.6 (7)	8.4 (8)	51.9 (13)	30.3 (10)						
Agency 50	2.7 (3)	61.4 (10)	24.1 (16)	43.7 (4)	16.3 (11)	20.0 (13)	35.2 (7)	26.2 (5)						

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

#### Comparison of Maternal Health Indicators by Local Agency

#### Combined 3 Years

<b>Comparison of Maternal Health Indicators <sup>(2,3)</sup></b>																
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy			
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
Agency 51	3.5	(9)	56.3	(1)	24.2	(17)	42.5	(2)	36.8	(18)	25.6	(18)	65.0	(16)	*	
Agency 52	3.7	(11)	59.7	(6)	20.4	(2)	48.4	(16)	7.9	(9)	4.9	(5)	28.9	(4)	29.4	(8)
Agency 53	4.6	(19)	62.7	(18)	20.7	(4)	50.1	(19)	5.5	(6)	8.1	(7)	*		*	
Agency 54	4.9	(20)	60.0	(7)	21.8	(9)	47.0	(12)	39.4	(19)	22.6	(15)	23.5	(2)	42.7	(15)
<b>Iowa</b>	<b>3.8</b>		<b>60.1</b>		<b>22.7</b>		<b>45.7</b>		<b>15.5</b>		<b>15.4</b>		<b>48.2</b>		<b>35.0</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre-pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre-pregnancy (ideal weight gain = 15 to 25 pounds), obese pre-pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

#### Comparison of Maternal Behavioral Indicators by Local Agency

#### Combined 3 Years

<b>Comparison of Maternal Behavioral Indicators <sup>(2,3)</sup></b>									
<b>Local Agency</b>	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31		29.7 (12)	14.6 (3)		6.8 (19)		15.1 (20)		70.0 (18)
Agency 33		28.4 (13)	21.3 (12)		9.4 (10)		20.1 (10)		64.6 (20)
Agency 34		23.7 (17)	16.8 (6)		9.0 (12)		15.7 (18)		80.9 (10)
Agency 35		32.0 (8)	20.0 (10)		10.4 (8)		17.9 (13)		88.8 (5)
Agency 36		*	*		*		*		*
Agency 37		33.5 (6)	27.2 (18)		12.3 (2)		23.1 (6)		91.1 (1)
Agency 38		34.4 (4)	16.0 (5)		11.2 (7)		17.0 (16)		76.8 (15)
Agency 39		38.5 (1)	11.7 (1)		8.0 (16)		24.8 (5)		76.3 (16)
Agency 41		32.1 (7)	31.1 (20)		12.3 (2)		22.8 (7)		85.4 (7)
Agency 42		27.1 (15)	19.9 (9)		8.6 (13)		26.3 (4)		89.7 (4)
Agency 43		36.3 (2)	25.8 (17)		12.1 (4)		18.2 (12)		86.4 (6)
Agency 45		31.1 (10)	13.9 (2)		9.1 (11)		15.9 (17)		90.1 (3)
Agency 46		30.2 (11)	19.8 (8)		11.4 (5)		22.0 (8)		79.9 (13)
Agency 47		21.7 (19)	22.3 (13)		6.4 (20)		36.7 (1)		90.3 (2)
Agency 48		36.2 (3)	16.8 (6)		7.1 (18)		31.6 (2)		73.4 (17)
Agency 49		26.5 (16)	24.6 (15)		8.6 (13)		22.0 (8)		84.5 (8)
Agency 50		27.4 (14)	30.9 (19)		12.4 (1)		17.2 (15)		77.1 (14)
Agency 51		20.3 (20)	20.4 (11)		7.6 (17)		18.6 (11)		80.0 (12)
Agency 52		34.2 (5)	23.4 (14)		9.6 (9)		17.8 (14)		83.3 (9)
Agency 53		23.6 (18)	15.5 (4)		8.5 (15)		29.0 (3)		80.2 (11)
Agency 54		31.8 (9)	24.7 (16)		11.4 (5)		15.7 (18)		67.5 (19)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6F

### 2016 - 2018 Pregnancy Nutrition Surveillance (1)

#### Iowa

#### Comparison of Maternal Behavioral Indicators by Local Agency

#### Combined 3 Years

Iowa	30.2	20.0	9.3	19.7	80.5
------	------	------	-----	------	------

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

#### Comparison of Smoking Indicators by Local Agency

#### Combined 3 Years

<b>Comparison of Smoking Indicators</b> <sup>(2,3)</sup>										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes <sup>(4)</sup>		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31		25.8 (5)		16.7 (7)		23.5 (17)		6.8 (8)		3.1 (13)
Agency 33		37.6 (18)		20.4 (14)		33.3 (5)		7.5 (9)		1.3 (5)
Agency 34		33.7 (12)		18.0 (8)		43.2 (1)		8.4 (10)		1.7 (7)
Agency 35		24.1 (3)		15.3 (4)		25.0 (14)		2.2 (1)		0.8 (2)
Agency 36		*		*		*		*		*
Agency 37		40.0 (20)		29.7 (20)		19.5 (19)		12.7 (16)		4.6 (17)
Agency 38		26.3 (6)		16.3 (5)		30.9 (8)		8.6 (11)		2.1 (10)
Agency 39		25.0 (4)		14.1 (3)		23.6 (16)		3.1 (2)		0.6 (1)
Agency 41		35.3 (15)		22.1 (17)		25.1 (13)		6.3 (5)		1.1 (3)
Agency 42		33.5 (11)		18.8 (10)		32.9 (6)		15.7 (20)		14.3 (20)
Agency 43		35.3 (15)		20.4 (14)		25.2 (12)		9.0 (13)		1.8 (9)
Agency 45		36.8 (17)		19.4 (12)		35.4 (4)		13.8 (18)		4.7 (18)
Agency 46		31.4 (9)		20.8 (16)		17.3 (20)		14.6 (19)		3.3 (14)
Agency 47		23.9 (2)		12.9 (2)		39.1 (2)		8.6 (11)		6.8 (19)
Agency 48		33.0 (10)		19.6 (13)		32.4 (7)		6.7 (7)		1.2 (4)
Agency 49		29.2 (7)		16.4 (6)		28.5 (10)		6.3 (5)		2.4 (11)
Agency 50		31.3 (8)		19.1 (11)		36.0 (3)		10.6 (14)		2.5 (12)
Agency 51		17.0 (1)		9.5 (1)		25.4 (11)		4.4 (3)		1.3 (5)
Agency 52		37.8 (19)		23.7 (18)		21.7 (18)		5.7 (4)		1.7 (7)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

#### Comparison of Smoking Indicators by Local Agency

#### Combined 3 Years

Comparison of Smoking Indicators <sup>(2,3)</sup>										
Local Agency	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes <sup>(4)</sup>	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (rank)	Number	% (rank)		Number	% (rank)	Number	% (rank)	
Agency 53		34.4 (14)		24.9 (19)		23.8 (15)		12.5 (15)		4.0 (16)
Agency 54		34.0 (13)		18.4 (9)		29.3 (9)		12.9 (17)		3.4 (15)
<b>Iowa</b>		<b>30.5</b>		<b>18.3</b>		<b>29.3</b>		<b>8.4</b>		<b>2.8</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

### Comparison of Infant Health Indicators by Local Agency

#### Combined 3 Years

Comparison of Infant Health Indicators <sup>(2,3)</sup>									
Local Agency	Birthweight			Preterm		Full Term LBW		Ever Breastfed	
	Low (4)		High (5)	(6)		(7)			
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)
Agency 31		8.0 (16)	8.4 (11)		10.1 (12)		3.1 (13)		78.7 (2)
Agency 33		9.1 (18)	9.5 (19)		11.0 (17)		4.0 (18)		67.1 (16)
Agency 34		7.9 (15)	8.2 (9)		10.4 (15)		2.6 (9)		68.1 (15)
Agency 35		8.0 (16)	9.4 (18)		8.3 (3)		3.0 (11)		60.4 (20)
Agency 36		*	*		*		*		*
Agency 37		6.2 (4)	6.5 (1)		9.0 (4)		3.0 (11)		63.9 (17)
Agency 38		7.5 (12)	9.2 (16)		9.6 (7)		3.2 (15)		70.0 (11)
Agency 39		7.0 (10)	8.5 (14)		11.4 (19)		2.6 (9)		78.4 (3)
Agency 41		6.3 (5)	9.8 (20)		9.8 (11)		2.3 (8)		69.3 (14)
Agency 42		5.4 (1)	8.7 (15)		8.0 (1)		2.0 (4)		77.0 (4)
Agency 43		7.2 (11)	8.1 (8)		9.4 (6)		3.1 (13)		69.7 (12)
Agency 45		7.6 (13)	8.4 (11)		9.3 (5)		3.5 (16)		63.9 (17)
Agency 46		6.7 (6)	7.9 (5)		10.8 (16)		2.2 (7)		69.4 (13)
Agency 47		5.6 (2)	7.7 (4)		9.7 (8)		2.1 (6)		70.2 (10)
Agency 48		6.1 (3)	8.4 (11)		10.3 (14)		1.5 (1)		70.4 (9)
Agency 49		6.8 (8)	8.0 (6)		9.7 (8)		1.9 (3)		71.8 (8)
Agency 50		6.8 (8)	8.2 (9)		8.1 (2)		3.6 (17)		76.7 (5)
Agency 51		11.4 (20)	6.8 (2)		11.3 (18)		4.0 (18)		81.2 (1)
Agency 52		7.6 (13)	8.0 (6)		12.3 (20)		1.8 (2)		74.9 (6)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8F

### 2016 - 2018 Pregnancy Nutrition Surveillance <sup>(1)</sup>

#### Iowa

#### Comparison of Infant Health Indicators by Local Agency

#### Combined 3 Years

<b>Comparison of Infant Health Indicators</b> <sup>(2,3)</sup>										
<b>Local Agency</b>	Birthweight			Preterm		Full Term LBW		Ever Breastfed		
	Low (4)		High (5)	(6)		(7)				
	Number	% (rank)	% (rank)	Number	% (rank)	Number	% (rank)	Number	% (rank)	
Agency 53	6.7	(6)	9.3	(17)	9.7	(8)	2.0	(4)	72.5	(7)
Agency 54	9.6	(19)	7.2	(3)	10.2	(13)	5.0	(20)	62.6	(19)
<b>Iowa</b>	<b>7.6</b>		<b>8.3</b>		<b>10.0</b>		<b>2.9</b>		<b>67.6</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one Local Agency's rate to other Local Agencies. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.