# WIC PNSS & PedNSS

Run Date: 2/7/2020

## Table 3A



## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Racial and Ethnic Distribution by Clinic

| **Clinic** | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Ackley | * | * | * | * | * | * | * |
| Adel | * | * | * | * | * | * | * |
| Albia | * | * | * | * | * | * | * |
| Algona | * | * | * | * | * | * | * |
| Allen WHC | * | * | * | * | * | * | * |
| Allison | * | * | * | * | * | * | * |
| Altoona | * | * | * | * | * | * | * |
| Ames | 75.5 | 7.3 | 5.5 | 0.0 | 10.0 | 1.8 | 0.0 |
| Anamosa | * | * | * | * | * | * | * |
| Ankeny | * | * | * | * | * | * | * |
| Atlantic | * | * | * | * | * | * | * |
| Audubon | * | * | * | * | * | * | * |
| Avoca | * | * | * | * | * | * | * |
| Bedford | * | * | * | * | * | * | * |
| Belle Plaine | * | * | * | * | * | * | * |
| Bloomfield | * | * | * | * | * | * | * |
| Boone | * | * | * | * | * | * | * |
| Brdlwns RP | 38.4 | 25.7 | 21.7 | 0.5 | 8.8 | 4.9 | 0.0 |
| Brooklyn | * | * | * | * | * | * | * |
| Burlington | 74.1 | 16.8 | 6.6 | 0.0 | 0.0 | 2.5 | 0.0 |
| CHC | 58.2 | 20.0 | 15.6 | 0.0 | 1.3 | 4.9 | 0.0 |
| CR Urban | 50.0 | 34.0 | 8.6 | 0.0 | 1.0 | 6.4 | 0.0 |
| Carlisle | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS



## Table 3A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Racial and Ethnic Distribution by Clinic

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Carroll | * | * | * | * | * | * | * |
| Cedar Falls | * | * | * | * | * | * | * |
| Centervill | * | * | * | * | * | * | * |
| Chariton | * | * | * | * | * | * | * |
| Charles City | * | * | * | * | * | * | * |
| Cherokee | * | * | * | * | * | * | * |
| Clarinda | * | * | * | * | * | * | * |
| Clarion | * | * | * | * | * | * | * |
| Clinton | 77.5 | 11.0 | 8.7 | 0.0 | 0.6 | 2.3 | 0.0 |
| Colfax | * | * | * | * | * | * | * |
| Columbus J | * | * | * | * | * | * | * |
| Corning | * | * | * | * | * | * | * |
| Corydon | * | * | * | * | * | * | * |
| Council Blfs | 79.2 | 2.3 | 14.4 | 1.0 | 0.8 | 2.3 | 0.0 |
| Cresco | * | * | * | * | * | * | * |
| Creston | * | * | * | * | * | * | * |
| Dallas Ctr | * | * | * | * | * | * | * |
| Decorah | * | * | * | * | * | * | * |
| Denison | 36.8 | 6.8 | 52.1 | 0.9 | 3.4 | 0.0 | 0.0 |
| Dewitt | * | * | * | * | * | * | * |
| Dubuque | 74.2 | 12.7 | 6.6 | 0.8 | 3.5 | 2.3 | 0.0 |
| Dyersville | * | * | * | * | * | * | * |
| Eagle Grove | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Racial and Ethnic Distribution by Clinic

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Edgerton | 59.0 | 24.8 | 9.1 | 0.7 | 0.9 | 5.5 | 0.0 |
| Eldora | * | * | * | * | * | * | * |
| Elkader | * | * | * | * | * | * | * |
| Emmetsburg | * | * | * | * | * | * | * |
| Esthervill | * | * | * | * | * | * | * |
| Fairfiel | * | * | * | * | * | * | * |
| Forest City | * | * | * | * | * | * | * |
| Fort Dodge | 79.7 | 8.4 | 8.4 | 0.5 | 0.0 | 3.0 | 0.0 |
| Ft Madison | * | * | * | * | * | * | * |
| Garner | * | * | * | * | * | * | * |
| Glenwood | * | * | * | * | * | * | * |
| Grand Junction | * | * | * | * | * | * | * |
| Greenfield | * | * | * | * | * | * | * |
| Grinnell | * | * | * | * | * | * | * |
| Grundy Ctr | * | * | * | * | * | * | * |
| Guthrie Ct | * | * | * | * | * | * | * |
| Hampton | * | * | * | * | * | * | * |
| Harlan | * | * | * | * | * | * | * |
| Hawarden | * | * | * | * | * | * | * |
| Hispanic Cln | * | * | * | * | * | * | * |
| Humboldt | * | * | * | * | * | * | * |
| ILH | 39.2 | 17.0 | 36.3 | 0.0 | 5.8 | 1.8 | 0.0 |
| Ida Grove | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Racial and Ethnic Distribution by Clinic

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Independen | * | * | * | * | * | * | * |
| Indianola | * | * | * | * | * | * | * |
| Iowa City | 36.1 | 40.2 | 19.1 | 0.3 | 2.1 | 2.3 | 0.0 |
| Iowa Falls | * | * | * | * | * | * | * |
| Jamaica | * | * | * | * | * | * | * |
| Jefferson | * | * | * | * | * | * | * |
| KR | * | * | * | * | * | * | * |
| Keokuk | * | * | * | * | * | * | * |
| Keosauqua | * | * | * | * | * | * | * |
| Knoxville | * | * | * | * | * | * | * |
| Lamoni | * | * | * | * | * | * | * |
| Le Mars | * | * | * | * | * | * | * |
| Lenox | * | * | * | * | * | * | * |
| Leon | * | * | * | * | * | * | * |
| Logan | * | * | * | * | * | * | * |
| Manchester | * | * | * | * | * | * | * |
| Maquoketa | * | * | * | * | * | * | * |
| Marengo | * | * | * | * | * | * | * |
| Marion | 84.5 | 7.8 | 4.7 | 0.0 | 1.6 | 1.6 | 0.0 |
| Marshalltown | 42.4 | 3.0 | 44.0 | 0.0 | 9.3 | 1.3 | 0.0 |
| Mason City | 77.2 | 4.3 | 12.1 | 0.0 | 3.9 | 2.6 | 0.0 |
| Monticello | * | * | * | * | * | * | * |
| Mt Ayr | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS



## Table 3A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Racial and Ethnic Distribution by Clinic

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Mt Pleasan | * | * | * | * | * | * | * |
| Muscatine | 58.3 | 6.8 | 32.6 | 0.0 | 2.3 | 0.0 | 0.0 |
| Nevada | * | * | * | * | * | * | * |
| New Hampto | * | * | * | * | * | * | * |
| Newton | * | * | * | * | * | * | * |
| North Towne | 61.8 | 18.0 | 7.8 | 0.0 | 7.4 | 5.1 | 0.0 |
| Northwood | * | * | * | * | * | * | * |
| Norwalk | * | * | * | * | * | * | * |
| Oelwein | * | * | * | * | * | * | * |
| Onawa | * | * | * | * | * | * | * |
| Orange Cty | * | * | * | * | * | * | * |
| Osage | * | * | * | * | * | * | * |
| Osceola | * | * | * | * | * | * | * |
| Oskaloosa | * | * | * | * | * | * | * |
| Ottumwa | 73.4 | 7.1 | 14.0 | 0.0 | 4.3 | 1.3 | 0.0 |
| Pella | * | * | * | * | * | * | * |
| Perry | * | * | * | * | * | * | * |
| Pocahontas | * | * | * | * | * | * | * |
| Pomeroy | * | * | * | * | * | * | * |
| Postville | * | * | * | * | * | * | * |
| Red Oak | * | * | * | * | * | * | * |
| Redfield | * | * | * | * | * | * | * |
| Refugee Cln | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Racial and Ethnic Distribution by Clinic

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Rock Rapid | * | * | * | * | * | * | * |
| Rock Valle | * | * | * | * | * | * | * |
| Rockwell C | * | * | * | * | * | * | * |
| SDM | 49.0 | 11.7 | 27.0 | 0.3 | 7.5 | 4.5 | 0.0 |
| Sac City | * | * | * | * | * | * | * |
| Sheldon | * | * | * | * | * | * | * |
| Shenandoah | * | * | * | * | * | * | * |
| Sibley | * | * | * | * | * | * | * |
| Sidney | * | * | * | * | * | * | * |
| Sigourney | * | * | * | * | * | * | * |
| Sioux Cent | * | * | * | * | * | * | * |
| Sioux City | 44.4 | 10.0 | 36.8 | 3.8 | 2.5 | 2.5 | 0.0 |
| Spencer | * | * | * | * | * | * | * |
| Spirit Lak | * | * | * | * | * | * | * |
| Storm Lake | 28.0 | 5.3 | 51.5 | 0.8 | 14.4 | 0.0 | 0.0 |
| Stuart | * | * | * | * | * | * | * |
| Tama | * | * | * | * | * | * | * |
| Tama/Settl | * | * | * | * | * | * | * |
| Tipton | * | * | * | * | * | * | * |
| Traer | * | * | * | * | * | * | * |
| UIHC | * | * | * | * | * | * | * |
| Vinton | * | * | * | * | * | * | * |
| WDM | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Racial and Ethnic Distribution by Clinic

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Wapello | * | * | * | * | * | * | * |
| Washington | * | * | * | * | * | * | * |
| Waterloo | 59.5 | 22.6 | 8.5 | 0.7 | 4.8 | 3.7 | 0.0 |
| Waukee | * | * | * | * | * | * | * |
| Waukon | * | * | * | * | * | * | * |
| Waverly | * | * | * | * | * | * | * |
| Wbstr Ct | * | * | * | * | * | * | * |
| West Union | * | * | * | * | * | * | * |
| Williamsburg | * | * | * | * | * | * | * |
| Winterset | * | * | * | * | * | * | * |
| **Iowa** | **61.3** | **13.2** | **18.7** | **0.5** | **3.7** | **2.6** | **0.0** |
| **National (Prior Year)** | **53.3** | **21.1** | **16.9** | **2.2** | **2.9** | **3.1** | **0.5** |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Ackley | * | * | * | * | * | * | * |
| Adel | * | * | * | * | * | * | * |
| Albia | * | * | * | * | * | * | * |
| Algona | * | * | * | * | * | * | * |
| Allen WHC | * | * | * | * | * | * | * |
| Allison | * | * | * | * | * | * | * |
| Altoona | * | * | * | * | * | * | * |
| Ames | 0.0 | 0.9 | 4.1 | 60.9 | 33.2 | 0.9 | 0.0 |
| Anamosa | * | * | * | * | * | * | * |
| Ankeny | * | * | * | * | * | * | * |
| Atlantic | * | * | * | * | * | * | * |
| Audubon | * | * | * | * | * | * | * |
| Avoca | * | * | * | * | * | * | * |
| Bedford | * | * | * | * | * | * | * |
| Belle Plaine | * | * | * | * | * | * | * |
| Bloomfield | * | * | * | * | * | * | * |
| Boone | * | * | * | * | * | * | * |
| Brdlwns RP | 0.0 | 1.8 | 6.8 | 56.4 | 31.6 | 3.3 | 0.0 |
| Brooklyn | * | * | * | * | * | * | * |
| Burlington | 0.0 | 5.1 | 9.1 | 55.8 | 28.9 | 1.0 | 0.0 |
| CHC | 0.4 | 3.1 | 5.3 | 66.7 | 23.6 | 0.9 | 0.0 |
| CR Urban | 0.0 | 1.0 | 6.8 | 62.6 | 27.0 | 2.7 | 0.0 |
| Carlisle | * | * | * | * | * | * | * |
| Carroll | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Cedar Falls | * | * | * | * | * | * | * |
| Centervill | * | * | * | * | * | * | * |
| Chariton | * | * | * | * | * | * | * |
| Charles City | * | * | * | * | * | * | * |
| Cherokee | * | * | * | * | * | * | * |
| Clarinda | * | * | * | * | * | * | * |
| Clarion | * | * | * | * | * | * | * |
| Clinton | 0.0 | 1.2 | 6.4 | 68.8 | 22.5 | 1.2 | 0.0 |
| Colfax | * | * | * | * | * | * | * |
| Columbus J | * | * | * | * | * | * | * |
| Corning | * | * | * | * | * | * | * |
| Corydon | * | * | * | * | * | * | * |
| Council Blfs | 0.0 | 3.1 | 5.8 | 64.0 | 26.1 | 1.0 | 0.0 |
| Cresco | * | * | * | * | * | * | * |
| Creston | * | * | * | * | * | * | * |
| Dallas Ctr | * | * | * | * | * | * | * |
| Decorah | * | * | * | * | * | * | * |
| Denison | 0.0 | 3.4 | 0.9 | 65.0 | 29.1 | 1.7 | 0.0 |
| Dewitt | * | * | * | * | * | * | * |
| Dubuque | 0.0 | 3.5 | 8.4 | 59.0 | 27.8 | 1.0 | 0.3 |
| Dyersville | * | * | * | * | * | * | * |
| Eagle Grove | * | * | * | * | * | * | * |
| Edgerton | 0.2 | 3.2 | 9.1 | 59.0 | 26.7 | 1.8 | 0.0 |
| Eldora | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Elkader | * | * | * | * | * | * | * |
| Emmetsburg | * | * | * | * | * | * | * |
| Esthervill | * | * | * | * | * | * | * |
| Fairfiel | * | * | * | * | * | * | * |
| Forest City | * | * | * | * | * | * | * |
| Fort Dodge | 0.5 | 5.0 | 6.9 | 60.4 | 26.2 | 1.0 | 0.0 |
| Ft Madison | * | * | * | * | * | * | * |
| Garner | * | * | * | * | * | * | * |
| Glenwood | * | * | * | * | * | * | * |
| Grand Junction | * | * | * | * | * | * | * |
| Greenfield | * | * | * | * | * | * | * |
| Grinnell | * | * | * | * | * | * | * |
| Grundy Ctr | * | * | * | * | * | * | * |
| Guthrie Ct | * | * | * | * | * | * | * |
| Hampton | * | * | * | * | * | * | * |
| Harlan | * | * | * | * | * | * | * |
| Hawarden | * | * | * | * | * | * | * |
| Hispanic Cln | * | * | * | * | * | * | * |
| Humboldt | * | * | * | * | * | * | * |
| ILH | 0.0 | 3.3 | 7.3 | 56.6 | 30.3 | 2.4 | 0.0 |
| Ida Grove | * | * | * | * | * | * | * |
| Independen | * | * | * | * | * | * | * |
| Indianola | * | * | * | * | * | * | * |
| Iowa City | 0.0 | 0.5 | 2.8 | 53.6 | 39.7 | 3.4 | 0.0 |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Iowa Falls | * | * | * | * | * | * | * |
| Jamaica | * | * | * | * | * | * | * |
| Jefferson | * | * | * | * | * | * | * |
| KR | * | * | * | * | * | * | * |
| Keokuk | * | * | * | * | * | * | * |
| Keosauqua | * | * | * | * | * | * | * |
| Knoxville | * | * | * | * | * | * | * |
| Lamoni | * | * | * | * | * | * | * |
| Le Mars | * | * | * | * | * | * | * |
| Lenox | * | * | * | * | * | * | * |
| Leon | * | * | * | * | * | * | * |
| Logan | * | * | * | * | * | * | * |
| Manchester | * | * | * | * | * | * | * |
| Maquoketa | * | * | * | * | * | * | * |
| Marengo | * | * | * | * | * | * | * |
| Marion | 0.0 | 0.0 | 4.7 | 63.6 | 28.7 | 3.1 | 0.0 |
| Marshalltown | 0.0 | 2.6 | 6.3 | 56.6 | 33.4 | 1.0 | 0.0 |
| Mason City | 0.0 | 0.4 | 5.6 | 67.7 | 24.1 | 2.2 | 0.0 |
| Monticello | * | * | * | * | * | * | * |
| Mt Ayr | * | * | * | * | * | * | * |
| Mt Pleasan | * | * | * | * | * | * | * |
| Muscatine | 0.0 | 2.7 | 8.7 | 60.2 | 26.1 | 2.3 | 0.0 |
| Nevada | * | * | * | * | * | * | * |
| New Hampto | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Newton | * | * | * | * | * | * | * |
| North Towne | 0.0 | 1.4 | 6.5 | 59.0 | 31.8 | 1.4 | 0.0 |
| Northwood | * | * | * | * | * | * | * |
| Norwalk | * | * | * | * | * | * | * |
| Oelwein | * | * | * | * | * | * | * |
| Onawa | * | * | * | * | * | * | * |
| Orange Cty | * | * | * | * | * | * | * |
| Osage | * | * | * | * | * | * | * |
| Osceola | * | * | * | * | * | * | * |
| Oskaloosa | * | * | * | * | * | * | * |
| Ottumwa | 0.0 | 1.8 | 7.4 | 64.7 | 24.6 | 1.3 | 0.3 |
| Pella | * | * | * | * | * | * | * |
| Perry | * | * | * | * | * | * | * |
| Pocahontas | * | * | * | * | * | * | * |
| Pomeroy | * | * | * | * | * | * | * |
| Postville | * | * | * | * | * | * | * |
| Red Oak | * | * | * | * | * | * | * |
| Redfield | * | * | * | * | * | * | * |
| Refugee Cln | * | * | * | * | * | * | * |
| Rock Rapid | * | * | * | * | * | * | * |
| Rock Valle | * | * | * | * | * | * | * |
| Rockwell C | * | * | * | * | * | * | * |
| SDM | 0.0 | 3.1 | 8.1 | 61.0 | 24.8 | 3.1 | 0.0 |
| Sac City | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Sheldon | * | * | * | * | * | * | * |
| Shenandoah | * | * | * | * | * | * | * |
| Sibley | * | * | * | * | * | * | * |
| Sidney | * | * | * | * | * | * | * |
| Sigourney | * | * | * | * | * | * | * |
| Sioux Cent | * | * | * | * | * | * | * |
| Sioux City | 0.2 | 2.2 | 6.5 | 57.1 | 32.4 | 1.7 | 0.0 |
| Spencer | * | * | * | * | * | * | * |
| Spirit Lak | * | * | * | * | * | * | * |
| Storm Lake | 0.0 | 3.8 | 3.8 | 56.8 | 31.1 | 4.5 | 0.0 |
| Stuart | * | * | * | * | * | * | * |
| Tama | * | * | * | * | * | * | * |
| Tama/Settl | * | * | * | * | * | * | * |
| Tipton | * | * | * | * | * | * | * |
| Traer | * | * | * | * | * | * | * |
| UIHC | * | * | * | * | * | * | * |
| Vinton | * | * | * | * | * | * | * |
| WDM | * | * | * | * | * | * | * |
| Wapello | * | * | * | * | * | * | * |
| Washington | * | * | * | * | * | * | * |
| Waterloo | 0.3 | 1.5 | 4.2 | 65.8 | 27.1 | 1.0 | 0.0 |
| Waukee | * | * | * | * | * | * | * |
| Waukon | * | * | * | * | * | * | * |
| Waverly | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Wbstr Ct | * | * | * | * | * | * | * |
| West Union | * | * | * | * | * | * | * |
| Williamsburg | * | * | * | * | * | * | * |
| Winterset | * | * | * | * | * | * | * |
| **Iowa** | **0.1** | **2.3** | **6.1** | **60.7** | **28.8** | **2.0** | **0.0** |
| **National (Prior Year)** | **0.1** | **2.9** | **8.1** | **62.2** | **24.7** | **1.8** | **0.1** |

(1) Reporting period is January 1 through December 31
(2) Excludes data with errors
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Ackley | | * | * | | * | * | | * | | * | | * | | * |
| Adel | | * | * | | * | * | | * | | * | | * | | * |
| Albia | | * | * | | * | * | | * | | * | | * | | * |
| Algona | | * | * | | * | * | | * | | * | | * | | * |
| Allen WHC | | * | * | | * | * | | * | | * | | * | | * |
| Allison | | * | * | | * | * | | * | | * | | * | | * |
| Altoona | | * | * | | * | * | | * | | * | | * | | * |
| Ames | | 4.4 (18) | 52.8 (1) | | 21.4 (8) | 40.9 (2) | | * | | * | | * | | * |
| Anamosa | | * | * | | * | * | | * | | * | | * | | * |
| Ankeny | | * | * | | * | * | | * | | * | | * | | * |
| Atlantic | | * | * | | * | * | | * | | * | | * | | * |
| Audubon | | * | * | | * | * | | * | | * | | * | | * |
| Avoca | | * | * | | * | * | | * | | * | | * | | * |
| Bedford | | * | * | | * | * | | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | * | | * | | * | | * | | * |
| Bloomfield | | * | * | | * | * | | * | | * | | * | | * |
| Boone | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Brdlwns RP | | 3.4 (8) | 55.2 (2) | | 21.9 (13) | 43.1 (6) | | 1.6 (1) | | * | | * | | * |
| Brooklyn | | * | * | | * | * | | * | | * | | * | | * |
| Burlington | | 4.3 (16) | 60.0 (7) | | 12.0 (1) | 56.4 (21) | | * | | * | | * | | * |
| CHC | | 2.5 (4) | 68.2 (21) | | 18.0 (3) | 53.3 (19) | | * | | * | | * | | * |
| CR Urban | | 2.9 (6) | 61.0 (10) | | 21.5 (10) | 49.1 (14) | | * | | * | | * | | * |
| Carlisle | | * | * | | * | * | | * | | * | | * | | * |
| Carroll | | * | * | | * | * | | * | | * | | * | | * |
| Cedar Falls | | * | * | | * | * | | * | | * | | * | | * |
| Centervill | | * | * | | * | * | | * | | * | | * | | * |
| Chariton | | * | * | | * | * | | * | | * | | * | | * |
| Charles City | | * | * | | * | * | | * | | * | | * | | * |
| Cherokee | | * | * | | * | * | | * | | * | | * | | * |
| Clarinda | | * | * | | * | * | | * | | * | | * | | * |
| Clarion | | * | * | | * | * | | * | | * | | * | | * |
| Clinton | | 4.3 (16) | 65.2 (19) | | 23.4 (15) | 53.9 (20) | | * | | * | | * | | * |
| Colfax | | * | * | | * | * | | * | | * | | * | | * |
| Columbus J | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Corning | | * | * | | * | * | | * | | * | | * | | * |
| Corydon | | * | * | | * | * | | * | | * | | * | | * |
| Council Blfs | | 5.2 (20) | 60.8 (9) | | 21.4 (8) | 49.5 (15) | | 7.3 (4) | | * | | * | | * |
| Cresco | | * | * | | * | * | | * | | * | | * | | * |
| Creston | | * | * | | * | * | | * | | * | | * | | * |
| Dallas Ctr | | * | * | | * | * | | * | | * | | * | | * |
| Decorah | | * | * | | * | * | | * | | * | | * | | * |
| Denison | | * | * | | * | * | | * | | * | | * | | * |
| Dewitt | | * | * | | * | * | | * | | * | | * | | * |
| Dubuque | | 4.9 (19) | 62.3 (13) | | 19.8 (6) | 51.7 (18) | | * | | * | | * | | * |
| Dyersville | | * | * | | * | * | | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | * | | * | | * | | * | | * |
| Edgerton | | 3.3 (7) | 59.4 (5) | | 18.6 (4) | 46.7 (11) | | * | | * | | * | | * |
| Eldora | | * | * | | * | * | | * | | * | | * | | * |
| Elkader | | * | * | | * | * | | * | | * | | * | | * |
| Emmetsburg | | * | * | | * | * | | * | | * | | * | | * |
| Esthervill | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Maternal Health Indicators by Clinic

### Comparison of Maternal Health Indicators (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Fairfiel | | * | * | | * | * | | * | | * | | * | | * |
| Forest City | | * | * | | * | * | | * | | * | | * | | * |
| Fort Dodge | | 2.4 (3) | 62.8 (15) | | 25.3 (21) | 44.3 (7) | | * | | * | | * | | * |
| Ft Madison | | * | * | | * | * | | * | | * | | * | | * |
| Garner | | * | * | | * | * | | * | | * | | * | | * |
| Glenwood | | * | * | | * | * | | * | | * | | * | | * |
| Grand Junction | | * | * | | * | * | | * | | * | | * | | * |
| Greenfield | | * | * | | * | * | | * | | * | | * | | * |
| Grinnell | | * | * | | * | * | | * | | * | | * | | * |
| Grundy Ctr | | * | * | | * | * | | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | * | | * | | * | | * | | * |
| Hampton | | * | * | | * | * | | * | | * | | * | | * |
| Harlan | | * | * | | * | * | | * | | * | | * | | * |
| Hawarden | | * | * | | * | * | | * | | * | | * | | * |
| Hispanic Cln | | * | * | | * | * | | * | | * | | * | | * |
| Humboldt | | * | * | | * | * | | * | | * | | * | | * |
| ILH | | 5.3 (21) | 57.8 (4) | | 24.4 (20) | 42.5 (5) | | 1.9 (3) | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Ida Grove | | * | * | | * | * | | * | | * | | * | | * |
| Independen | | * | * | | * | * | | * | | * | | * | | * |
| Indianola | | * | * | | * | * | | * | | * | | * | | * |
| Iowa City | | 2.5 (4) | 59.7 (6) | | 23.5 (16) | 42.0 (4) | | * | | 28.6 (2) | | * | | * |
| Iowa Falls | | * | * | | * | * | | * | | * | | * | | * |
| Jamaica | | * | * | | * | * | | * | | * | | * | | * |
| Jefferson | | * | * | | * | * | | * | | * | | * | | * |
| KR | | * | * | | * | * | | * | | * | | * | | * |
| Keokuk | | * | * | | * | * | | * | | * | | * | | * |
| Keosauqua | | * | * | | * | * | | * | | * | | * | | * |
| Knoxville | | * | * | | * | * | | * | | * | | * | | * |
| Lamoni | | * | * | | * | * | | * | | * | | * | | * |
| Le Mars | | * | * | | * | * | | * | | * | | * | | * |
| Lenox | | * | * | | * | * | | * | | * | | * | | * |
| Leon | | * | * | | * | * | | * | | * | | * | | * |
| Logan | | * | * | | * | * | | * | | * | | * | | * |
| Manchester | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Maquoketa | | * | * | | * | * | | * | | * | | * | | * |
| Marengo | | * | * | | * | * | | * | | * | | * | | * |
| Marion | | 1.9 (2) | 64.5 (17) | | 23.3 (14) | 45.6 (9) | | * | | * | | * | | * |
| Marshalltown | | 3.7 (11) | 60.7 (8) | | 24.0 (18) | 39.1 (1) | | * | | * | | * | | * |
| Mason City | | 3.6 (10) | 65.8 (20) | | 16.7 (2) | 51.6 (17) | | * | | * | | * | | * |
| Monticello | | * | * | | * | * | | * | | * | | * | | * |
| Mt Ayr | | * | * | | * | * | | * | | * | | * | | * |
| Mt Pleasan | | * | * | | * | * | | * | | * | | * | | * |
| Muscatine | | 3.8 (12) | 62.6 (14) | | 19.5 (5) | 47.8 (13) | | * | | * | | * | | * |
| Nevada | | * | * | | * | * | | * | | * | | * | | * |
| New Hampto | | * | * | | * | * | | * | | * | | * | | * |
| Newton | | * | * | | * | * | | * | | * | | * | | * |
| North Towne | | 1.8 (1) | 64.5 (17) | | 21.1 (7) | 50.9 (16) | | * | | * | | * | | * |
| Northwood | | * | * | | * | * | | * | | * | | * | | * |
| Norwalk | | * | * | | * | * | | * | | * | | * | | * |
| Oelwein | | * | * | | * | * | | * | | * | | * | | * |
| Onawa | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Orange Cty | | * | * | | * | * | | * | | * | | * | | * |
| Osage | | * | * | | * | * | | * | | * | | * | | * |
| Osceola | | * | * | | * | * | | * | | * | | * | | * |
| Oskaloosa | | * | * | | * | * | | * | | * | | * | | * |
| Ottumwa | | 4.1 (14) | 61.5 (12) | | 24.3 (19) | 45.8 (10) | | 31.7 (6) | | * | | * | | * |
| Pella | | * | * | | * | * | | * | | * | | * | | * |
| Perry | | * | * | | * | * | | * | | * | | * | | * |
| Pocahontas | | * | * | | * | * | | * | | * | | * | | * |
| Pomeroy | | * | * | | * | * | | * | | * | | * | | * |
| Postville | | * | * | | * | * | | * | | * | | * | | * |
| Red Oak | | * | * | | * | * | | * | | * | | * | | * |
| Redfield | | * | * | | * | * | | * | | * | | * | | * |
| Refugee Cln | | * | * | | * | * | | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | * | | * | | * | | * | | * |
| Rock Valle | | * | * | | * | * | | * | | * | | * | | * |
| Rockwell C | | * | * | | * | * | | * | | * | | * | | * |
| SDM | | 3.8 (12) | 57.4 (3) | | 23.7 (17) | 45.5 (8) | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Sac City | | * | * | | * | * | | * | | * | | * | | * |
| Sheldon | | * | * | | * | * | | * | | * | | * | | * |
| Shenandoah | | * | * | | * | * | | * | | * | | * | | * |
| Sibley | | * | * | | * | * | | * | | * | | * | | * |
| Sidney | | * | * | | * | * | | * | | * | | * | | * |
| Sigourney | | * | * | | * | * | | * | | * | | * | | * |
| Sioux Cent | | * | * | | * | * | | * | | * | | * | | * |
| Sioux City | | 3.5 (9) | 63.2 (16) | | 21.6 (12) | 41.9 (3) | | 1.7 (2) | | 3.5 (1) | | * | | * |
| Spencer | | * | * | | * | * | | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | * | | * | | * | | * | | * |
| Storm Lake | | * | * | | * | * | | * | | * | | * | | * |
| Stuart | | * | * | | * | * | | * | | * | | * | | * |
| Tama | | * | * | | * | * | | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | * | | * | | * | | * | | * |
| Tipton | | * | * | | * | * | | * | | * | | * | | * |
| Traer | | * | * | | * | * | | * | | * | | * | | * |
| UIHC | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Vinton | | * | * | | * | * | | * | | * | | * | | * |
| WDM | | * | * | | * | * | | * | | * | | * | | * |
| Wapello | | * | * | | * | * | | * | | * | | * | | * |
| Washington | | * | * | | * | * | | * | | * | | * | | * |
| Waterloo | | 4.2 (15) | 61.2 (11) | | 21.5 (10) | 47.2 (12) | | 10.3 (5) | | * | | * | | * |
| Waukee | | * | * | | * | * | | * | | * | | * | | * |
| Waukon | | * | * | | * | * | | * | | * | | * | | * |
| Waverly | | * | * | | * | * | | * | | * | | * | | * |
| Wbstr Ct | | * | * | | * | * | | * | | * | | * | | * |
| West Union | | * | * | | * | * | | * | | * | | * | | * |
| Williamsburg | | * | * | | * | * | | * | | * | | * | | * |
| Winterset | | * | * | | * | * | | * | | * | | * | | * |
| **Iowa** | | **3.5** | **61.2** | | **22.5** | **45.7** | | **14.4** | | **14.4** | | **100.0** | | **100.0** |
| **National (Prior Year)** | **120,741** | **4.0** | **59.6** | **97,554** | **22.4** | **47.7** | **26,239** | **27.4** | **53,856** | **17.5** | **104,317** | **9.9** | **103,483** | **9.3** |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

| Comparison of Maternal Behavioral Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ackley | | * | * | | * | | * | | * |
| Adel | | * | * | | * | | * | | * |
| Albia | | * | * | | * | | * | | * |
| Algona | | * | * | | * | | * | | * |
| Allen WHC | | * | * | | * | | * | | * |
| Allison | | * | * | | * | | * | | * |
| Altoona | | * | * | | * | | * | | * |
| Ames | | 29.8 (14) | 16.3 (14) | | 1.2 (11) | | 26.6 (1) | | 87.3 (11) |
| Anamosa | | * | * | | * | | * | | * |
| Ankeny | | * | * | | * | | * | | * |
| Atlantic | | * | * | | * | | * | | * |
| Audubon | | * | * | | * | | * | | * |
| Avoca | | * | * | | * | | * | | * |
| Bedford | | * | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | | * | | * |
| Bloomfield | | * | * | | * | | * | | * |
| Boone | | * | * | | * | | * | | * |
| Brdlwns RP | | 28.4 (16) | 11.8 (11) | | 1.4 (9) | | 10.4 (11) | | 84.9 (16) |
| Brooklyn | | * | * | | * | | * | | * |
| Burlington | | 35.3 (6) | 6.5 (2) | | 0.0 (20) | | 22.5 (2) | | 92.0 (3) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

| Comparison of Maternal Behavioral Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| CHC | | 33.8 (9) | 22.1 (17) | | 0.6 (18) | | * | | * |
| CR Urban | | 23.7 (19) | 10.5 (8) | | 1.3 (10) | | 11.9 (8) | | 87.9 (9) |
| Carlisle | | * | * | | * | | * | | * |
| Carroll | | * | * | | * | | * | | * |
| Cedar Falls | | * | * | | * | | * | | * |
| Centervill | | * | * | | * | | * | | * |
| Chariton | | * | * | | * | | * | | * |
| Charles City | | * | * | | * | | * | | * |
| Cherokee | | * | * | | * | | * | | * |
| Clarinda | | * | * | | * | | * | | * |
| Clarion | | * | * | | * | | * | | * |
| Clinton | | 42.1 (2) | 11.3 (10) | | 0.7 (15) | | * | | 88.0 (8) |
| Colfax | | * | * | | * | | * | | * |
| Columbus J | | * | * | | * | | * | | * |
| Corning | | * | * | | * | | * | | * |
| Corydon | | * | * | | * | | * | | * |
| Council Blfs | | 33.5 (11) | 6.6 (3) | | 0.9 (13) | | 20.3 (4) | | 87.6 (10) |
| Cresco | | * | * | | * | | * | | * |
| Creston | | * | * | | * | | * | | * |
| Dallas Ctr | | * | * | | * | | * | | * |
| Decorah | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

| Comparison of Maternal Behavioral Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Denison | | * | * | | 0.0 (20) | | * | | * |
| Dewitt | | * | * | | * | | * | | * |
| Dubuque | | 33.7 (10) | 14.8 (13) | | 4.1 (1) | | 10.4 (11) | | 90.0 (5) |
| Dyersville | | * | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | | * | | * |
| Edgerton | | 35.8 (5) | 10.9 (9) | | 2.6 (4) | | * | | * |
| Eldora | | * | * | | * | | * | | * |
| Elkader | | * | * | | * | | * | | * |
| Emmetsburg | | * | * | | * | | * | | * |
| Esthervill | | * | * | | * | | * | | * |
| Fairfiel | | * | * | | * | | * | | * |
| Forest City | | * | * | | * | | * | | * |
| Fort Dodge | | 34.4 (8) | 21.3 (16) | | 2.2 (6) | | 17.4 (6) | | 82.3 (17) |
| Ft Madison | | * | * | | * | | * | | * |
| Garner | | * | * | | * | | * | | * |
| Glenwood | | * | * | | * | | * | | * |
| Grand Junction | | * | * | | * | | * | | * |
| Greenfield | | * | * | | * | | * | | * |
| Grinnell | | * | * | | * | | * | | * |
| Grundy Ctr | | * | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

| Comparison of Maternal Behavioral Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Hampton | | * | * | | * | | * | | * |
| Harlan | | * | * | | * | | * | | * |
| Hawarden | | * | * | | * | | * | | * |
| Hispanic Cln | | * | * | | * | | * | | * |
| Humboldt | | * | * | | * | | * | | * |
| ILH | | 31.1 (13) | 12.1 (12) | | 0.8 (14) | | 11.7 (9) | | 88.3 (7) |
| Ida Grove | | * | * | | * | | * | | * |
| Independen | | * | * | | * | | * | | * |
| Indianola | | * | * | | * | | * | | * |
| Iowa City | | 15.2 (20) | 23.4 (18) | | 1.7 (8) | | * | | 87.1 (12) |
| Iowa Falls | | * | * | | * | | * | | * |
| Jamaica | | * | * | | * | | * | | * |
| Jefferson | | * | * | | * | | * | | * |
| KR | | * | * | | * | | * | | * |
| Keokuk | | * | * | | * | | * | | * |
| Keosauqua | | * | * | | * | | * | | * |
| Knoxville | | * | * | | * | | * | | * |
| Lamoni | | * | * | | * | | * | | * |
| Le Mars | | * | * | | * | | * | | * |
| Lenox | | * | * | | * | | * | | * |
| Leon | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

| Comparison of Maternal Behavioral Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Logan | | * | * | | * | | * | | * |
| Manchester | | * | * | | * | | * | | * |
| Maquoketa | | * | * | | * | | * | | * |
| Marengo | | * | * | | * | | * | | * |
| Marion | | * | * | | 0.0 (20) | | * | | * |
| Marshalltown | | 47.1 (1) | 6.3 (1) | | 0.7 (15) | | 11.1 (10) | | 85.8 (13) |
| Mason City | | 37.3 (4) | 21.1 (15) | | 2.0 (7) | | 21.5 (3) | | 85.2 (15) |
| Monticello | | * | * | | * | | * | | * |
| Mt Ayr | | * | * | | * | | * | | * |
| Mt Pleasan | | * | * | | * | | * | | * |
| Muscatine | | 39.5 (3) | 9.3 (5) | | 2.8 (2) | | 16.3 (7) | | 90.8 (4) |
| Nevada | | * | * | | * | | * | | * |
| New Hampto | | * | * | | * | | * | | * |
| Newton | | * | * | | * | | * | | * |
| North Towne | | 28.1 (17) | 8.9 (4) | | 0.5 (19) | | * | | 92.4 (2) |
| Northwood | | * | * | | * | | * | | * |
| Norwalk | | * | * | | * | | * | | * |
| Oelwein | | * | * | | * | | * | | * |
| Onawa | | * | * | | * | | * | | * |
| Orange Cty | | * | * | | * | | * | | * |
| Osage | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

| Comparison of Maternal Behavioral Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Osceola | | * | * | | * | | * | | * |
| Oskaloosa | | * | * | | * | | * | | * |
| Ottumwa | | 24.7 (18) | 24.4 (20) | | 2.3 (5) | | * | | * |
| Pella | | * | * | | * | | * | | * |
| Perry | | * | * | | * | | * | | * |
| Pocahontas | | * | * | | * | | * | | * |
| Pomeroy | | * | * | | * | | * | | * |
| Postville | | * | * | | * | | * | | * |
| Red Oak | | * | * | | * | | * | | * |
| Redfield | | * | * | | * | | * | | * |
| Refugee Cln | | * | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | | * | | * |
| Rock Valle | | * | * | | * | | * | | * |
| Rockwell C | | * | * | | * | | * | | * |
| SDM | | 34.8 (7) | 10.2 (7) | | 1.0 (12) | | * | | 90.0 (5) |
| Sac City | | * | * | | * | | * | | * |
| Sheldon | | * | * | | * | | * | | * |
| Shenandoah | | * | * | | * | | * | | * |
| Sibley | | * | * | | * | | * | | * |
| Sidney | | * | * | | * | | * | | * |
| Sigourney | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

| Comparison of Maternal Behavioral Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Sioux Cent | | * | * | | * | | * | | * |
| Sioux City | | 28.5 (15) | 9.8 (6) | | 0.7 (15) | | * | | 93.4 (1) |
| Spencer | | * | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | | * | | * |
| Storm Lake | | * | * | | 0.0 (20) | | * | | * |
| Stuart | | * | * | | * | | * | | * |
| Tama | | * | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | | * | | * |
| Tipton | | * | * | | * | | * | | * |
| Traer | | * | * | | * | | * | | * |
| UIHC | | * | * | | * | | * | | * |
| Vinton | | * | * | | * | | * | | * |
| WDM | | * | * | | * | | * | | * |
| Wapello | | * | * | | * | | * | | * |
| Washington | | * | * | | * | | * | | * |
| Waterloo | | 33.5 (11) | 23.9 (19) | | 2.7 (3) | | 17.7 (5) | | 85.6 (14) |
| Waukee | | * | * | | * | | * | | * |
| Waukon | | * | * | | * | | * | | * |
| Waverly | | * | * | | * | | * | | * |
| Wbstr Ct | | * | * | | * | | * | | * |
| West Union | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

| Comparison of Maternal Behavioral Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Williamsburg | | * | * | | * | | * | | * |
| Winterset | | * | * | | * | | * | | * |
| **Iowa** | | **30.9** | **13.1** | | **1.5** | | **19.6** | | **5317.0** |
| **National (Prior Year)** | **109,069** | **69.2** | **7.4** | **136,111** | **23.4** | **99,278** | **22.2** | **98,734** | **88.0** |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ackley | | * | | * | | * | | * | | * |
| Adel | | * | | * | | * | | * | | * |
| Albia | | * | | * | | * | | * | | * |
| Algona | | * | | * | | * | | * | | * |
| Allen WHC | | * | | * | | * | | * | | * |
| Allison | | * | | * | | * | | * | | * |
| Altoona | | * | | * | | * | | * | | * |
| Ames | | 26.3 (9) | | 16.8 (11) | | * | | 10.1 (16) | | 1.1 (5) |
| Anamosa | | * | | * | | * | | * | | * |
| Ankeny | | * | | * | | * | | * | | * |
| Atlantic | | * | | * | | * | | * | | * |
| Audubon | | * | | * | | * | | * | | * |
| Avoca | | * | | * | | * | | * | | * |
| Bedford | | * | | * | | * | | * | | * |
| Belle Plaine | | * | | * | | * | | * | | * |
| Bloomfield | | * | | * | | * | | * | | * |
| Boone | | * | | * | | * | | * | | * |
| Brdlwns RP | | 21.3 (5) | | 10.8 (5) | | 29.0 (1) | | 4.7 (4) | | 1.4 (6) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Brooklyn | | * | | * | | * | | * | | * |
| Burlington | | 43.1 (20) | | 23.4 (19) | | * | | 17.5 (22) | | 2.8 (13) |
| CHC | | 34.3 (16) | | 20.4 (15) | | * | | 17.4 (21) | | 5.4 (21) |
| CR Urban | | 23.9 (7) | | 13.5 (7) | | * | | 7.7 (14) | | 0.7 (3) |
| Carlisle | | * | | * | | * | | * | | * |
| Carroll | | * | | * | | * | | * | | * |
| Cedar Falls | | * | | * | | * | | * | | * |
| Centervill | | * | | * | | * | | * | | * |
| Chariton | | * | | * | | * | | * | | * |
| Charles City | | * | | * | | * | | * | | * |
| Cherokee | | * | | * | | * | | * | | * |
| Clarinda | | * | | * | | * | | * | | * |
| Clarion | | * | | * | | * | | * | | * |
| Clinton | | 38.7 (18) | | 27.7 (22) | | * | | 4.9 (6) | | 2.9 (15) |
| Colfax | | * | | * | | * | | * | | * |
| Columbus J | | * | | * | | * | | * | | * |
| Corning | | * | | * | | * | | * | | * |
| Corydon | | * | | * | | * | | * | | * |
| Council Blfs | | 32.6 (15) | | 23.1 (18) | | 19.8 (2) | | 5.5 (8) | | 0.9 (4) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Cresco | | * | | * | | * | | * | | * |
| Creston | | * | | * | | * | | * | | * |
| Dallas Ctr | | * | | * | | * | | * | | * |
| Decorah | | * | | * | | * | | * | | * |
| Denison | | * | | * | | * | | * | | * |
| Dewitt | | * | | * | | * | | * | | * |
| Dubuque | | 15.8 (3) | | 9.7 (4) | | * | | 0.0 (1) | | 0.0 (1) |
| Dyersville | | * | | * | | * | | * | | * |
| Eagle Grove | | * | | * | | * | | * | | * |
| Edgerton | | 29.1 (14) | | 17.2 (12) | | * | | 11.0 (19) | | 2.0 (10) |
| Eldora | | * | | * | | * | | * | | * |
| Elkader | | * | | * | | * | | * | | * |
| Emmetsburg | | * | | * | | * | | * | | * |
| Esthervill | | * | | * | | * | | * | | * |
| Fairfiel | | * | | * | | * | | * | | * |
| Forest City | | * | | * | | * | | * | | * |
| Fort Dodge | | 43.5 (21) | | 22.9 (17) | | * | | 6.0 (10) | | 2.8 (13) |
| Ft Madison | | * | | * | | * | | * | | * |
| Garner | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Glenwood | | * | | * | | * | | * | | * |
| Grand Junction | | * | | * | | * | | * | | * |
| Greenfield | | * | | * | | * | | * | | * |
| Grinnell | | * | | * | | * | | * | | * |
| Grundy Ctr | | * | | * | | * | | * | | * |
| Guthrie Ct | | * | | * | | * | | * | | * |
| Hampton | | * | | * | | * | | * | | * |
| Harlan | | * | | * | | * | | * | | * |
| Hawarden | | * | | * | | * | | * | | * |
| Hispanic Cln | | * | | * | | * | | * | | * |
| Humboldt | | * | | * | | * | | * | | * |
| ILH | | 27.1 (11) | | 17.9 (13) | | * | | 7.0 (12) | | 1.5 (8) |
| Ida Grove | | * | | * | | * | | * | | * |
| Independen | | * | | * | | * | | * | | * |
| Indianola | | * | | * | | * | | * | | * |
| Iowa City | | 14.2 (2) | | 5.0 (1) | | * | | 3.3 (2) | | 0.0 (1) |
| Iowa Falls | | * | | * | | * | | * | | * |
| Jamaica | | * | | * | | * | | * | | * |
| Jefferson | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| KR | | * | | * | | * | | * | | * |
| Keokuk | | * | | * | | * | | * | | * |
| Keosauqua | | * | | * | | * | | * | | * |
| Knoxville | | * | | * | | * | | * | | * |
| Lamoni | | * | | * | | * | | * | | * |
| Le Mars | | * | | * | | * | | * | | * |
| Lenox | | * | | * | | * | | * | | * |
| Leon | | * | | * | | * | | * | | * |
| Logan | | * | | * | | * | | * | | * |
| Manchester | | * | | * | | * | | * | | * |
| Maquoketa | | * | | * | | * | | * | | * |
| Marengo | | * | | * | | * | | * | | * |
| Marion | | * | | 19.3 (14) | | * | | 4.8 (5) | | 2.6 (12) |
| Marshalltown | | 17.8 (4) | | 9.6 (3) | | * | | 5.8 (9) | | 1.7 (9) |
| Mason City | | 35.7 (17) | | 23.7 (20) | | * | | 7.2 (13) | | 1.4 (6) |
| Monticello | | * | | * | | * | | * | | * |
| Mt Ayr | | * | | * | | * | | * | | * |
| Mt Pleasan | | * | | * | | * | | * | | * |
| Muscatine | | 23.1 (6) | | 12.3 (6) | | * | | 10.9 (18) | | 4.4 (20) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Nevada | | * | | * | | * | | * | | * |
| New Hampto | | * | | * | | * | | * | | * |
| Newton | | * | | * | | * | | * | | * |
| North Towne | | 27.4 (12) | | 13.7 (8) | | * | | 5.0 (7) | | 2.0 (10) |
| Northwood | | * | | * | | * | | * | | * |
| Norwalk | | * | | * | | * | | * | | * |
| Oelwein | | * | | * | | * | | * | | * |
| Onawa | | * | | * | | * | | * | | * |
| Orange Cty | | * | | * | | * | | * | | * |
| Osage | | * | | * | | * | | * | | * |
| Osceola | | * | | * | | * | | * | | * |
| Oskaloosa | | * | | * | | * | | * | | * |
| Ottumwa | | 27.5 (13) | | 14.0 (9) | | * | | 11.6 (20) | | 4.2 (19) |
| Pella | | * | | * | | * | | * | | * |
| Perry | | * | | * | | * | | * | | * |
| Pocahontas | | * | | * | | * | | * | | * |
| Pomeroy | | * | | * | | * | | * | | * |
| Postville | | * | | * | | * | | * | | * |
| Red Oak | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Redfield | | * | | * | | * | | * | | * |
| Refugee Cln | | * | | * | | * | | * | | * |
| Rock Rapid | | * | | * | | * | | * | | * |
| Rock Valle | | * | | * | | * | | * | | * |
| Rockwell C | | * | | * | | * | | * | | * |
| SDM | | 27.0 (10) | | 24.8 (21) | | * | | 7.8 (15) | | 3.9 (18) |
| Sac City | | * | | * | | * | | * | | * |
| Sheldon | | * | | * | | * | | * | | * |
| Shenandoah | | * | | * | | * | | * | | * |
| Sibley | | * | | * | | * | | * | | * |
| Sidney | | * | | * | | * | | * | | * |
| Sigourney | | * | | * | | * | | * | | * |
| Sioux Cent | | * | | * | | * | | * | | * |
| Sioux City | | 25.3 (8) | | 14.5 (10) | | * | | 6.1 (11) | | 2.9 (15) |
| Spencer | | * | | * | | * | | * | | * |
| Spirit Lak | | * | | * | | * | | * | | * |
| Storm Lake | | 11.0 (1) | | 5.4 (2) | | * | | 4.0 (3) | | 5.9 (22) |
| Stuart | | * | | * | | * | | * | | * |
| Tama | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Tama/Settl | | * | | * | | * | | * | | * |
| Tipton | | * | | * | | * | | * | | * |
| Traer | | * | | * | | * | | * | | * |
| UIHC | | * | | * | | * | | * | | * |
| Vinton | | * | | * | | * | | * | | * |
| WDM | | * | | * | | * | | * | | * |
| Wapello | | * | | * | | * | | * | | * |
| Washington | | * | | * | | * | | * | | * |
| Waterloo | | 38.7 (18) | | 21.1 (16) | | 18.3 (3) | | 10.3 (17) | | 3.5 (17) |
| Waukee | | * | | * | | * | | * | | * |
| Waukon | | * | | * | | * | | * | | * |
| Waverly | | * | | * | | * | | * | | * |
| Wbstr Ct | | * | | * | | * | | * | | * |
| West Union | | * | | * | | * | | * | | * |
| Williamsburg | | * | | * | | * | | * | | * |
| Winterset | | * | | * | | * | | * | | * |
| **Iowa** | | **28.1** | | **16.7** | | **26.3** | | **7.3** | | **2.3** |
| **National (Prior Year)** | **115,645** | **28.4** | **127,715** | **14.9** | **28,227** | **36.2** | **104,358** | **10.5** | **128,247** | **3.4** |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ackley | | * | * | | * | | * | | * |
| Adel | | * | * | | * | | * | | * |
| Albia | | * | * | | * | | * | | * |
| Algona | | * | * | | * | | * | | * |
| Allen WHC | | * | * | | * | | * | | * |
| Allison | | * | * | | * | | * | | * |
| Altoona | | * | * | | * | | * | | * |
| Ames | | 10.4 (22) | 10.0 (20) | | 13.5 (21) | | 3.5 (16) | | 79.1 (4) |
| Anamosa | | * | * | | * | | * | | * |
| Ankeny | | * | * | | * | | * | | * |
| Atlantic | | * | * | | * | | * | | * |
| Audubon | | * | * | | * | | * | | * |
| Avoca | | * | * | | * | | * | | * |
| Bedford | | * | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | | * | | * |
| Bloomfield | | * | * | | * | | * | | * |
| Boone | | * | * | | * | | * | | * |
| Brdlwns RP | | 7.7 (12) | 6.3 (4) | | 9.2 (8) | | 2.9 (12) | | 81.3 (2) |
| Brooklyn | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Burlington | | 6.9 (8) | 12.2 (23) | | 8.6 (4) | | 2.2 (3) | | 61.6 (21) |
| CHC | | 9.9 (20) | 6.6 (5) | | 12.4 (18) | | 5.4 (21) | | 60.1 (22) |
| CR Urban | | 5.8 (3) | 8.7 (15) | | 7.6 (2) | | 2.9 (12) | | 65.0 (19) |
| Carlisle | | * | * | | * | | * | | * |
| Carroll | | * | * | | * | | * | | * |
| Cedar Falls | | * | * | | * | | * | | * |
| Centervill | | * | * | | * | | * | | * |
| Chariton | | * | * | | * | | * | | * |
| Charles City | | * | * | | * | | * | | * |
| Cherokee | | * | * | | * | | * | | * |
| Clarinda | | * | * | | * | | * | | * |
| Clarion | | * | * | | * | | * | | * |
| Clinton | | 12.2 (23) | 10.3 (21) | | 13.1 (19) | | 2.7 (9) | | 71.2 (12) |
| Colfax | | * | * | | * | | * | | * |
| Columbus J | | * | * | | * | | * | | * |
| Corning | | * | * | | * | | * | | * |
| Corydon | | * | * | | * | | * | | * |
| Council Blfs | | 7.6 (10) | 7.0 (7) | | 13.4 (20) | | 1.3 (1) | | 75.7 (7) |
| Cresco | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Creston | | * | * | | * | | * | | * |
| Dallas Ctr | | * | * | | * | | * | | * |
| Decorah | | * | * | | * | | * | | * |
| Denison | | 8.8 (17) | 6.2 (3) | | 15.0 (22) | | * | | 65.5 (17) |
| Dewitt | | * | * | | * | | * | | * |
| Dubuque | | 6.9 (8) | 7.8 (12) | | 5.7 (1) | | 2.2 (3) | | 55.9 (23) |
| Dyersville | | * | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | | * | | * |
| Edgerton | | 9.7 (19) | 6.8 (6) | | 10.6 (15) | | 5.3 (20) | | 65.9 (16) |
| Eldora | | * | * | | * | | * | | * |
| Elkader | | * | * | | * | | * | | * |
| Emmetsburg | | * | * | | * | | * | | * |
| Esthervill | | * | * | | * | | * | | * |
| Fairfiel | | * | * | | * | | * | | * |
| Forest City | | * | * | | * | | * | | * |
| Fort Dodge | | 6.4 (7) | 5.3 (1) | | 7.6 (2) | | 2.5 (7) | | 70.6 (13) |
| Ft Madison | | * | * | | * | | * | | * |
| Garner | | * | * | | * | | * | | * |
| Glenwood | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

| Comparison of Infant Health Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Grand Junction | | * | * | | * | | * | | * |
| Greenfield | | * | * | | * | | * | | * |
| Grinnell | | * | * | | * | | * | | * |
| Grundy Ctr | | * | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | | * | | * |
| Hampton | | * | * | | * | | * | | * |
| Harlan | | * | * | | * | | * | | * |
| Hawarden | | * | * | | * | | * | | * |
| Hispanic Cln | | * | * | | * | | * | | * |
| Humboldt | | * | * | | * | | * | | * |
| ILH | | 7.6 (10) | 9.3 (16) | | 11.9 (17) | | 2.4 (5) | | 75.6 (8) |
| Ida Grove | | * | * | | * | | * | | * |
| Independen | | * | * | | * | | * | | * |
| Indianola | | * | * | | * | | * | | * |
| Iowa City | | 10.3 (21) | 7.1 (8) | | 10.5 (14) | | 4.4 (19) | | 86.8 (1) |
| Iowa Falls | | * | * | | * | | * | | * |
| Jamaica | | * | * | | * | | * | | * |
| Jefferson | | * | * | | * | | * | | * |
| KR | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Keokuk | | * | * | | * | | * | | * |
| Keosauqua | | * | * | | * | | * | | * |
| Knoxville | | * | * | | * | | * | | * |
| Lamoni | | * | * | | * | | * | | * |
| Le Mars | | * | * | | * | | * | | * |
| Lenox | | * | * | | * | | * | | * |
| Leon | | * | * | | * | | * | | * |
| Logan | | * | * | | * | | * | | * |
| Manchester | | * | * | | * | | * | | * |
| Maquoketa | | * | * | | * | | * | | * |
| Marengo | | * | * | | * | | * | | * |
| Marion | | 9.3 (18) | 9.3 (16) | | 15.1 (23) | | * | | 69.7 (14) |
| Marshalltown | | 5.9 (4) | 8.0 (13) | | 9.3 (9) | | 2.4 (5) | | 72.6 (10) |
| Mason City | | 6.0 (5) | 9.7 (19) | | 11.2 (16) | | 2.9 (12) | | 65.3 (18) |
| Monticello | | * | * | | * | | * | | * |
| Mt Ayr | | * | * | | * | | * | | * |
| Mt Pleasan | | * | * | | * | | * | | * |
| Muscatine | | 4.8 (1) | 6.0 (2) | | 10.4 (13) | | 2.6 (8) | | 79.6 (3) |
| Nevada | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| New Hampto | | * | * | | * | | * | | * |
| Newton | | * | * | | * | | * | | * |
| North Towne | | 7.7 (12) | 11.5 (22) | | 9.1 (6) | | 3.6 (17) | | 72.7 (9) |
| Northwood | | * | * | | * | | * | | * |
| Norwalk | | * | * | | * | | * | | * |
| Oelwein | | * | * | | * | | * | | * |
| Onawa | | * | * | | * | | * | | * |
| Orange Cty | | * | * | | * | | * | | * |
| Osage | | * | * | | * | | * | | * |
| Osceola | | * | * | | * | | * | | * |
| Oskaloosa | | * | * | | * | | * | | * |
| Ottumwa | | 6.1 (6) | 7.5 (11) | | 9.1 (6) | | 2.7 (9) | | 75.8 (6) |
| Pella | | * | * | | * | | * | | * |
| Perry | | * | * | | * | | * | | * |
| Pocahontas | | * | * | | * | | * | | * |
| Pomeroy | | * | * | | * | | * | | * |
| Postville | | * | * | | * | | * | | * |
| Red Oak | | * | * | | * | | * | | * |
| Redfield | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

| Comparison of Infant Health Indicators (2,3) | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| **Clinic** | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Refugee Cln | | * | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | | * | | * |
| Rock Valle | | * | * | | * | | * | | * |
| Rockwell C | | * | * | | * | | * | | * |
| SDM | | 7.8 (14) | 9.3 (16) | | 10.0 (11) | | 2.7 (9) | | 76.1 (5) |
| Sac City | | * | * | | * | | * | | * |
| Sheldon | | * | * | | * | | * | | * |
| Shenandoah | | * | * | | * | | * | | * |
| Sibley | | * | * | | * | | * | | * |
| Sidney | | * | * | | * | | * | | * |
| Sigourney | | * | * | | * | | * | | * |
| Sioux Cent | | * | * | | * | | * | | * |
| Sioux City | | 8.3 (15) | 8.3 (14) | | 10.1 (12) | | 3.0 (15) | | 71.6 (11) |
| Spencer | | * | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | | * | | * |
| Storm Lake | | 4.8 (1) | 7.1 (8) | | 8.7 (5) | | 1.9 (2) | | 61.7 (20) |
| Stuart | | * | * | | * | | * | | * |
| Tama | | * | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Tipton | | * | * | | * | | * | | * |
| Traer | | * | * | | * | | * | | * |
| UIHC | | * | * | | * | | * | | * |
| Vinton | | * | * | | * | | * | | * |
| WDM | | * | * | | * | | * | | * |
| Wapello | | * | * | | * | | * | | * |
| Washington | | * | * | | * | | * | | * |
| Waterloo | | 8.6 (16) | 7.1 (8) | | 9.5 (10) | | 3.8 (18) | | 68.8 (15) |
| Waukee | | * | * | | * | | * | | * |
| Waukon | | * | * | | * | | * | | * |
| Waverly | | * | * | | * | | * | | * |
| Wbstr Ct | | * | * | | * | | * | | * |
| West Union | | * | * | | * | | * | | * |
| Williamsburg | | * | * | | * | | * | | * |
| Winterset | | * | * | | * | | * | | * |
| **Iowa** | | **7.6** | **8.0** | | **10.2** | | **2.9** | | **72.0** |
| **National (Prior Year)** | **125,347** | **9.3** | **6.9** | **125,751** | **11.6** | **106,512** | **3.9** | **126,356** | **53.2** |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* National includes previous year's data for all participating States and ITOs.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Racial and Ethnic Distribution by Clinic

## Combined 3 Years

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Ackley | * | * | * | * | * | * | * |
| Adel | * | * | * | * | * | * | * |
| Albia | * | * | * | * | * | * | * |
| Algona | 79.8 | 3.5 | 14.0 | 0.0 | 0.0 | 2.6 | 0.0 |
| Allen WHC | 56.7 | 28.8 | 4.0 | 0.3 | 6.3 | 3.7 | 0.3 |
| Allison | * | * | * | * | * | * | * |
| Altoona | 82.1 | 7.7 | 5.1 | 0.0 | 2.6 | 2.6 | 0.0 |
| Ames | 68.7 | 7.0 | 10.6 | 0.2 | 11.9 | 1.5 | 0.0 |
| Anamosa | * | * | * | * | * | * | * |
| Ankeny | 80.5 | 6.8 | 6.0 | 0.0 | 3.0 | 3.8 | 0.0 |
| Atlantic | 88.9 | 0.6 | 9.9 | 0.0 | 0.0 | 0.6 | 0.0 |
| Audubon | * | * | * | * | * | * | * |
| Avoca | * | * | * | * | * | * | * |
| Bedford | * | * | * | * | * | * | * |
| Belle Plaine | * | * | * | * | * | * | * |
| Bloomfield | * | * | * | * | * | * | * |
| Boone | * | * | * | * | * | * | * |
| Brdlwns RP | 43.7 | 20.7 | 20.4 | 0.4 | 9.4 | 5.4 | 0.0 |
| Brooklyn | * | * | * | * | * | * | * |
| Burlington | 75.5 | 16.8 | 5.6 | 0.0 | 0.5 | 1.6 | 0.0 |
| CHC | 58.8 | 22.9 | 13.3 | 0.0 | 1.7 | 3.4 | 0.0 |
| CR Urban | 54.2 | 29.6 | 7.6 | 0.3 | 0.9 | 5.5 | 1.7 |
| Carlisle | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Racial and Ethnic Distribution by Clinic

## Combined 3 Years

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Carroll | 78.6 | 8.0 | 12.5 | 0.0 | 0.9 | 0.0 | 0.0 |
| Cedar Falls | * | * | * | * | * | * | * |
| Centervill | * | * | * | * | * | * | * |
| Chariton | * | * | * | * | * | * | * |
| Charles City | 77.2 | 8.9 | 7.9 | 0.0 | 4.5 | 1.5 | 0.0 |
| Cherokee | 90.7 | 1.6 | 7.8 | 0.0 | 0.0 | 0.0 | 0.0 |
| Clarinda | * | * | * | * | * | * | * |
| Clarion | 39.3 | 0.0 | 60.7 | 0.0 | 0.0 | 0.0 | 0.0 |
| Clinton | 78.4 | 11.2 | 7.9 | 0.0 | 0.3 | 2.1 | 0.0 |
| Colfax | * | * | * | * | * | * | * |
| Columbus J | * | * | * | * | * | * | * |
| Corning | * | * | * | * | * | * | * |
| Correctionvl | * | * | * | * | * | * | * |
| Corydon | * | * | * | * | * | * | * |
| Council Blfs | 77.9 | 3.4 | 15.1 | 0.5 | 0.5 | 2.5 | 0.0 |
| Cresco | * | * | * | * | * | * | * |
| Creston | 88.5 | 2.4 | 8.5 | 0.7 | 0.0 | 0.0 | 0.0 |
| Dallas Ctr | * | * | * | * | * | * | * |
| Decorah | * | * | * | * | * | * | * |
| Denison | 34.0 | 3.8 | 55.8 | 0.2 | 6.2 | 0.0 | 0.0 |
| Dewitt | * | * | * | * | * | * | * |
| Dubuque | 74.5 | 12.5 | 7.1 | 0.3 | 3.2 | 2.4 | 0.0 |
| Dyersville | * | * | * | * | * | * | * |
| Eagle Grove | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Racial and Ethnic Distribution by Clinic

## Combined 3 Years

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Edgerton | 61.8 | 22.7 | 9.4 | 0.4 | 0.9 | 4.7 | 0.0 |
| Edgerton CHC | * | * | * | * | * | * | * |
| Eldora | * | * | * | * | * | * | * |
| Elkader | 87.4 | 0.0 | 7.9 | 0.0 | 2.4 | 2.4 | 0.0 |
| Emmetsburg | * | * | * | * | * | * | * |
| Esthervill | 60.0 | 5.0 | 32.9 | 0.0 | 0.7 | 1.4 | 0.0 |
| Fairfiel | 84.6 | 1.9 | 6.7 | 0.0 | 4.8 | 1.9 | 0.0 |
| Fayette | * | * | * | * | * | * | * |
| Forest City | 87.8 | 0.8 | 9.2 | 0.8 | 0.0 | 1.5 | 0.0 |
| Fort Dodge | 79.7 | 6.7 | 11.0 | 0.3 | 0.0 | 2.2 | 0.0 |
| Ft Madison | 89.7 | 3.0 | 4.7 | 0.0 | 0.0 | 2.6 | 0.0 |
| Garner | * | * | * | * | * | * | * |
| Glenwood | * | * | * | * | * | * | * |
| Grand Junction | * | * | * | * | * | * | * |
| Greenfield | * | * | * | * | * | * | * |
| Grinnell | 87.7 | 2.3 | 7.7 | 0.0 | 0.8 | 1.5 | 0.0 |
| Grundy Ctr | * | * | * | * | * | * | * |
| Guthrie Ct | * | * | * | * | * | * | * |
| Hampton | 55.4 | 0.7 | 43.9 | 0.0 | 0.0 | 0.0 | 0.0 |
| Harlan | 85.6 | 0.0 | 12.5 | 1.9 | 0.0 | 0.0 | 0.0 |
| Hawarden | 64.7 | 0.0 | 34.3 | 1.0 | 0.0 | 0.0 | 0.0 |
| Hispanic Cln | * | * | * | * | * | * | * |
| Humboldt | * | * | * | * | * | * | * |
| ILH | 42.7 | 14.4 | 34.2 | 0.3 | 5.4 | 3.1 | 0.0 |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Racial and Ethnic Distribution by Clinic

## Combined 3 Years

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Ida Grove | * | * | * | * | * | * | * |
| Independen | 94.3 | 0.7 | 5.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Indianola | 93.5 | 1.5 | 3.0 | 0.0 | 0.5 | 1.0 | 0.5 |
| Iowa City | 35.9 | 40.4 | 18.4 | 0.1 | 3.2 | 1.9 | 0.0 |
| Iowa Falls | * | * | * | * | * | * | * |
| Jamaica | * | * | * | * | * | * | * |
| Jefferson | * | * | * | * | * | * | * |
| KR | * | * | * | * | * | * | * |
| Keokuk | 87.3 | 5.3 | 3.9 | 0.0 | 0.9 | 2.6 | 0.0 |
| Keosauqua | * | * | * | * | * | * | * |
| Knoxville | 89.8 | 2.3 | 6.1 | 0.4 | 0.8 | 0.8 | 0.0 |
| Lamoni | * | * | * | * | * | * | * |
| Le Mars | 72.8 | 1.2 | 20.7 | 1.8 | 2.4 | 1.2 | 0.0 |
| Lenox | * | * | * | * | * | * | * |
| Leon | * | * | * | * | * | * | * |
| Logan | 93.9 | 0.9 | 5.3 | 0.0 | 0.0 | 0.0 | 0.0 |
| Manchester | * | * | * | * | * | * | * |
| Maquoketa | 85.5 | 1.7 | 1.7 | 0.0 | 9.4 | 1.7 | 0.0 |
| Marengo | * | * | * | * | * | * | * |
| Marion | 81.9 | 8.9 | 6.9 | 0.0 | 0.9 | 1.3 | 0.2 |
| Marshalltown | 39.7 | 3.1 | 46.0 | 0.4 | 9.7 | 1.0 | 0.0 |
| Mason City | 80.2 | 4.7 | 10.1 | 0.3 | 2.6 | 2.1 | 0.0 |
| Monticello | * | * | * | * | * | * | * |
| Mt Ayr | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Racial and Ethnic Distribution by Clinic

## Combined 3 Years

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Mt Pleasan | 77.0 | 6.5 | 12.9 | 0.0 | 2.3 | 1.4 | 0.0 |
| Murray | * | * | * | * | * | * | * |
| Muscatine | 60.6 | 4.8 | 32.8 | 0.0 | 1.8 | 0.1 | 0.0 |
| Nevada | * | * | * | * | * | * | * |
| New Hampto | * | * | * | * | * | * | * |
| Newton | 85.3 | 4.1 | 8.8 | 0.0 | 0.6 | 0.9 | 0.3 |
| North Towne | 64.6 | 17.2 | 7.2 | 0.1 | 6.9 | 3.8 | 0.1 |
| Northwood | * | * | * | * | * | * | * |
| Norwalk | * | * | * | * | * | * | * |
| Oelwein | 87.6 | 3.5 | 8.0 | 0.0 | 0.0 | 0.9 | 0.0 |
| Onawa | * | * | * | * | * | * | * |
| Orange Cty | * | * | * | * | * | * | * |
| Osage | * | * | * | * | * | * | * |
| Osceola | 63.4 | 0.0 | 34.3 | 0.0 | 1.7 | 0.6 | 0.0 |
| Oskaloosa | 88.4 | 3.9 | 6.8 | 0.0 | 0.5 | 0.5 | 0.0 |
| Ottumwa | 73.0 | 7.7 | 15.2 | 0.1 | 2.7 | 1.3 | 0.0 |
| Pella | * | * | * | * | * | * | * |
| Perry | 42.9 | 3.4 | 48.8 | 0.0 | 2.5 | 2.5 | 0.0 |
| Pocahontas | * | * | * | * | * | * | * |
| Pomeroy | * | * | * | * | * | * | * |
| Postville | 35.8 | 15.3 | 48.2 | 0.0 | 0.0 | 0.7 | 0.0 |
| Red Oak | 91.2 | 0.0 | 7.7 | 0.0 | 0.6 | 0.6 | 0.0 |
| Redfield | * | * | * | * | * | * | * |
| Refugee Cln | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Racial and Ethnic Distribution by Clinic

## Combined 3 Years

| Clinic | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Rock Rapid | * | * | * | * | * | * | * |
| Rock Valle | 41.7 | 0.0 | 56.5 | 0.9 | 0.0 | 0.9 | 0.0 |
| Rockwell C | * | * | * | * | * | * | * |
| SDM | 49.0 | 11.7 | 27.0 | 0.3 | 7.5 | 4.5 | 0.0 |
| Sac City | * | * | * | * | * | * | * |
| Sheldon | 56.5 | 1.9 | 38.3 | 0.0 | 2.6 | 0.6 | 0.0 |
| Shenandoah | 88.6 | 0.0 | 11.4 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sibley | * | * | * | * | * | * | * |
| Sidney | * | * | * | * | * | * | * |
| Sigourney | * | * | * | * | * | * | * |
| Sioux Cent | 29.7 | 0.0 | 69.7 | 0.0 | 0.0 | 0.6 | 0.0 |
| Sioux City | 44.0 | 10.3 | 36.7 | 3.3 | 3.3 | 2.4 | 0.0 |
| Southridge | 49.0 | 13.5 | 27.7 | 0.3 | 5.8 | 3.7 | 0.0 |
| Spencer | 82.1 | 0.0 | 14.5 | 0.0 | 2.1 | 1.3 | 0.0 |
| Spirit Lak | * | * | * | * | * | * | * |
| Storm Lake | 25.3 | 4.2 | 52.4 | 0.4 | 17.1 | 0.6 | 0.0 |
| Stuart | * | * | * | * | * | * | * |
| Tama | 41.4 | 1.0 | 39.0 | 16.2 | 0.5 | 1.9 | 0.0 |
| Tama/Settl | * | * | * | * | * | * | * |
| Tipton | * | * | * | * | * | * | * |
| Traer | * | * | * | * | * | * | * |
| UIHC | * | * | * | * | * | * | * |
| Vinton | * | * | * | * | * | * | * |
| WDM | 54.0 | 16.3 | 25.1 | 0.5 | 0.9 | 3.3 | 0.0 |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 3A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Racial and Ethnic Distribution by Clinic

### Combined 3 Years

| **Clinic** | White, Not Hispanic % | Black, Not Hispanic % | Hispanic % | American Indian/Alaska Native % | Asian/Pacific Islander % | Multiple Races % | All Other/Unknown % |
|---|---|---|---|---|---|---|---|
| Wapello | * | * | * | * | * | * | * |
| Washington | 65.7 | 0.7 | 30.6 | 0.0 | 0.0 | 3.0 | 0.0 |
| Waterloo | 57.9 | 22.1 | 10.4 | 0.5 | 4.7 | 4.2 | 0.3 |
| Waukee | 80.7 | 6.4 | 7.3 | 0.0 | 3.7 | 1.8 | 0.0 |
| Waukon | * | * | * | * | * | * | * |
| Waverly | * | * | * | * | * | * | * |
| Wbstr Ct | 67.2 | 0.5 | 29.9 | 0.0 | 2.0 | 0.5 | 0.0 |
| West Union | * | * | * | * | * | * | * |
| Williamsburg | * | * | * | * | * | * | * |
| Winterset | * | * | * | * | * | * | * |
| **Iowa** | **62.3** | **12.3** | **18.4** | **0.5** | **3.7** | **2.6** | **0.1** |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

### Combined 3 Years

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Ackley | * | * | * | * | * | * | * |
| Adel | * | * | * | * | * | * | * |
| Albia | * | * | * | * | * | * | * |
| Algona | 0.0 | 1.8 | 7.9 | 64.0 | 24.6 | 1.8 | 0.0 |
| Allen WHC | 1.1 | 2.9 | 6.9 | 58.3 | 29.6 | 1.3 | 0.0 |
| Allison | * | * | * | * | * | * | * |
| Altoona | 0.0 | 4.3 | 1.7 | 53.8 | 40.2 | 0.0 | 0.0 |
| Ames | 0.0 | 1.2 | 5.4 | 56.2 | 34.6 | 2.5 | 0.0 |
| Anamosa | * | * | * | * | * | * | * |
| Ankeny | 0.0 | 2.3 | 3.8 | 64.7 | 25.6 | 3.8 | 0.0 |
| Atlantic | 0.0 | 1.9 | 11.7 | 61.7 | 24.1 | 0.0 | 0.6 |
| Audubon | * | * | * | * | * | * | * |
| Avoca | * | * | * | * | * | * | * |
| Bedford | * | * | * | * | * | * | * |
| Belle Plaine | * | * | * | * | * | * | * |
| Bloomfield | * | * | * | * | * | * | * |
| Boone | * | * | * | * | * | * | * |
| Brdlwns RP | 0.1 | 2.3 | 6.8 | 58.0 | 29.8 | 3.0 | 0.0 |
| Brooklyn | * | * | * | * | * | * | * |
| Burlington | 0.0 | 4.5 | 10.2 | 62.6 | 21.4 | 1.3 | 0.0 |
| CHC | 0.3 | 3.8 | 6.5 | 66.1 | 22.3 | 1.1 | 0.0 |
| CR Urban | 0.1 | 1.5 | 6.1 | 62.8 | 27.8 | 1.7 | 0.1 |
| Carlisle | * | * | * | * | * | * | * |
| Carroll | 0.0 | 0.0 | 5.4 | 62.9 | 29.9 | 1.8 | 0.0 |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

### Combined 3 Years

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Cedar Falls | * | * | * | * | * | * | * |
| Centervill | * | * | * | * | * | * | * |
| Chariton | * | * | * | * | * | * | * |
| Charles City | 0.0 | 0.0 | 2.5 | 65.3 | 31.2 | 1.0 | 0.0 |
| Cherokee | 0.0 | 0.8 | 10.1 | 66.7 | 20.9 | 1.6 | 0.0 |
| Clarinda | * | * | * | * | * | * | * |
| Clarion | 0.7 | 4.1 | 11.0 | 55.9 | 26.9 | 1.4 | 0.0 |
| Clinton | 0.0 | 2.8 | 7.8 | 65.8 | 22.6 | 0.8 | 0.2 |
| Colfax | * | * | * | * | * | * | * |
| Columbus J | * | * | * | * | * | * | * |
| Corning | * | * | * | * | * | * | * |
| Correctionvl | * | * | * | * | * | * | * |
| Corydon | * | * | * | * | * | * | * |
| Council Blfs | 0.1 | 3.2 | 7.2 | 66.4 | 21.9 | 1.2 | 0.0 |
| Cresco | * | * | * | * | * | * | * |
| Creston | 0.0 | 4.7 | 8.1 | 63.4 | 21.7 | 1.7 | 0.3 |
| Dallas Ctr | * | * | * | * | * | * | * |
| Decorah | * | * | * | * | * | * | * |
| Denison | 0.0 | 3.1 | 7.1 | 58.2 | 29.5 | 2.1 | 0.0 |
| Dewitt | * | * | * | * | * | * | * |
| Dubuque | 0.1 | 3.2 | 8.5 | 63.0 | 23.7 | 1.2 | 0.3 |
| Dyersville | * | * | * | * | * | * | * |
| Eagle Grove | * | * | * | * | * | * | * |
| Edgerton | 0.1 | 3.3 | 9.1 | 62.2 | 24.3 | 1.0 | 0.0 |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

### Combined 3 Years

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Edgerton CHC | * | * | * | * | * | * | * |
| Eldora | * | * | * | * | * | * | * |
| Elkader | 0.0 | 0.8 | 0.8 | 70.9 | 26.0 | 1.6 | 0.0 |
| Emmetsburg | * | * | * | * | * | * | * |
| Esthervill | 0.0 | 0.7 | 7.1 | 68.6 | 22.9 | 0.7 | 0.0 |
| Fairfiel | 0.0 | 1.9 | 1.9 | 68.3 | 26.0 | 1.9 | 0.0 |
| Fayette | * | * | * | * | * | * | * |
| Forest City | 0.0 | 1.5 | 6.1 | 71.0 | 20.6 | 0.8 | 0.0 |
| Fort Dodge | 0.2 | 4.3 | 7.1 | 63.9 | 23.6 | 0.9 | 0.0 |
| Ft Madison | 0.0 | 6.4 | 9.0 | 65.8 | 18.4 | 0.4 | 0.0 |
| Garner | * | * | * | * | * | * | * |
| Glenwood | * | * | * | * | * | * | * |
| Grand Junction | * | * | * | * | * | * | * |
| Greenfield | * | * | * | * | * | * | * |
| Grinnell | 0.0 | 2.3 | 3.8 | 68.5 | 24.6 | 0.8 | 0.0 |
| Grundy Ctr | * | * | * | * | * | * | * |
| Guthrie Ct | * | * | * | * | * | * | * |
| Hampton | 0.7 | 2.2 | 5.8 | 55.4 | 32.4 | 3.6 | 0.0 |
| Harlan | 0.0 | 1.0 | 4.8 | 73.1 | 19.2 | 1.9 | 0.0 |
| Hawarden | 0.0 | 1.0 | 4.9 | 62.7 | 29.4 | 2.0 | 0.0 |
| Hispanic Cln | * | * | * | * | * | * | * |
| Humboldt | * | * | * | * | * | * | * |
| ILH | 0.1 | 3.7 | 7.2 | 57.7 | 29.3 | 1.9 | 0.0 |
| Ida Grove | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

### Combined 3 Years

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Independen | 0.0 | 3.5 | 5.7 | 61.7 | 27.7 | 1.4 | 0.0 |
| Indianola | 0.0 | 1.0 | 4.5 | 68.5 | 25.0 | 1.0 | 0.0 |
| Iowa City | 0.0 | 0.9 | 4.6 | 53.4 | 37.6 | 3.5 | 0.0 |
| Iowa Falls | * | * | * | * | * | * | * |
| Jamaica | * | * | * | * | * | * | * |
| Jefferson | * | * | * | * | * | * | * |
| KR | * | * | * | * | * | * | * |
| Keokuk | 0.0 | 3.5 | 13.6 | 69.3 | 13.6 | 0.0 | 0.0 |
| Keosauqua | * | * | * | * | * | * | * |
| Knoxville | 0.0 | 3.0 | 6.8 | 64.4 | 23.9 | 1.9 | 0.0 |
| Lamoni | * | * | * | * | * | * | * |
| Le Mars | 0.0 | 1.2 | 10.7 | 55.0 | 30.8 | 2.4 | 0.0 |
| Lenox | * | * | * | * | * | * | * |
| Leon | * | * | * | * | * | * | * |
| Logan | 0.0 | 0.0 | 3.5 | 69.3 | 27.2 | 0.0 | 0.0 |
| Manchester | * | * | * | * | * | * | * |
| Maquoketa | 0.0 | 0.9 | 7.7 | 69.2 | 22.2 | 0.0 | 0.0 |
| Marengo | * | * | * | * | * | * | * |
| Marion | 0.4 | 2.2 | 6.3 | 62.0 | 27.5 | 1.6 | 0.0 |
| Marshalltown | 0.1 | 2.6 | 7.9 | 59.0 | 29.1 | 1.3 | 0.0 |
| Mason City | 0.0 | 1.7 | 8.3 | 66.5 | 22.2 | 1.3 | 0.0 |
| Monticello | * | * | * | * | * | * | * |
| Mt Ayr | * | * | * | * | * | * | * |
| Mt Pleasan | 0.0 | 2.3 | 5.1 | 69.6 | 21.7 | 1.4 | 0.0 |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

### Combined 3 Years

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Murray | * | * | * | * | * | * | * |
| Muscatine | 0.1 | 3.0 | 8.2 | 62.2 | 24.3 | 2.2 | 0.0 |
| Nevada | * | * | * | * | * | * | * |
| New Hampto | * | * | * | * | * | * | * |
| Newton | 0.0 | 2.4 | 6.2 | 68.4 | 21.8 | 1.2 | 0.0 |
| North Towne | 0.0 | 1.7 | 6.9 | 64.8 | 25.4 | 1.3 | 0.0 |
| Northwood | * | * | * | * | * | * | * |
| Norwalk | * | * | * | * | * | * | * |
| Oelwein | 0.0 | 2.7 | 8.8 | 61.1 | 26.5 | 0.9 | 0.0 |
| Onawa | * | * | * | * | * | * | * |
| Orange Cty | * | * | * | * | * | * | * |
| Osage | * | * | * | * | * | * | * |
| Osceola | 0.0 | 0.6 | 5.2 | 61.6 | 27.3 | 5.2 | 0.0 |
| Oskaloosa | 0.0 | 2.9 | 8.7 | 68.1 | 16.9 | 2.4 | 1.0 |
| Ottumwa | 0.1 | 2.7 | 8.1 | 63.3 | 24.5 | 1.3 | 0.1 |
| Pella | * | * | * | * | * | * | * |
| Perry | 0.5 | 4.4 | 5.4 | 56.7 | 28.1 | 4.9 | 0.0 |
| Pocahontas | * | * | * | * | * | * | * |
| Pomeroy | * | * | * | * | * | * | * |
| Postville | 0.0 | 2.2 | 5.1 | 55.5 | 36.5 | 0.7 | 0.0 |
| Red Oak | 0.0 | 1.7 | 11.0 | 64.1 | 21.5 | 1.7 | 0.0 |
| Redfield | * | * | * | * | * | * | * |
| Refugee Cln | * | * | * | * | * | * | * |
| Rock Rapid | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

### Combined 3 Years

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Rock Valle | 0.0 | 1.9 | 8.3 | 62.0 | 23.1 | 2.8 | 1.9 |
| Rockwell C | * | * | * | * | * | * | * |
| SDM | 0.0 | 3.1 | 8.1 | 61.0 | 24.8 | 3.1 | 0.0 |
| Sac City | * | * | * | * | * | * | * |
| Sheldon | 0.0 | 3.9 | 6.5 | 58.4 | 29.9 | 1.3 | 0.0 |
| Shenandoah | 0.0 | 3.8 | 6.7 | 69.5 | 19.0 | 1.0 | 0.0 |
| Sibley | * | * | * | * | * | * | * |
| Sidney | * | * | * | * | * | * | * |
| Sigourney | * | * | * | * | * | * | * |
| Sioux Cent | 0.6 | 4.8 | 9.7 | 48.5 | 33.3 | 2.4 | 0.6 |
| Sioux City | 0.0 | 2.1 | 6.7 | 61.7 | 27.5 | 1.9 | 0.0 |
| Southridge | 0.3 | 2.6 | 6.3 | 59.1 | 29.1 | 2.6 | 0.0 |
| Spencer | 0.0 | 1.3 | 5.1 | 71.5 | 20.9 | 1.3 | 0.0 |
| Spirit Lak | * | * | * | * | * | * | * |
| Storm Lake | 0.0 | 2.6 | 6.6 | 62.9 | 25.1 | 2.8 | 0.0 |
| Stuart | * | * | * | * | * | * | * |
| Tama | 0.0 | 1.9 | 5.7 | 62.9 | 29.0 | 0.5 | 0.0 |
| Tama/Settl | * | * | * | * | * | * | * |
| Tipton | * | * | * | * | * | * | * |
| Traer | * | * | * | * | * | * | * |
| UIHC | * | * | * | * | * | * | * |
| Vinton | * | * | * | * | * | * | * |
| WDM | 0.0 | 0.0 | 5.6 | 59.5 | 32.1 | 2.8 | 0.0 |
| Wapello | * | * | * | * | * | * | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 4A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Age Distribution by Clinic

### Combined 3 Years

| Clinic | < 15 Years % | 15 - 17 Years % | 18 - 19 Years % | 20 - 29 Years % | 30 - 39 Years % | >= 40 Years % | Unknown % |
|---|---|---|---|---|---|---|---|
| Washington | 0.0 | 1.5 | 4.5 | 62.7 | 29.9 | 0.7 | 0.7 |
| Waterloo | 0.1 | 1.6 | 5.1 | 66.4 | 25.8 | 1.0 | 0.0 |
| Waukee | 0.0 | 3.7 | 2.8 | 54.1 | 37.6 | 1.8 | 0.0 |
| Waukon | * | * | * | * | * | * | * |
| Waverly | * | * | * | * | * | * | * |
| Wbstr Ct | 0.0 | 3.9 | 11.3 | 57.8 | 22.5 | 4.4 | 0.0 |
| West Union | * | * | * | * | * | * | * |
| Williamsburg | * | * | * | * | * | * | * |
| Winterset | * | * | * | * | * | * | * |
| **Iowa** | **0.1** | **2.4** | **6.8** | **62.1** | **26.8** | **1.8** | **0.0** |

(1) Reporting period is January 1 through December 31.
(2) Excludes data with errors.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Ackley | | * | * | | * | * | | * | | * | | * | | * |
| Adel | | * | * | | * | * | | * | | * | | * | | * |
| Adel | | * | * | | * | * | | * | | * | | * | | * |
| Adel | | * | * | | * | * | | * | | * | | * | | * |
| Albia | | * | * | | * | * | | * | | * | | * | | * |
| Albia | | * | * | | * | * | | * | | * | | * | | * |
| Algona | | * | * | | * | * | | * | | * | | * | | * |
| Algona | | * | * | | * | * | | * | | * | | * | | * |
| Allen WHC | | * | * | | * | * | | * | | * | | * | | * |
| Allen WHC | | 3.1 (16) | 60.2 (30) | | 19.0 (5) | 45.7 (18) | | * | | * | | * | | * |
| Allison | | * | * | | * | * | | * | | * | | * | | * |
| Allison | | * | * | | * | * | | * | | * | | * | | * |
| Altoona | | * | * | | * | * | | * | | * | | * | | * |
| Altoona | | * | * | | * | * | | * | | * | | * | | * |
| Ames | | 5.4 (54) | 54.7 (4) | | 21.2 (15) | 42.7 (8) | | 16.8 (13) | | 15.8 (7) | | * | | * |
| Ames | | 3.8 (27) | 54.3 (2) | | * | * | | * | | * | | * | | * |
| Anamosa | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Anamosa | | * | * | | * | * | | * | | * | | * | | * |
| Anamosa | | * | * | | * | * | | * | | * | | * | | * |
| Ankeny | | * | * | | * | * | | * | | * | | * | | * |
| Ankeny | | * | * | | * | * | | * | | * | | * | | * |
| Atlantic | | * | * | | * | * | | * | | * | | * | | * |
| Atlantic | | 8.9 (59) | 57.4 (10) | | * | * | | * | | * | | * | | * |
| Audubon | | * | * | | * | * | | * | | * | | * | | * |
| Audubon | | * | * | | * | * | | * | | * | | * | | * |
| Avoca | | * | * | | * | * | | * | | * | | * | | * |
| Avoca | | * | * | | * | * | | * | | * | | * | | * |
| Bedford | | * | * | | * | * | | * | | * | | * | | * |
| Bedford | | * | * | | * | * | | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | * | | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | * | | * | | * | | * | | * |
| Bloomfield | | * | * | | * | * | | * | | * | | * | | * |
| Bloomfield | | * | * | | * | * | | * | | * | | * | | * |
| Boone | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Boone | | * | * | | * | * | | * | | * | | * | | * |
| Brdlwns RP | | 4.0 (36) | 60.3 (31) | | 25.7 (40) | 44.8 (13) | | * | | * | | 12.7 (6) | | * |
| Brdlwns RP | | 3.8 (27) | 55.0 (5) | | 25.4 (39) | 41.6 (7) | | 3.0 (4) | | 3.5 (1) | | 100.0 (8) | | * |
| Brooklyn | | * | * | | * | * | | * | | * | | * | | * |
| Brooklyn | | * | * | | * | * | | * | | * | | * | | * |
| Burlington | | 6.5 (57) | 58.2 (17) | | 16.3 (2) | 53.1 (44) | | 6.5 (8) | | * | | * | | * |
| Burlington | | * | * | | * | * | | * | | * | | * | | * |
| CHC | | 5.2 (52) | 62.1 (41) | | * | * | | * | | * | | * | | * |
| CHC | | 4.7 (46) | 65.7 (56) | | 21.2 (15) | 48.7 (31) | | * | | * | | * | | * |
| CR Urban | | 3.7 (26) | 62.6 (46) | | 23.7 (32) | 45.6 (17) | | * | | * | | 8.4 (3) | | 16.9 (4) |
| CR Urban | | 4.0 (36) | 59.2 (22) | | 21.1 (14) | 48.7 (31) | | 32.6 (16) | | 30.8 (11) | | 100.0 (8) | | * |
| Carlisle | | * | * | | * | * | | * | | * | | * | | * |
| Carlisle | | * | * | | * | * | | * | | * | | * | | * |
| Carroll | | 1.3 (3) | 63.8 (50) | | 16.7 (3) | 51.4 (41) | | * | | * | | * | | * |
| Carroll | | * | * | | * | * | | * | | * | | * | | * |
| Cedar Falls | | * | * | | * | * | | * | | * | | * | | * |
| Cedar Falls | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Centervill | | * | * | | * | * | | * | | * | | * | | * |
| Centervill | | * | * | | * | * | | * | | * | | * | | * |
| Chariton | | * | * | | * | * | | * | | * | | * | | * |
| Chariton | | * | * | | * | * | | * | | * | | * | | * |
| Chariton | | * | * | | * | * | | * | | * | | * | | * |
| Charles City | | 4.0 (36) | 59.2 (22) | | 22.3 (25) | 46.3 (22) | | * | | * | | * | | * |
| Charles City | | * | * | | * | * | | * | | * | | * | | * |
| Cherokee | | * | * | | * | * | | * | | * | | * | | * |
| Cherokee | | * | * | | * | * | | * | | * | | * | | * |
| Clarinda | | * | * | | * | * | | * | | * | | * | | * |
| Clarinda | | * | * | | * | * | | * | | * | | * | | * |
| Clarion | | * | * | | * | * | | * | | * | | * | | * |
| Clarion | | * | * | | * | * | | * | | * | | * | | * |
| Clinton | | * | * | | * | * | | * | | * | | * | | * |
| Clinton | | 4.9 (47) | 61.8 (40) | | 20.5 (11) | 53.4 (46) | | 30.4 (15) | | * | | * | | * |
| Colfax | | * | * | | * | * | | * | | * | | * | | * |
| Colfax | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Columbus J | | * | * | | * | * | | * | | * | | * | | * |
| Columbus J | | * | * | | * | * | | * | | * | | * | | * |
| Corning | | * | * | | * | * | | * | | * | | * | | * |
| Corning | | * | * | | * | * | | * | | * | | * | | * |
| Correctionvl | | * | * | | * | * | | * | | * | | * | | * |
| Correctionvl | | * | * | | * | * | | * | | * | | * | | * |
| Corydon | | * | * | | * | * | | * | | * | | * | | * |
| Corydon | | * | * | | * | * | | * | | * | | * | | * |
| Council Blfs | | 3.9 (33) | 59.9 (26) | | 21.4 (19) | 48.1 (27) | | 8.6 (9) | | 3.9 (2) | | * | | * |
| Council Blfs | | 3.3 (21) | 60.3 (31) | | 15.4 (1) | 49.8 (37) | | * | | * | | 11.1 (5) | | 13.3 (2) |
| Cresco | | * | * | | * | * | | * | | * | | * | | * |
| Cresco | | * | * | | * | * | | * | | * | | * | | * |
| Creston | | 3.0 (14) | 62.7 (47) | | 19.0 (5) | 53.2 (45) | | * | | * | | * | | * |
| Creston | | * | * | | * | * | | * | | * | | * | | * |
| Dallas Ctr | | * | * | | * | * | | * | | * | | * | | * |
| Decorah | | * | * | | * | * | | * | | * | | * | | * |
| Decorah | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Denison | | 2.5 (9) | 60.7 (34) | | 26.5 (44) | 37.6 (1) | | 3.9 (5) | | * | | * | | * |
| Denison | | * | * | | * | * | | * | | * | | * | | * |
| Dewitt | | * | * | | * | * | | * | | * | | * | | * |
| Dewitt | | * | * | | * | * | | * | | * | | * | | * |
| Dubuque | | 5.1 (48) | 56.6 (7) | | * | * | | * | | * | | 8.1 (2) | | 4.5 (1) |
| Dubuque | | 4.2 (42) | 58.0 (14) | | 20.2 (9) | 49.9 (38) | | 47.8 (20) | | * | | * | | * |
| Dyersville | | * | * | | * | * | | * | | * | | * | | * |
| Dyersville | | * | * | | * | * | | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | * | | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | * | | * | | * | | * | | * |
| Edgerton | | 5.1 (48) | 57.3 (8) | | 21.6 (22) | 45.8 (20) | | 36.9 (18) | | 21.8 (8) | | * | | * |
| Edgerton | | 3.9 (33) | 62.4 (44) | | 21.0 (13) | 52.0 (43) | | * | | * | | 8.0 (1) | | 20.5 (6) |
| Edgerton CHC | | * | * | | * | * | | * | | * | | * | | * |
| Eldora | | * | * | | * | * | | * | | * | | * | | * |
| Eldora | | * | * | | * | * | | * | | * | | * | | * |
| Elkader | | * | * | | * | * | | * | | * | | * | | * |
| Elkader | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Emmetsburg | | * | * | | * | * | | * | | * | | * | | * |
| Emmetsburg | | * | * | | * | * | | * | | * | | * | | * |
| Esthervill | | * | * | | * | * | | * | | * | | * | | * |
| Esthervill | | * | * | | * | * | | * | | * | | * | | * |
| Fairfiel | | * | * | | * | * | | * | | * | | * | | * |
| Fairfiel | | * | * | | * | * | | * | | * | | * | | * |
| Fayette | | * | * | | * | * | | * | | * | | * | | * |
| Fayette | | * | * | | * | * | | * | | * | | * | | * |
| Forest City | | * | * | | * | * | | * | | * | | * | | * |
| Forest City | | * | * | | * | * | | * | | * | | * | | * |
| Fort Dodge | | 3.8 (27) | 60.6 (33) | | * | * | | * | | * | | * | | * |
| Fort Dodge | | 2.8 (11) | 63.7 (49) | | 21.2 (15) | 47.6 (25) | | 12.4 (11) | | * | | * | | * |
| Ft Madison | | 3.4 (22) | 66.9 (58) | | 21.7 (23) | 48.6 (29) | | * | | * | | * | | * |
| Ft Madison | | * | * | | * | * | | * | | * | | * | | * |
| Garner | | * | * | | * | * | | * | | * | | * | | * |
| Garner | | * | * | | * | * | | * | | * | | * | | * |
| Glenwood | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Glenwood | | * | * | | * | * | | * | | * | | * | | * |
| Grand Junction | | * | * | | * | * | | * | | * | | * | | * |
| Greenfield | | * | * | | * | * | | * | | * | | * | | * |
| Greenfield | | * | * | | * | * | | * | | * | | * | | * |
| Grinnell | | * | * | | * | * | | * | | * | | * | | * |
| Grinnell | | * | * | | * | * | | * | | * | | * | | * |
| Grundy Ctr | | * | * | | * | * | | * | | * | | * | | * |
| Grundy Ctr | | * | * | | * | * | | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | * | | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | * | | * | | * | | * | | * |
| Hampton | | * | * | | * | * | | * | | * | | * | | * |
| Hampton | | * | * | | * | * | | * | | * | | * | | * |
| Harlan | | * | * | | * | * | | * | | * | | * | | * |
| Harlan | | * | * | | * | * | | * | | * | | * | | * |
| Hawarden | | * | * | | * | * | | * | | * | | * | | * |
| Hawarden | | * | * | | * | * | | * | | * | | * | | * |
| Hispanic Cln | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Hispanic Cln | | * | * | | * | * | | * | | * | | * | | * |
| Humboldt | | * | * | | * | * | | * | | * | | * | | * |
| Humboldt | | * | * | | * | * | | * | | * | | * | | * |
| ILH | | 5.1 (48) | 58.0 (14) | | 25.8 (41) | 40.5 (4) | | 2.2 (2) | | 5.1 (3) | | * | | * |
| ILH | | 4.2 (42) | 65.1 (54) | | * | * | | * | | * | | * | | * |
| Ida Grove | | * | * | | * | * | | * | | * | | * | | * |
| Ida Grove | | * | * | | * | * | | * | | * | | * | | * |
| Independen | | * | * | | * | * | | * | | * | | * | | * |
| Independen | | * | * | | * | * | | * | | * | | * | | * |
| Indianola | | * | * | | * | * | | * | | * | | * | | * |
| Indianola | | 2.9 (13) | 62.5 (45) | | 20.3 (10) | 47.7 (26) | | * | | * | | * | | * |
| Iowa City | | 3.2 (17) | 57.4 (10) | | * | * | | * | | * | | * | | * |
| Iowa City | | 3.8 (27) | 55.3 (6) | | 25.8 (41) | 40.5 (4) | | 37.1 (19) | | 27.2 (10) | | * | | * |
| Iowa Falls | | * | * | | * | * | | * | | * | | * | | * |
| Iowa Falls | | * | * | | * | * | | * | | * | | * | | * |
| Jamaica | | * | * | | * | * | | * | | * | | * | | * |
| Jefferson | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Jefferson | | * | * | | * | * | | * | | * | | * | | * |
| KR | | * | * | | * | * | | * | | * | | * | | * |
| KR | | * | * | | * | * | | * | | * | | * | | * |
| Keokuk | | * | * | | * | * | | * | | * | | * | | * |
| Keokuk | | 5.3 (53) | 59.9 (26) | | 21.5 (21) | 45.1 (14) | | * | | * | | * | | * |
| Keosauqua | | * | * | | * | * | | * | | * | | * | | * |
| Keosauqua | | * | * | | * | * | | * | | * | | * | | * |
| Knoxville | | * | * | | * | * | | * | | * | | * | | * |
| Knoxville | | 5.1 (48) | 59.5 (24) | | 22.0 (24) | 49.3 (35) | | * | | * | | * | | * |
| Lamoni | | * | * | | * | * | | * | | * | | * | | * |
| Lamoni | | * | * | | * | * | | * | | * | | * | | * |
| Le Mars | | 6.6 (58) | 62.3 (43) | | 23.1 (29) | 49.0 (34) | | * | | * | | * | | * |
| Le Mars | | * | * | | * | * | | * | | * | | * | | * |
| Lenox | | * | * | | * | * | | * | | * | | * | | * |
| Lenox | | * | * | | * | * | | * | | * | | * | | * |
| Leon | | * | * | | * | * | | * | | * | | * | | * |
| Leon | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Logan | | * | * | | * | * | | * | | * | | * | | * |
| Logan | | * | * | | * | * | | * | | * | | * | | * |
| Manchester | | * | * | | * | * | | * | | * | | * | | * |
| Manchester | | * | * | | * | * | | * | | * | | * | | * |
| Maquoketa | | * | * | | * | * | | * | | * | | * | | * |
| Maquoketa | | * | * | | * | * | | * | | * | | * | | * |
| Marengo | | * | * | | * | * | | * | | * | | * | | * |
| Marengo | | * | * | | * | * | | * | | * | | * | | * |
| Marion | | 1.8 (5) | 63.8 (50) | | 19.9 (8) | 51.4 (41) | | * | | * | | * | | * |
| Marion | | * | * | | * | * | | * | | * | | * | | * |
| Marshalltown | | 3.8 (27) | 59.5 (24) | | * | * | | * | | * | | * | | * |
| Marshalltown | | 4.6 (45) | 60.1 (29) | | 25.8 (41) | 39.2 (2) | | 15.2 (12) | | 14.6 (6) | | * | | * |
| Mason City | | 3.2 (17) | 63.6 (48) | | 18.8 (4) | 50.4 (40) | | 4.4 (6) | | * | | * | | * |
| Mason City | | 2.3 (8) | 57.4 (10) | | * | * | | * | | * | | * | | * |
| Monticello | | * | * | | * | * | | * | | * | | * | | * |
| Monticello | | * | * | | * | * | | * | | * | | * | | * |
| Monticello | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Mt Ayr | | * | * | | * | * | | * | | * | | * | | * |
| Mt Ayr | | * | * | | * | * | | * | | * | | * | | * |
| Mt Pleasan | | 2.2 (7) | 59.0 (20) | | 22.6 (26) | 48.9 (33) | | * | | * | | * | | * |
| Mt Pleasan | | * | * | | * | * | | * | | * | | * | | * |
| Murray | | * | * | | * | * | | * | | * | | * | | * |
| Muscatine | | 6.1 (56) | 60.9 (37) | | * | * | | * | | * | | * | | * |
| Muscatine | | 3.6 (25) | 59.9 (26) | | 20.5 (11) | 47.5 (24) | | 1.9 (1) | | * | | * | | * |
| Nevada | | * | * | | * | * | | * | | * | | * | | * |
| Nevada | | * | * | | * | * | | * | | * | | * | | * |
| New Hampto | | * | * | | * | * | | * | | * | | * | | * |
| New Hampto | | * | * | | * | * | | * | | * | | * | | * |
| Newton | | * | * | | * | * | | * | | * | | * | | * |
| Newton | | 3.2 (17) | 59.1 (21) | | 23.2 (30) | 46.4 (23) | | * | | * | | * | | * |
| North Towne | | 2.0 (6) | 58.8 (19) | | * | * | | * | | * | | * | | * |
| North Towne | | 3.0 (14) | 64.0 (52) | | 19.7 (7) | 49.4 (36) | | 36.4 (17) | | 36.5 (12) | | * | | * |
| Northwood | | * | * | | * | * | | * | | * | | * | | * |
| Northwood | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Norwalk | | * | * | | * | * | | * | | * | | * | | * |
| Oelwein | | * | * | | * | * | | * | | * | | * | | * |
| Oelwein | | * | * | | * | * | | * | | * | | * | | * |
| Onawa | | * | * | | * | * | | * | | * | | * | | * |
| Onawa | | * | * | | * | * | | * | | * | | * | | * |
| Orange Cty | | * | * | | * | * | | * | | * | | * | | * |
| Orange Cty | | * | * | | * | * | | * | | * | | * | | * |
| Osage | | * | * | | * | * | | * | | * | | * | | * |
| Osage | | * | * | | * | * | | * | | * | | * | | * |
| Osceola | | * | * | | * | * | | * | | * | | * | | * |
| Osceola | | * | * | | * | * | | * | | * | | * | | * |
| Oskaloosa | | * | * | | * | * | | * | | * | | * | | * |
| Oskaloosa | | * | * | | * | * | | * | | * | | * | | * |
| Oskaloosa | | * | * | | * | * | | * | | * | | * | | * |
| Ottumwa | | 4.3 (44) | 66.1 (57) | | * | * | | * | | * | | * | | * |
| Ottumwa | | 4.1 (40) | 60.8 (35) | | 23.3 (31) | 45.7 (18) | | 22.9 (14) | | 22.8 (9) | | * | | * |
| Pella | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Pella | | * | * | | * | * | | * | | * | | * | | * |
| Perry | | * | * | | * | * | | * | | * | | * | | * |
| Perry | | * | * | | * | * | | * | | * | | * | | * |
| Perry | | * | * | | * | * | | * | | * | | * | | * |
| Pocahontas | | * | * | | * | * | | * | | * | | * | | * |
| Pocahontas | | * | * | | * | * | | * | | * | | * | | * |
| Pocahontas | | * | * | | * | * | | * | | * | | * | | * |
| Pomeroy | | * | * | | * | * | | * | | * | | * | | * |
| Postville | | * | * | | * | * | | * | | * | | * | | * |
| Postville | | * | * | | * | * | | * | | * | | * | | * |
| Red Oak | | 5.8 (55) | 64.5 (53) | | 24.6 (35) | 50.0 (39) | | * | | * | | * | | * |
| Red Oak | | * | * | | * | * | | * | | * | | * | | * |
| Redfield | | * | * | | * | * | | * | | * | | * | | * |
| Refugee Cln | | * | * | | * | * | | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | * | | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | * | | * | | * | | * | | * |
| Rock Valle | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Rock Valle | | * | * | | * | * | | * | | * | | * | | * |
| Rockwell C | | * | * | | * | * | | * | | * | | * | | * |
| Rockwell C | | * | * | | * | * | | * | | * | | * | | * |
| SDM | | 3.8 (27) | 57.4 (10) | | 23.7 (32) | 45.5 (16) | | * | | * | | * | | * |
| Sac City | | * | * | | * | * | | * | | * | | * | | * |
| Sac City | | * | * | | * | * | | * | | * | | * | | * |
| Sheldon | | * | * | | * | * | | * | | * | | * | | * |
| Sheldon | | * | * | | * | * | | * | | * | | * | | * |
| Shenandoah | | * | * | | * | * | | * | | * | | * | | * |
| Shenandoah | | * | * | | * | * | | * | | * | | * | | * |
| Sibley | | * | * | | * | * | | * | | * | | * | | * |
| Sibley | | * | * | | * | * | | * | | * | | * | | * |
| Sidney | | * | * | | * | * | | * | | * | | * | | * |
| Sidney | | * | * | | * | * | | * | | * | | * | | * |
| Sigourney | | * | * | | * | * | | * | | * | | * | | * |
| Sigourney | | * | * | | * | * | | * | | * | | * | | * |
| Sioux Cent | | 0.0 (1) | 52.5 (1) | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Sioux Cent | | * | * | | * | * | | * | | * | | * | | * |
| Sioux City | | 3.9 (33) | 58.1 (16) | | 21.3 (18) | 48.6 (29) | | * | | * | | 19.3 (7) | | 17.2 (5) |
| Sioux City | | 3.4 (22) | 60.8 (35) | | 21.4 (19) | 43.5 (9) | | 5.6 (7) | | 7.4 (4) | | 100.0 (8) | | * |
| Southridge | | 3.5 (24) | 54.6 (3) | | 24.7 (36) | 44.7 (11) | | * | | * | | * | | * |
| Spencer | | * | * | | * | * | | * | | * | | * | | * |
| Spencer | | 1.2 (2) | 62.1 (41) | | 29.0 (46) | 40.6 (6) | | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | * | | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | * | | * | | * | | * | | * |
| Storm Lake | | 4.1 (40) | 58.3 (18) | | 25.2 (37) | 39.9 (3) | | 2.4 (3) | | * | | * | | * |
| Storm Lake | | * | * | | * | * | | * | | * | | * | | * |
| Stuart | | * | * | | * | * | | * | | * | | * | | * |
| Stuart | | * | * | | * | * | | * | | * | | * | | * |
| Tama | | 2.8 (11) | 65.3 (55) | | 25.2 (37) | 48.1 (27) | | * | | * | | * | | * |
| Tama | | * | * | | * | * | | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | * | | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | * | | * | | * | | * | | * |
| Tipton | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Tipton | | * | * | | * | * | | * | | * | | * | | * |
| Traer | | * | * | | * | * | | * | | * | | * | | * |
| Traer | | * | * | | * | * | | * | | * | | * | | * |
| UIHC | | * | * | | * | * | | * | | * | | * | | * |
| Vinton | | * | * | | * | * | | * | | * | | * | | * |
| Vinton | | * | * | | * | * | | * | | * | | * | | * |
| WDM | | 1.4 (4) | 61.2 (38) | | 22.7 (27) | 44.7 (11) | | * | | * | | * | | * |
| WDM | | * | * | | * | * | | * | | * | | * | | * |
| Wapello | | * | * | | * | * | | * | | * | | * | | * |
| Wapello | | * | * | | * | * | | * | | * | | * | | * |
| Washington | | * | * | | * | * | | * | | * | | * | | * |
| Washington | | * | * | | * | * | | * | | * | | * | | * |
| Waterloo | | 2.5 (9) | 67.2 (59) | | 28.4 (45) | 44.4 (10) | | * | | * | | 10.1 (4) | | 15.5 (3) |
| Waterloo | | 3.2 (17) | 61.5 (39) | | 22.7 (27) | 46.2 (21) | | 9.2 (10) | | 7.5 (5) | | * | | * |
| Waukee | | * | * | | * | * | | * | | * | | * | | * |
| Waukee | | * | * | | * | * | | * | | * | | * | | * |
| Waukee | | * | * | | * | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 5A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Health Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Health Indicators** (2,3)

| Clinic | Prepregnancy BMI (4) | | | Weight Gain (5) | | | Anemia (Low Hb/Hct) (6) | | | | Gestational Diabetes During Pregnancy | | Hypertension During Pregnancy | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Under-weight | Over-weight | | < Ideal | > Ideal | 3rd Trimester | | Postpartum | | | | | |
| | Number | % (Rank) | % (Rank) | Number | % (Rank) | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) | Number | % (Rank) |
| Waukon | | * | * | | * | * | | * | | * | | * | | * |
| Waukon | | * | * | | * | * | | * | | * | | * | | * |
| Waverly | | * | * | | * | * | | * | | * | | * | | * |
| Waverly | | * | * | | * | * | | * | | * | | * | | * |
| Wbstr Ct | | 4.0 (36) | 57.3 (8) | | 23.9 (34) | 45.3 (15) | | * | | * | | * | | * |
| Wbstr Ct | | * | * | | * | * | | * | | * | | * | | * |
| West Union | | * | * | | * | * | | * | | * | | * | | * |
| Williamsburg | | * | * | | * | * | | * | | * | | * | | * |
| Winterset | | * | * | | * | * | | * | | * | | * | | * |
| Winterset | | * | * | | * | * | | * | | * | | * | | * |
| **Iowa** | | **3.8** | **60.1** | | **22.7** | **45.7** | | **15.5** | | **15.4** | | **48.2** | | **35.0** |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ackley | | * | * | | * | | * | | * |
| Adel | | * | * | | * | | * | | * |
| Adel | | * | * | | * | | * | | * |
| Adel | | * | * | | * | | * | | * |
| Albia | | * | * | | * | | * | | * |
| Albia | | * | * | | * | | * | | * |
| Algona | | * | * | | * | | * | | * |
| Algona | | * | * | | * | | * | | * |
| Allen WHC | | 42.2 (3) | 23.6 (36) | | 4.3 (21) | | 18.5 (22) | | 72.9 (41) |
| Allen WHC | | * | * | | 63.1 (2) | | * | | * |
| Allison | | * | * | | * | | * | | * |
| Allison | | * | * | | * | | * | | * |
| Altoona | | * | * | | * | | * | | * |
| Altoona | | * | * | | * | | * | | * |
| Ames | | * | * | | 53.3 (6) | | * | | * |
| Ames | | 29.6 (27) | 15.3 (28) | | 1.5 (39) | | 21.7 (16) | | 71.7 (42) |
| Anamosa | | * | * | | * | | * | | * |
| Anamosa | | * | * | | * | | * | | * |
| Anamosa | | * | * | | * | | * | | * |
| Ankeny | | * | * | | * | | * | | * |
| Ankeny | | * | * | | 1.9 (34) | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Atlantic | | * | * | | * | | * | | * |
| Atlantic | | * | * | | 0.8 (54) | | * | | * |
| Audubon | | * | * | | * | | * | | * |
| Audubon | | * | * | | * | | * | | * |
| Avoca | | * | * | | * | | * | | * |
| Avoca | | * | * | | * | | * | | * |
| Bedford | | * | * | | * | | * | | * |
| Bedford | | * | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | | * | | * |
| Bloomfield | | * | * | | * | | * | | * |
| Bloomfield | | * | * | | * | | * | | * |
| Boone | | * | * | | * | | * | | * |
| Boone | | * | * | | * | | * | | * |
| Brdlwns RP | | 5.1 (46) | 94.9 (46) | | 42.5 (15) | | 25.2 (6) | | 96.8 (5) |
| Brdlwns RP | | 29.2 (28) | 11.3 (20) | | 1.3 (43) | | 13.7 (39) | | 62.1 (44) |
| Brooklyn | | * | * | | * | | * | | * |
| Brooklyn | | * | * | | * | | * | | * |
| Burlington | | * | * | | * | | * | | * |
| Burlington | | 30.4 (23) | 8.4 (10) | | 0.5 (61) | | 16.5 (30) | | 89.5 (17) |
| CHC | | * | * | | 48.7 (10) | | 10.3 (45) | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| CHC | | 33.2 (16) | 23.6 (36) | | 4.2 (22) | | 18.1 (23) | | 55.4 (47) |
| CR Urban | | 22.1 (37) | 9.1 (12) | | 1.5 (39) | | 13.1 (42) | | 77.7 (33) |
| CR Urban | | 19.7 (39) | 73.8 (41) | | 41.8 (16) | | 8.6 (46) | | 97.0 (4) |
| Carlisle | | * | * | | * | | * | | * |
| Carlisle | | * | * | | * | | * | | * |
| Carroll | | * | * | | * | | * | | * |
| Carroll | | 23.3 (36) | 6.0 (3) | | 0.5 (61) | | * | | * |
| Cedar Falls | | * | * | | * | | * | | * |
| Cedar Falls | | * | * | | * | | * | | * |
| Centervill | | * | * | | * | | * | | * |
| Centervill | | * | * | | * | | * | | * |
| Chariton | | * | * | | * | | * | | * |
| Chariton | | * | * | | * | | * | | * |
| Chariton | | * | * | | * | | * | | * |
| Charles City | | 45.2 (1) | 17.7 (31) | | 0.7 (56) | | 26.9 (5) | | 82.9 (27) |
| Charles City | | * | * | | * | | * | | * |
| Cherokee | | * | * | | * | | * | | * |
| Cherokee | | * | * | | * | | * | | * |
| Clarinda | | * | * | | * | | * | | * |
| Clarinda | | * | * | | * | | * | | * |
| Clarion | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Clarion | | * | * | | 5.7 (19) | | * | | * |
| Clinton | | 35.6 (9) | 7.9 (8) | | 1.5 (39) | | 23.3 (10) | | 84.9 (25) |
| Clinton | | * | * | | * | | * | | * |
| Colfax | | * | * | | * | | * | | * |
| Colfax | | * | * | | * | | * | | * |
| Columbus J | | * | * | | * | | * | | * |
| Columbus J | | * | * | | * | | * | | * |
| Corning | | * | * | | * | | * | | * |
| Corning | | * | * | | * | | * | | * |
| Correctionvl | | * | * | | * | | * | | * |
| Correctionvl | | * | * | | * | | * | | * |
| Corydon | | * | * | | * | | * | | * |
| Corydon | | * | * | | * | | * | | * |
| Council Blfs | | 10.7 (43) | 88.4 (44) | | 44.7 (14) | | 15.0 (35) | | 98.7 (1) |
| Council Blfs | | 40.5 (5) | 6.6 (5) | | 1.2 (45) | | 18.6 (21) | | 77.4 (34) |
| Cresco | | * | * | | * | | * | | * |
| Cresco | | * | * | | * | | * | | * |
| Creston | | 44.3 (2) | 7.1 (7) | | 2.8 (24) | | * | | * |
| Creston | | * | * | | * | | * | | * |
| Dallas Ctr | | * | * | | * | | * | | * |
| Decorah | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Decorah | | * | * | | * | | * | | * |
| Denison | | * | * | | * | | * | | * |
| Denison | | 31.0 (20) | 7.9 (8) | | 0.6 (58) | | 29.6 (3) | | 57.6 (46) |
| Dewitt | | * | * | | * | | * | | * |
| Dewitt | | * | * | | * | | * | | * |
| Dubuque | | 10.3 (44) | 85.5 (43) | | 47.4 (11) | | 17.1 (26) | | 96.6 (6) |
| Dubuque | | 34.3 (14) | 15.5 (29) | | 3.7 (23) | | 15.6 (34) | | 87.8 (19) |
| Dyersville | | * | * | | * | | * | | * |
| Dyersville | | * | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | | * | | * |
| Edgerton | | 35.3 (11) | 12.9 (23) | | 2.0 (33) | | 17.9 (24) | | 48.9 (48) |
| Edgerton | | 14.7 (42) | 82.2 (42) | | 47.3 (12) | | 13.5 (40) | | 98.1 (2) |
| Edgerton CHC | | * | * | | * | | * | | * |
| Eldora | | * | * | | * | | * | | * |
| Eldora | | * | * | | * | | * | | * |
| Elkader | | * | * | | * | | * | | * |
| Elkader | | * | * | | * | | * | | * |
| Emmetsburg | | * | * | | * | | * | | * |
| Emmetsburg | | * | * | | * | | * | | * |
| Esthervill | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Esthervill | | * | * | | 0.0 (63) | | * | | 92.0 (10) |
| Fairfiel | | * | * | | * | | * | | * |
| Fairfiel | | * | * | | * | | * | | * |
| Fayette | | * | * | | * | | * | | * |
| Fayette | | * | * | | * | | * | | * |
| Forest City | | * | * | | * | | * | | * |
| Forest City | | * | * | | * | | * | | * |
| Fort Dodge | | * | * | | 49.5 (9) | | 15.7 (33) | | * |
| Fort Dodge | | 29.7 (26) | 24.1 (38) | | 2.8 (24) | | 16.8 (29) | | 75.9 (38) |
| Ft Madison | | 28.9 (30) | 4.9 (2) | | 0.6 (58) | | 19.1 (18) | | 89.4 (18) |
| Ft Madison | | * | * | | * | | * | | * |
| Garner | | * | * | | * | | * | | * |
| Garner | | * | * | | * | | * | | * |
| Glenwood | | * | * | | * | | * | | * |
| Glenwood | | * | * | | * | | * | | * |
| Grand Junction | | * | * | | * | | * | | * |
| Greenfield | | * | * | | * | | * | | * |
| Greenfield | | * | * | | * | | * | | * |
| Grinnell | | * | * | | * | | * | | * |
| Grinnell | | * | * | | * | | * | | * |
| Grundy Ctr | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Grundy Ctr | | * | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | | * | | * |
| Hampton | | * | * | | * | | * | | * |
| Hampton | | * | * | | * | | * | | * |
| Harlan | | * | * | | * | | * | | * |
| Harlan | | * | * | | * | | * | | * |
| Hawarden | | * | * | | * | | * | | * |
| Hawarden | | * | * | | * | | * | | * |
| Hispanic Cln | | * | * | | * | | * | | * |
| Hispanic Cln | | * | * | | * | | * | | * |
| Humboldt | | * | * | | * | | * | | * |
| Humboldt | | * | * | | * | | * | | * |
| ILH | | 33.7 (15) | 12.5 (21) | | 0.9 (53) | | 12.1 (43) | | 59.5 (45) |
| ILH | | * | * | | 50.2 (8) | | * | | 95.5 (8) |
| Ida Grove | | * | * | | * | | * | | * |
| Ida Grove | | * | * | | * | | * | | * |
| Independen | | * | * | | 0.0 (63) | | * | | * |
| Independen | | * | * | | * | | * | | * |
| Indianola | | * | * | | * | | * | | * |
| Indianola | | 24.2 (34) | 14.4 (27) | | 1.3 (43) | | 27.2 (4) | | 86.0 (23) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Iowa City | | 19.0 (40) | 18.6 (33) | | 1.9 (34) | | 19.1 (18) | | 76.2 (36) |
| Iowa City | | * | * | | 38.7 (17) | | 16.3 (31) | | 90.2 (13) |
| Iowa Falls | | * | * | | * | | * | | * |
| Iowa Falls | | * | * | | * | | * | | * |
| Jamaica | | * | * | | * | | * | | * |
| Jefferson | | * | * | | * | | * | | * |
| Jefferson | | * | * | | * | | * | | * |
| KR | | * | * | | * | | * | | * |
| KR | | * | * | | * | | * | | * |
| Keokuk | | * | * | | * | | * | | * |
| Keokuk | | 20.8 (38) | 10.7 (18) | | 1.2 (45) | | 19.1 (18) | | 90.8 (12) |
| Keosauqua | | * | * | | * | | * | | * |
| Keosauqua | | * | * | | * | | * | | * |
| Knoxville | | * | * | | * | | * | | * |
| Knoxville | | 24.5 (33) | 17.3 (30) | | 1.1 (47) | | 24.8 (7) | | 79.1 (30) |
| Lamoni | | * | * | | * | | * | | * |
| Lamoni | | * | * | | * | | * | | * |
| Le Mars | | * | * | | 0.0 (63) | | * | | * |
| Le Mars | | * | * | | * | | * | | * |
| Lenox | | * | * | | * | | * | | * |
| Lenox | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Leon | | * | * | | * | | * | | * |
| Leon | | * | * | | * | | * | | * |
| Logan | | * | * | | * | | * | | * |
| Logan | | * | * | | * | | * | | * |
| Manchester | | * | * | | * | | * | | * |
| Manchester | | * | * | | * | | * | | * |
| Maquoketa | | * | * | | * | | * | | * |
| Maquoketa | | * | * | | * | | * | | * |
| Marengo | | * | * | | * | | * | | * |
| Marengo | | * | * | | * | | * | | * |
| Marion | | * | * | | * | | * | | * |
| Marion | | 29.2 (28) | 13.7 (25) | | 0.8 (54) | | 24.7 (8) | | 78.8 (31) |
| Marshalltown | | 42.2 (3) | 10.3 (17) | | 1.1 (47) | | 11.2 (44) | | 70.0 (43) |
| Marshalltown | | * | * | | 60.5 (3) | | 13.2 (41) | | 96.1 (7) |
| Mason City | | 34.6 (13) | 24.8 (39) | | 1.8 (38) | | 23.2 (11) | | 82.2 (28) |
| Mason City | | * | * | | 58.1 (5) | | 22.8 (12) | | 94.4 (9) |
| Monticello | | * | * | | * | | * | | * |
| Monticello | | * | * | | * | | * | | * |
| Monticello | | * | * | | * | | * | | * |
| Mt Ayr | | * | * | | * | | * | | * |
| Mt Ayr | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Mt Pleasan | | 31.9 (19) | 9.2 (13) | | 2.5 (27) | | 22.3 (14) | | 85.3 (24) |
| Mt Pleasan | | * | * | | * | | * | | * |
| Murray | | * | * | | * | | * | | * |
| Muscatine | | * | * | | 66.4 (1) | | 14.8 (38) | | 87.6 (20) |
| Muscatine | | 38.5 (8) | 7.0 (6) | | 2.2 (30) | | 14.9 (37) | | 90.1 (14) |
| Nevada | | * | * | | * | | * | | * |
| Nevada | | * | * | | * | | * | | * |
| New Hampto | | * | * | | * | | * | | * |
| New Hampto | | * | * | | * | | * | | * |
| Newton | | 30.0 (24) | 6.3 (4) | | 2.1 (32) | | 15.0 (35) | | 80.7 (29) |
| Newton | | * | * | | * | | * | | * |
| North Towne | | 23.7 (35) | 9.7 (15) | | 0.7 (56) | | 16.9 (28) | | 76.8 (35) |
| North Towne | | * | * | | 50.5 (7) | | * | | * |
| Northwood | | * | * | | * | | * | | * |
| Northwood | | * | * | | * | | * | | * |
| Norwalk | | * | * | | * | | * | | * |
| Oelwein | | * | * | | * | | * | | * |
| Oelwein | | * | * | | * | | * | | * |
| Onawa | | * | * | | * | | * | | * |
| Onawa | | * | * | | * | | * | | * |
| Orange Cty | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Orange Cty | | * | * | | * | | * | | * |
| Osage | | * | * | | * | | * | | * |
| Osage | | * | * | | * | | * | | * |
| Osceola | | * | * | | 1.9 (34) | | * | | * |
| Osceola | | * | * | | * | | * | | * |
| Oskaloosa | | * | * | | * | | * | | * |
| Oskaloosa | | * | * | | 1.0 (49) | | * | | * |
| Oskaloosa | | * | * | | * | | * | | * |
| Ottumwa | | 30.6 (21) | 18.8 (34) | | 2.8 (24) | | 21.9 (15) | | 73.9 (40) |
| Ottumwa | | * | * | | 59.9 (4) | | 24.0 (9) | | 90.9 (11) |
| Pella | | * | * | | * | | * | | * |
| Pella | | * | * | | * | | * | | * |
| Perry | | * | * | | 1.0 (49) | | * | | * |
| Perry | | * | * | | * | | * | | * |
| Perry | | * | * | | * | | * | | * |
| Pocahontas | | * | * | | * | | * | | * |
| Pocahontas | | * | * | | * | | * | | * |
| Pocahontas | | * | * | | * | | * | | * |
| Pomeroy | | * | * | | * | | * | | * |
| Postville | | * | * | | * | | * | | * |
| Postville | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Red Oak | | 33.0 (17) | 2.0 (1) | | 0.0 (63) | | * | | * |
| Red Oak | | * | * | | * | | * | | * |
| Redfield | | * | * | | * | | * | | * |
| Refugee Cln | | * | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | | * | | * |
| Rock Valle | | * | * | | * | | * | | * |
| Rock Valle | | * | * | | * | | * | | * |
| Rockwell C | | * | * | | * | | * | | * |
| Rockwell C | | * | * | | * | | * | | * |
| SDM | | 34.8 (12) | 10.2 (16) | | 1.0 (49) | | * | | 90.0 (15) |
| Sac City | | * | * | | * | | * | | * |
| Sac City | | * | * | | * | | * | | * |
| Sheldon | | * | * | | * | | * | | * |
| Sheldon | | * | * | | 1.0 (49) | | * | | * |
| Shenandoah | | * | * | | * | | * | | * |
| Shenandoah | | * | * | | * | | * | | * |
| Sibley | | * | * | | * | | * | | * |
| Sibley | | * | * | | * | | * | | * |
| Sidney | | * | * | | * | | * | | * |
| Sidney | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Sigourney | | * | * | | * | | * | | * |
| Sigourney | | * | * | | * | | * | | * |
| Sioux Cent | | * | * | | 0.0 (63) | | * | | * |
| Sioux Cent | | * | * | | * | | * | | * |
| Sioux City | | 6.6 (45) | 91.8 (45) | | 36.7 (18) | | 20.9 (17) | | 78.6 (32) |
| Sioux City | | 30.5 (22) | 10.9 (19) | | 1.5 (39) | | 22.8 (12) | | 87.2 (21) |
| Southridge | | 26.2 (32) | 12.7 (22) | | 1.9 (34) | | * | | * |
| Spencer | | 27.9 (31) | 13.0 (24) | | 0.6 (58) | | 41.8 (1) | | 86.2 (22) |
| Spencer | | * | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | | * | | * |
| Storm Lake | | 15.8 (41) | 9.4 (14) | | 0.0 (63) | | 39.6 (2) | | 90.0 (15) |
| Storm Lake | | * | * | | * | | * | | * |
| Stuart | | * | * | | * | | * | | * |
| Stuart | | * | * | | * | | * | | * |
| Tama | | 30.0 (24) | 14.2 (26) | | 4.6 (20) | | * | | 74.4 (39) |
| Tama | | * | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | | * | | * |
| Tipton | | * | * | | * | | * | | * |
| Tipton | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Traer | | * | * | | * | | * | | * |
| Traer | | * | * | | * | | * | | * |
| UIHC | | * | * | | * | | * | | * |
| Vinton | | * | * | | * | | * | | * |
| Vinton | | * | * | | * | | * | | * |
| WDM | | 32.8 (18) | 8.6 (11) | | 2.4 (28) | | * | | * |
| WDM | | * | * | | * | | * | | * |
| Wapello | | * | * | | * | | * | | * |
| Wapello | | * | * | | * | | * | | * |
| Washington | | * | * | | * | | * | | * |
| Washington | | * | * | | * | | * | | * |
| Waterloo | | 38.9 (7) | 57.1 (40) | | 45.9 (13) | | 15.8 (32) | | 98.1 (2) |
| Waterloo | | 35.5 (10) | 19.5 (35) | | 2.3 (29) | | 17.0 (27) | | 83.1 (26) |
| Waukee | | * | * | | * | | * | | * |
| Waukee | | * | * | | * | | * | | * |
| Waukee | | * | * | | * | | * | | * |
| Waukon | | * | * | | * | | * | | * |
| Waukon | | * | * | | * | | * | | * |
| Waverly | | * | * | | * | | * | | * |
| Waverly | | * | * | | * | | * | | * |
| Wbstr Ct | | 39.5 (6) | 18.5 (32) | | 2.2 (30) | | 17.9 (24) | | 76.0 (37) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 6A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Maternal Behavioral Indicators by Clinic

### Combined 3 Years

**Comparison of Maternal Behavioral Indicators** (2,3)

| Clinic | Medical Care | | | WIC Enrollment | | Multivitamin Consumption | | | |
|---|---|---|---|---|---|---|---|---|---|
| | 1st Trimester | | No Care | 1st Trimester | | Prepregnancy (4) | | During Pregnancy (5) | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Wbstr Ct | | * | * | | * | | * | | * |
| West Union | | * | * | | * | | * | | * |
| Williamsburg | | * | * | | * | | * | | * |
| Winterset | | * | * | | * | | * | | * |
| Winterset | | * | * | | * | | * | | * |
| **Iowa** | | **30.2** | **20.0** | | **9.3** | | **19.7** | | **80.5** |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Women who reported taking a multivitamin four times a weeks or more in the month before pregnancy; proxy for folic acid consumption.
(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ackley | | * | | * | | * | | * | | * |
| Adel | | * | | * | | * | | * | | * |
| Adel | | * | | * | | * | | * | | * |
| Adel | | * | | * | | * | | * | | * |
| Albia | | * | | * | | * | | * | | * |
| Albia | | * | | * | | * | | * | | * |
| Algona | | * | | * | | * | | * | | * |
| Algona | | * | | * | | * | | * | | * |
| Allen WHC | | 30.2 (21) | | 14.3 (14) | | * | | 7.2 (20) | | 2.0 (18) |
| Allen WHC | | * | | * | | * | | * | | * |
| Allison | | * | | * | | * | | * | | * |
| Allison | | * | | * | | * | | * | | * |
| Altoona | | * | | * | | * | | * | | * |
| Altoona | | * | | * | | * | | * | | * |
| Ames | | 27.8 (17) | | 14.8 (15) | | 38.6 (4) | | 10.2 (33) | | 1.2 (9) |
| Ames | | 38.8 (45) | | * | | * | | * | | * |
| Anamosa | | * | | * | | * | | * | | * |
| Anamosa | | * | | * | | * | | * | | * |
| Anamosa | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ankeny | | * | | * | | * | | * | | * |
| Ankeny | | * | | * | | * | | * | | * |
| Atlantic | | * | | * | | * | | * | | * |
| Atlantic | | * | | * | | * | | * | | * |
| Audubon | | * | | * | | * | | * | | * |
| Audubon | | * | | * | | * | | * | | * |
| Avoca | | * | | * | | * | | * | | * |
| Avoca | | * | | * | | * | | * | | * |
| Bedford | | * | | * | | * | | * | | * |
| Bedford | | * | | * | | * | | * | | * |
| Belle Plaine | | * | | * | | * | | * | | * |
| Belle Plaine | | * | | * | | * | | * | | * |
| Bloomfield | | * | | * | | * | | * | | * |
| Bloomfield | | * | | * | | * | | * | | * |
| Boone | | * | | * | | * | | * | | * |
| Boone | | * | | * | | * | | * | | * |
| Brdlwns RP | | 25.5 (12) | | 15.8 (17) | | 22.8 (16) | | 6.1 (16) | | 2.5 (24) |
| Brdlwns RP | | 25.4 (11) | | * | | * | | 7.9 (24) | | * |
| Brooklyn | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Brooklyn | | * | | * | | * | | * | | * |
| Burlington | | 45.3 (54) | | 22.9 (37) | | 31.8 (7) | | 15.3 (52) | | 3.8 (36) |
| Burlington | | * | | * | | * | | * | | * |
| CHC | | 37.7 (40) | | 25.0 (43) | | 22.1 (17) | | 14.4 (49) | | 3.9 (37) |
| CHC | | 37.6 (39) | | * | | * | | 15.2 (51) | | * |
| CR Urban | | 33.2 (31) | | 18.7 (28) | | 36.2 (5) | | 9.2 (31) | | 1.3 (12) |
| CR Urban | | 43.8 (52) | | 16.0 (18) | | * | | 11.8 (41) | | 2.5 (24) |
| Carlisle | | * | | * | | * | | * | | * |
| Carlisle | | * | | * | | * | | * | | * |
| Carroll | | 40.3 (48) | | 21.9 (36) | | * | | 10.8 (36) | | 1.1 (8) |
| Carroll | | * | | * | | * | | * | | * |
| Cedar Falls | | * | | * | | * | | * | | * |
| Cedar Falls | | * | | * | | * | | * | | * |
| Centervill | | * | | * | | * | | * | | * |
| Centervill | | * | | * | | * | | * | | * |
| Chariton | | * | | * | | * | | * | | * |
| Chariton | | * | | * | | * | | * | | * |
| Chariton | | * | | * | | * | | * | | * |
| Charles City | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS
## Table 7A
## 2016 - 2018 Pregnancy Nutrition Surveillance (1)
## Iowa
### Comparison of Smoking Indicators by Clinic
### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Charles City | | 31.0 (25) | | 23.6 (40) | | * | | 4.7 (8) | | 1.4 (14) |
| Cherokee | | * | | * | | * | | * | | * |
| Cherokee | | * | | * | | * | | * | | * |
| Clarinda | | * | | * | | * | | * | | * |
| Clarinda | | * | | * | | * | | * | | * |
| Clarion | | * | | * | | * | | * | | * |
| Clarion | | * | | 14.0 (11) | | * | | 6.0 (15) | | 2.0 (18) |
| Clinton | | * | | * | | * | | * | | * |
| Clinton | | 39.0 (46) | | 25.5 (44) | | 26.0 (12) | | 5.3 (11) | | 2.1 (20) |
| Colfax | | * | | * | | * | | * | | * |
| Colfax | | * | | * | | * | | * | | * |
| Columbus J | | * | | * | | * | | * | | * |
| Columbus J | | * | | * | | * | | * | | * |
| Corning | | * | | * | | * | | * | | * |
| Corning | | * | | * | | * | | * | | * |
| Correctionvl | | * | | * | | * | | * | | * |
| Correctionvl | | * | | * | | * | | * | | * |
| Corydon | | * | | * | | * | | * | | * |
| Corydon | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Council Blfs | | 41.2 (49) | | 25.7 (45) | | * | | 5.7 (12) | | 2.7 (26) |
| Council Blfs | | 37.0 (37) | | 23.4 (39) | | 20.5 (18) | | 5.8 (13) | | 1.5 (15) |
| Cresco | | * | | * | | * | | * | | * |
| Cresco | | * | | * | | * | | * | | * |
| Creston | | 38.6 (44) | | 32.9 (49) | | * | | 15.6 (53) | | 3.3 (33) |
| Creston | | * | | * | | * | | * | | * |
| Dallas Ctr | | * | | * | | * | | * | | * |
| Decorah | | * | | * | | * | | * | | * |
| Decorah | | * | | * | | * | | * | | * |
| Denison | | 12.1 (3) | | 7.1 (3) | | * | | 3.5 (5) | | 0.5 (5) |
| Denison | | * | | * | | * | | * | | * |
| Dewitt | | * | | * | | * | | * | | * |
| Dewitt | | * | | * | | * | | * | | * |
| Dubuque | | 19.6 (8) | | 13.5 (10) | | * | | 0.6 (2) | | 0.0 (1) |
| Dubuque | | 14.6 (6) | | 9.1 (4) | | 25.8 (13) | | 0.1 (1) | | 0.0 (1) |
| Dyersville | | * | | * | | * | | * | | * |
| Dyersville | | * | | * | | * | | * | | * |
| Eagle Grove | | * | | * | | * | | * | | * |
| Eagle Grove | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indi ators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Edgerton | | 36.9 (36) | | 17.5 (22) | | * | | 14.6 (50) | | 5.2 (43) |
| Edgerton | | 31.6 (27) | | 15.7 (16) | | 29.9 (9) | | 11.7 (40) | | 2.8 (29) |
| Edgerton CHC | | * | | * | | * | | * | | * |
| Eldora | | * | | * | | * | | * | | * |
| Eldora | | * | | * | | * | | * | | * |
| Elkader | | * | | * | | * | | * | | * |
| Elkader | | * | | * | | * | | * | | * |
| Emmetsburg | | * | | * | | * | | * | | * |
| Emmetsburg | | * | | * | | * | | * | | * |
| Esthervill | | * | | * | | * | | * | | * |
| Esthervill | | 26.5 (14) | | 18.1 (26) | | * | | 13.7 (47) | | 7.0 (46) |
| Fairfiel | | * | | * | | * | | * | | * |
| Fairfiel | | * | | * | | * | | * | | * |
| Fayette | | * | | * | | * | | * | | * |
| Fayette | | * | | * | | * | | * | | * |
| Forest City | | * | | 18.6 (27) | | * | | * | | * |
| Forest City | | * | | * | | * | | * | | * |
| Fort Dodge | | 38.2 (43) | | 22.9 (37) | | 33.7 (6) | | 10.2 (33) | | 2.7 (26) |
| Fort Dodge | | 31.4 (26) | | * | | * | | 13.7 (47) | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ft Madison | | 43.8 (52) | | 20.3 (32) | | * | | 13.2 (45) | | 2.8 (29) |
| Ft Madison | | * | | * | | * | | * | | * |
| Garner | | * | | * | | * | | * | | * |
| Garner | | * | | * | | * | | * | | * |
| Glenwood | | * | | * | | * | | * | | * |
| Glenwood | | * | | * | | * | | * | | * |
| Grand Junction | | * | | * | | * | | * | | * |
| Greenfield | | * | | * | | * | | * | | * |
| Greenfield | | * | | * | | * | | * | | * |
| Grinnell | | * | | * | | * | | * | | * |
| Grinnell | | * | | * | | * | | * | | * |
| Grundy Ctr | | * | | * | | * | | * | | * |
| Grundy Ctr | | * | | * | | * | | * | | * |
| Guthrie Ct | | * | | * | | * | | * | | * |
| Guthrie Ct | | * | | * | | * | | * | | * |
| Hampton | | * | | 11.8 (7) | | * | | * | | 0.0 (1) |
| Hampton | | * | | * | | * | | * | | * |
| Harlan | | * | | * | | * | | * | | * |
| Harlan | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Hawarden | | * | | * | | * | | * | | * |
| Hawarden | | * | | * | | * | | * | | * |
| Hispanic Cln | | * | | * | | * | | * | | * |
| Hispanic Cln | | * | | * | | * | | * | | * |
| Humboldt | | * | | * | | * | | * | | * |
| Humboldt | | * | | * | | * | | * | | * |
| ILH | | 26.2 (13) | | 17.7 (23) | | 26.4 (11) | | 8.0 (26) | | 4.2 (41) |
| ILH | | 32.0 (28) | | * | | * | | 9.7 (32) | | * |
| Ida Grove | | * | | * | | * | | * | | * |
| Ida Grove | | * | | * | | * | | * | | * |
| Independen | | * | | * | | * | | * | | * |
| Independen | | * | | * | | * | | * | | * |
| Indianola | | * | | * | | * | | * | | * |
| Indianola | | 32.9 (30) | | 21.0 (35) | | * | | 11.9 (42) | | 7.2 (47) |
| Iowa City | | 13.3 (4) | | 6.8 (2) | | * | | 4.2 (7) | | 0.8 (7) |
| Iowa City | | 14.1 (5) | | * | | * | | 3.9 (6) | | * |
| Iowa Falls | | * | | * | | * | | * | | * |
| Iowa Falls | | * | | * | | * | | * | | * |
| Jamaica | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Jefferson | | * | | * | | * | | * | | * |
| Jefferson | | * | | * | | * | | * | | * |
| KR | | * | | * | | * | | * | | * |
| KR | | * | | * | | * | | * | | * |
| Keokuk | | * | | * | | * | | * | | * |
| Keokuk | | 51.6 (56) | | 31.5 (48) | | * | | 18.6 (55) | | 4.0 (39) |
| Keosauqua | | * | | * | | * | | * | | * |
| Keosauqua | | * | | * | | * | | * | | * |
| Knoxville | | * | | * | | * | | * | | * |
| Knoxville | | 38.1 (42) | | 28.9 (47) | | * | | 11.2 (38) | | 4.0 (39) |
| Lamoni | | * | | * | | * | | * | | * |
| Lamoni | | * | | * | | * | | * | | * |
| Le Mars | | * | | * | | * | | * | | * |
| Le Mars | | * | | * | | * | | * | | * |
| Lenox | | * | | * | | * | | * | | * |
| Lenox | | * | | * | | * | | * | | * |
| Leon | | * | | * | | * | | * | | * |
| Leon | | * | | * | | * | | * | | * |
| Logan | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Logan | | * | | * | | * | | * | | * |
| Manchester | | * | | * | | * | | * | | * |
| Manchester | | * | | * | | * | | * | | * |
| Maquoketa | | * | | * | | * | | * | | * |
| Maquoketa | | * | | * | | * | | * | | * |
| Marengo | | * | | * | | * | | * | | * |
| Marengo | | * | | * | | * | | * | | * |
| Marion | | * | | * | | * | | * | | * |
| Marion | | 35.0 (33) | | 19.0 (30) | | 40.3 (3) | | 7.4 (21) | | 1.9 (17) |
| Marshalltown | | 20.5 (9) | | * | | * | | 10.3 (35) | | * |
| Marshalltown | | 18.0 (7) | | 11.2 (6) | | 31.8 (7) | | 6.2 (17) | | 1.3 (12) |
| Mason City | | 43.5 (51) | | * | | * | | 4.8 (9) | | * |
| Mason City | | 37.4 (38) | | 24.1 (41) | | 25.0 (14) | | 8.0 (26) | | 1.2 (9) |
| Monticello | | * | | * | | * | | * | | * |
| Monticello | | * | | * | | * | | * | | * |
| Monticello | | * | | * | | * | | * | | * |
| Mt Ayr | | * | | * | | * | | * | | * |
| Mt Ayr | | * | | * | | * | | * | | * |
| Mt Pleasan | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indi ators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Mt Pleasan | | 37.9 (41) | | 20.1 (31) | | * | | 9.0 (29) | | 1.2 (9) |
| Murray | | * | | * | | * | | * | | * |
| Muscatine | | 30.7 (24) | | * | | * | | 16.5 (54) | | * |
| Muscatine | | 26.7 (15) | | 14.1 (12) | | 40.6 (2) | | 13.1 (44) | | 6.6 (45) |
| Nevada | | * | | * | | * | | * | | * |
| Nevada | | * | | * | | * | | * | | * |
| New Hampto | | * | | * | | * | | * | | * |
| New Hampto | | * | | * | | * | | * | | * |
| Newton | | * | | * | | * | | * | | * |
| Newton | | 35.2 (34) | | 25.9 (46) | | * | | 8.3 (28) | | 2.3 (23) |
| North Towne | | 30.2 (21) | | 16.3 (20) | | 46.7 (1) | | 6.8 (19) | | 1.8 (16) |
| North Towne | | * | | * | | * | | * | | * |
| Northwood | | * | | * | | * | | * | | * |
| Northwood | | * | | * | | * | | * | | * |
| Norwalk | | * | | * | | * | | * | | * |
| Oelwein | | * | | * | | * | | * | | * |
| Oelwein | | * | | * | | * | | * | | * |
| Onawa | | * | | * | | * | | * | | * |
| Onawa | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Orange Cty | | * | | * | | * | | * | | * |
| Orange Cty | | * | | * | | * | | * | | * |
| Osage | | * | | * | | * | | * | | * |
| Osage | | * | | * | | * | | * | | * |
| Osceola | | * | | * | | * | | * | | * |
| Osceola | | * | | * | | * | | * | | * |
| Oskaloosa | | * | | * | | * | | * | | * |
| Oskaloosa | | * | | 20.6 (33) | | * | | * | | * |
| Oskaloosa | | * | | * | | * | | * | | * |
| Ottumwa | | 40.1 (47) | | * | | * | | 18.6 (55) | | * |
| Ottumwa | | 28.0 (18) | | 18.8 (29) | | 15.8 (19) | | 13.4 (46) | | 2.7 (26) |
| Pella | | * | | * | | * | | * | | * |
| Pella | | * | | * | | * | | * | | * |
| Perry | | * | | 10.9 (5) | | * | | * | | 0.0 (1) |
| Perry | | * | | * | | * | | * | | * |
| Perry | | * | | * | | * | | * | | * |
| Pocahontas | | * | | * | | * | | * | | * |
| Pocahontas | | * | | * | | * | | * | | * |
| Pocahontas | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS
## Table 7A
## 2016 - 2018 Pregnancy Nutrition Surveillance (1)
## Iowa
### Comparison of Smoking Indicators by Clinic
### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Pomeroy | | * | | * | | * | | * | | * |
| Postville | | * | | * | | * | | * | | * |
| Postville | | * | | * | | * | | * | | * |
| Red Oak | | 47.6 (55) | | * | | * | | 7.7 (22) | | * |
| Red Oak | | * | | * | | * | | * | | * |
| Redfield | | * | | * | | * | | * | | * |
| Refugee Cln | | * | | * | | * | | * | | * |
| Rock Rapid | | * | | * | | * | | * | | * |
| Rock Rapid | | * | | * | | * | | * | | * |
| Rock Valle | | * | | * | | * | | * | | * |
| Rock Valle | | * | | * | | * | | * | | * |
| Rockwell C | | * | | * | | * | | * | | * |
| Rockwell C | | * | | * | | * | | * | | * |
| SDM | | 27.0 (16) | | 24.8 (42) | | * | | 7.8 (23) | | 3.9 (37) |
| Sac City | | * | | * | | * | | * | | * |
| Sac City | | * | | * | | * | | * | | * |
| Sheldon | | * | | * | | * | | * | | * |
| Sheldon | | * | | 14.2 (13) | | * | | * | | 6.0 (44) |
| Shenandoah | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Shenandoah | | * | | * | | * | | * | | * |
| Sibley | | * | | * | | * | | * | | * |
| Sibley | | * | | * | | * | | * | | * |
| Sidney | | * | | * | | * | | * | | * |
| Sidney | | * | | * | | * | | * | | * |
| Sigourney | | * | | * | | * | | * | | * |
| Sigourney | | * | | * | | * | | * | | * |
| Sioux Cent | | * | | * | | * | | * | | * |
| Sioux Cent | | 5.9 (1) | | * | | * | | 2.0 (3) | | * |
| Sioux City | | 32.1 (29) | | 17.9 (25) | | * | | 7.9 (24) | | 3.2 (32) |
| Sioux City | | 28.2 (19) | | 16.0 (18) | | 27.5 (10) | | 5.9 (14) | | 2.2 (22) |
| Southridge | | 29.5 (20) | | * | | * | | 6.7 (18) | | 3.4 (35) |
| Spencer | | 41.9 (50) | | 17.7 (23) | | * | | 12.6 (43) | | 7.8 (48) |
| Spencer | | * | | * | | * | | * | | * |
| Spirit Lak | | * | | * | | * | | * | | * |
| Spirit Lak | | * | | * | | * | | * | | * |
| Storm Lake | | 8.8 (2) | | 3.8 (1) | | * | | 2.9 (4) | | 4.2 (41) |
| Storm Lake | | * | | * | | * | | * | | * |
| Stuart | | * | | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indicators** (2,3)

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Stuart | | * | | * | | * | | * | | * |
| Tama | | 23.1 (10) | | 12.2 (8) | | * | | 5.0 (10) | | 3.3 (33) |
| Tama | | * | | * | | * | | * | | * |
| Tama/Settl | | * | | * | | * | | * | | * |
| Tama/Settl | | * | | * | | * | | * | | * |
| Tipton | | * | | * | | * | | * | | * |
| Tipton | | * | | * | | * | | * | | * |
| Traer | | * | | * | | * | | * | | * |
| Traer | | * | | * | | * | | * | | * |
| UIHC | | * | | * | | * | | * | | * |
| Vinton | | * | | * | | * | | * | | * |
| Vinton | | * | | * | | * | | * | | * |
| WDM | | * | | * | | * | | * | | * |
| WDM | | * | | * | | * | | * | | * |
| Wapello | | * | | * | | * | | * | | * |
| Wapello | | * | | * | | * | | * | | * |
| Washington | | * | | * | | * | | * | | * |
| Washington | | * | | * | | * | | * | | * |
| Waterloo | | 35.8 (35) | | 20.6 (33) | | 24.7 (15) | | 9.1 (30) | | 2.1 (20) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 7A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Smoking Indicators by Clinic

### Combined 3 Years

**Comparison of Smoking Indi ators (2,3)**

| Clinic | Smoking | | | | Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4) | | Smoking in Household | | | |
|---|---|---|---|---|---|---|---|---|---|---|
| | 3 Months Prior to Pregnancy | | Last 3 Months of Pregnancy | | | | Prenatal | | Postpartum | |
| | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Waterloo | | 34.9 (32) | | 17.1 (21) | | * | | 11.3 (39) | | 3.0 (31) |
| Waukee | | * | | * | | * | | * | | * |
| Waukee | | * | | * | | * | | * | | * |
| Waukee | | * | | * | | * | | * | | * |
| Waukon | | * | | * | | * | | * | | * |
| Waukon | | * | | * | | * | | * | | * |
| Waverly | | * | | * | | * | | * | | * |
| Waverly | | * | | * | | * | | * | | * |
| Wbstr Ct | | 30.4 (23) | | 12.8 (9) | | * | | 11.1 (37) | | 0.7 (6) |
| Wbstr Ct | | * | | * | | * | | * | | * |
| West Union | | * | | * | | * | | * | | * |
| Williamsburg | | * | | * | | * | | * | | * |
| Winterset | | * | | * | | * | | * | | * |
| Winterset | | * | | * | | * | | * | | * |
| **Iowa** | | **30.5** | | **18.3** | | **29.3** | | **8.4** | | **2.8** |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) Among women who smoked 3 months prior to pregnancy.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Infant Health Indicators by Clinic

## Combined 3 Years

### Comparison of Infant Health Indicators (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ackley | | * | * | | * | | * | | * |
| Adel | | * | * | | * | | * | | * |
| Adel | | * | * | | * | | * | | * |
| Adel | | * | * | | * | | * | | * |
| Albia | | * | * | | * | | * | | * |
| Albia | | * | * | | * | | * | | * |
| Algona | | * | * | | * | | * | | * |
| Algona | | * | * | | * | | * | | * |
| Allen WHC | | 5.9 (13) | 7.2 (13) | | 8.1 (9) | | 2.4 (14) | | 68.7 (38) |
| Allen WHC | | * | * | | * | | * | | * |
| Allison | | * | * | | * | | * | | * |
| Allison | | * | * | | * | | * | | * |
| Altoona | | * | * | | * | | * | | * |
| Altoona | | * | * | | * | | * | | * |
| Ames | | 7.4 (28) | 9.1 (38) | | 8.9 (13) | | 3.0 (25) | | 78.9 (9) |
| Ames | | * | * | | * | | * | | * |
| Anamosa | | * | * | | * | | * | | * |
| Anamosa | | * | * | | * | | * | | * |
| Anamosa | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

## Comparison of Infant Health Indicators by Clinic

## Combined 3 Years

### Comparison of Infant Health Indicators (2,3)

| Clinic | Birthweight | | | Preterm | | Full Term LBW | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | (6) | | (7) | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ankeny | | * | * | | * | | * | | * |
| Ankeny | | 5.2 (9) | 16.5 (58) | | * | | * | | 84.3 (1) |
| Atlantic | | * | * | | * | | * | | * |
| Atlantic | | 5.1 (7) | 11.0 (49) | | 6.0 (1) | | 1.8 (6) | | 73.7 (24) |
| Audubon | | * | * | | * | | * | | * |
| Audubon | | * | * | | * | | * | | * |
| Avoca | | * | * | | * | | * | | * |
| Avoca | | * | * | | * | | * | | * |
| Bedford | | * | * | | * | | * | | * |
| Bedford | | * | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | | * | | * |
| Belle Plaine | | * | * | | * | | * | | * |
| Bloomfield | | * | * | | * | | * | | * |
| Bloomfield | | * | * | | * | | * | | * |
| Boone | | * | * | | * | | * | | * |
| Boone | | * | * | | * | | * | | * |
| Brdlwns RP | | 7.8 (32) | 7.5 (21) | | 9.6 (18) | | 3.3 (30) | | 80.4 (7) |
| Brdlwns RP | | * | * | | * | | * | | * |
| Brooklyn | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Brooklyn | | * | * | | * | | * | | * |
| Burlington | | 8.7 (46) | 9.9 (43) | | 9.1 (14) | | 3.7 (34) | | 60.9 (54) |
| Burlington | | * | * | | * | | * | | * |
| CHC | | * | * | | * | | * | | * |
| CHC | | 8.6 (45) | 6.4 (6) | | 10.4 (26) | | 3.2 (27) | | 61.3 (53) |
| CR Urban | | * | * | | * | | * | | * |
| CR Urban | | 7.7 (31) | 7.9 (25) | | 10.1 (21) | | 2.6 (18) | | 66.6 (42) |
| Carlisle | | * | * | | * | | * | | * |
| Carlisle | | * | * | | * | | * | | * |
| Carroll | | 7.5 (29) | 7.5 (21) | | 11.4 (36) | | 3.7 (34) | | 69.4 (33) |
| Carroll | | * | * | | * | | * | | * |
| Cedar Falls | | * | * | | * | | * | | * |
| Cedar Falls | | * | * | | * | | * | | * |
| Centervill | | * | * | | * | | * | | * |
| Centervill | | * | * | | * | | * | | * |
| Chariton | | * | * | | * | | * | | * |
| Chariton | | * | * | | * | | * | | * |
| Chariton | | * | * | | * | | * | | * |
| Charles City | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Charles City | | 10.9 (56) | 11.5 (51) | | 12.8 (44) | | 5.2 (43) | | 67.3 (40) |
| Cherokee | | 8.9 (49) | 5.0 (3) | | * | | * | | 74.3 (23) |
| Cherokee | | * | * | | * | | * | | * |
| Clarinda | | * | * | | * | | * | | * |
| Clarinda | | * | * | | * | | * | | * |
| Clarion | | 2.7 (1) | 7.3 (15) | | * | | * | | 74.5 (21) |
| Clarion | | * | * | | * | | * | | * |
| Clinton | | 10.1 (54) | 8.8 (34) | | 11.7 (38) | | 3.3 (30) | | 68.8 (36) |
| Clinton | | * | * | | * | | * | | * |
| Colfax | | * | * | | * | | * | | * |
| Colfax | | * | * | | * | | * | | * |
| Columbus J | | * | * | | * | | * | | * |
| Columbus J | | * | * | | * | | * | | * |
| Corning | | * | * | | * | | * | | * |
| Corning | | * | * | | * | | * | | * |
| Correctionvl | | * | * | | * | | * | | * |
| Correctionvl | | * | * | | * | | * | | * |
| Corydon | | * | * | | * | | * | | * |
| Corydon | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Council Blfs | | 7.5 (29) | 7.8 (24) | | 12.0 (40) | | 1.9 (7) | | 75.3 (17) |
| Council Blfs | | 8.3 (41) | 9.3 (39) | | 14.9 (47) | | 1.2 (4) | | 72.8 (25) |
| Cresco | | * | * | | * | | * | | * |
| Cresco | | * | * | | * | | * | | * |
| Creston | | 8.3 (41) | 7.4 (19) | | 10.5 (27) | | 3.9 (37) | | 63.8 (50) |
| Creston | | * | * | | * | | * | | * |
| Dallas Ctr | | * | * | | * | | * | | * |
| Decorah | | * | * | | * | | * | | * |
| Decorah | | * | * | | * | | * | | * |
| Denison | | * | * | | * | | * | | * |
| Denison | | 8.0 (37) | 7.3 (15) | | 12.6 (42) | | 1.1 (3) | | 68.8 (36) |
| Dewitt | | * | * | | * | | * | | * |
| Dewitt | | * | * | | * | | * | | * |
| Dubuque | | 6.9 (24) | 11.9 (53) | | * | | * | | 51.5 (58) |
| Dubuque | | 6.4 (19) | 9.4 (41) | | 6.5 (3) | | 2.3 (13) | | 57.1 (56) |
| Dyersville | | * | * | | * | | * | | * |
| Dyersville | | * | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | | * | | * |
| Eagle Grove | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm | | Full Term LBW | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | (6) | | (7) | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Edgerton | | * | * | | * | | * | | 59.2 (55) |
| Edgerton | | 9.9 (52) | 7.4 (19) | | 10.2 (23) | | 5.2 (43) | | 63.5 (51) |
| Edgerton CHC | | * | * | | * | | * | | * |
| Eldora | | * | * | | * | | * | | * |
| Eldora | | * | * | | * | | * | | * |
| Elkader | | * | * | | * | | * | | * |
| Elkader | | * | * | | * | | * | | * |
| Emmetsburg | | * | * | | * | | * | | * |
| Emmetsburg | | * | * | | * | | * | | * |
| Esthervill | | * | * | | * | | * | | * |
| Esthervill | | 11.6 (57) | 8.0 (26) | | * | | * | | 66.1 (44) |
| Fairfiel | | * | * | | * | | * | | * |
| Fairfiel | | * | * | | * | | * | | * |
| Fayette | | * | * | | * | | * | | * |
| Fayette | | * | * | | * | | * | | * |
| Forest City | | * | * | | * | | * | | * |
| Forest City | | * | * | | * | | * | | * |
| Fort Dodge | | * | * | | * | | * | | * |
| Fort Dodge | | 7.2 (27) | 8.5 (30) | | 6.3 (2) | | 3.2 (27) | | 74.8 (19) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm | | Full Term LBW | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | (6) | | (7) | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Ft Madison | | * | * | | * | | * | | * |
| Ft Madison | | 8.2 (38) | 8.7 (33) | | 7.2 (5) | | 4.2 (40) | | 55.1 (57) |
| Garner | | * | * | | * | | * | | * |
| Garner | | * | * | | * | | * | | * |
| Glenwood | | * | * | | * | | * | | * |
| Glenwood | | * | * | | * | | * | | * |
| Grand Junction | | * | * | | * | | * | | * |
| Greenfield | | * | * | | * | | * | | * |
| Greenfield | | * | * | | * | | * | | * |
| Grinnell | | * | * | | * | | * | | * |
| Grinnell | | 7.1 (25) | 7.1 (12) | | * | | * | | 75.0 (18) |
| Grundy Ctr | | * | * | | * | | * | | * |
| Grundy Ctr | | * | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | | * | | * |
| Guthrie Ct | | * | * | | * | | * | | * |
| Hampton | | 4.5 (4) | 9.0 (36) | | * | | * | | 75.9 (14) |
| Hampton | | * | * | | * | | * | | * |
| Harlan | | * | * | | * | | * | | * |
| Harlan | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Hawarden | | * | * | | * | | * | | * |
| Hawarden | | * | * | | * | | * | | * |
| Hispanic Cln | | * | * | | * | | * | | * |
| Hispanic Cln | | * | * | | * | | * | | * |
| Humboldt | | * | * | | * | | * | | * |
| Humboldt | | * | * | | * | | * | | * |
| ILH | | * | * | | * | | * | | * |
| ILH | | 8.7 (46) | 8.8 (34) | | 11.6 (37) | | 3.0 (25) | | 74.4 (22) |
| Ida Grove | | * | * | | * | | * | | * |
| Ida Grove | | * | * | | * | | * | | * |
| Independen | | * | * | | * | | * | | * |
| Independen | | 4.8 (5) | 10.5 (46) | | * | | * | | 69.5 (29) |
| Indianola | | * | * | | * | | * | | * |
| Indianola | | 6.0 (15) | 12.0 (55) | | 7.2 (5) | | 2.8 (24) | | 78.1 (10) |
| Iowa City | | 10.0 (53) | 6.9 (9) | | 10.5 (27) | | 4.0 (39) | | 82.0 (2) |
| Iowa City | | * | * | | * | | * | | * |
| Iowa Falls | | * | * | | * | | * | | * |
| Iowa Falls | | * | * | | * | | * | | * |
| Jamaica | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm | | Full Term LBW | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | (6) | | (7) | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Jefferson | | * | * | | * | | * | | * |
| Jefferson | | * | * | | * | | * | | * |
| KR | | * | * | | * | | * | | * |
| KR | | * | * | | * | | * | | * |
| Keokuk | | * | * | | * | | * | | * |
| Keokuk | | 5.0 (6) | 7.2 (13) | | 6.8 (4) | | 3.3 (30) | | 44.4 (59) |
| Keosauqua | | * | * | | * | | * | | * |
| Keosauqua | | * | * | | * | | * | | * |
| Knoxville | | * | * | | * | | * | | * |
| Knoxville | | 7.8 (32) | 7.3 (15) | | 11.1 (35) | | 2.5 (15) | | 69.5 (29) |
| Lamoni | | * | * | | * | | * | | * |
| Lamoni | | * | * | | * | | * | | * |
| Le Mars | | * | * | | * | | * | | * |
| Le Mars | | 6.2 (16) | 6.9 (9) | | 13.0 (45) | | 2.0 (8) | | 77.7 (11) |
| Lenox | | * | * | | * | | * | | * |
| Lenox | | * | * | | * | | * | | * |
| Leon | | * | * | | * | | * | | * |
| Leon | | * | * | | * | | * | | * |
| Logan | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm | | Full Term LBW | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | (6) | | (7) | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Logan | | * | * | | * | | * | | * |
| Manchester | | * | * | | * | | * | | * |
| Manchester | | * | * | | * | | * | | * |
| Maquoketa | | * | * | | * | | * | | * |
| Maquoketa | | * | * | | * | | * | | * |
| Marengo | | * | * | | * | | * | | * |
| Marengo | | * | * | | * | | * | | * |
| Marion | | * | * | | * | | * | | * |
| Marion | | 7.1 (25) | 10.6 (48) | | 10.9 (33) | | 2.1 (10) | | 69.5 (29) |
| Marshalltown | | * | * | | * | | * | | * |
| Marshalltown | | 6.2 (16) | 9.0 (36) | | 8.3 (12) | | 2.7 (20) | | 67.1 (41) |
| Mason City | | 5.1 (7) | 10.5 (46) | | 9.3 (15) | | 2.5 (15) | | 69.5 (29) |
| Mason City | | * | * | | * | | * | | * |
| Monticello | | * | * | | * | | * | | * |
| Monticello | | * | * | | * | | * | | * |
| Monticello | | * | * | | * | | * | | * |
| Mt Ayr | | * | * | | * | | * | | * |
| Mt Ayr | | * | * | | * | | * | | * |
| Mt Pleasan | | 8.5 (44) | 10.2 (44) | | 7.9 (8) | | 4.3 (41) | | 65.5 (45) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Mt Pleasan | | * | * | | * | | * | | * |
| Murray | | * | * | | * | | * | | * |
| Muscatine | | 6.5 (20) | 6.9 (9) | | 10.6 (30) | | 2.6 (18) | | 74.6 (20) |
| Muscatine | | * | * | | * | | * | | * |
| Nevada | | * | * | | * | | * | | * |
| Nevada | | * | * | | * | | * | | * |
| New Hampto | | * | * | | * | | * | | * |
| New Hampto | | * | * | | * | | * | | * |
| Newton | | * | * | | * | | * | | * |
| Newton | | 10.4 (55) | 10.4 (45) | | 10.8 (32) | | 3.9 (37) | | 64.2 (49) |
| North Towne | | 8.2 (38) | 8.5 (30) | | 9.5 (17) | | 2.7 (20) | | 70.7 (28) |
| North Towne | | * | * | | * | | * | | * |
| Northwood | | * | * | | * | | * | | * |
| Northwood | | * | * | | * | | * | | * |
| Norwalk | | * | * | | * | | * | | * |
| Oelwein | | * | * | | * | | * | | * |
| Oelwein | | * | * | | * | | * | | * |
| Onawa | | * | * | | * | | * | | * |
| Onawa | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm | | Full Term LBW | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | (6) | | (7) | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Orange Cty | | * | * | | * | | * | | * |
| Orange Cty | | * | * | | * | | * | | * |
| Osage | | * | * | | * | | * | | * |
| Osage | | * | * | | * | | * | | * |
| Osceola | | * | * | | * | | * | | * |
| Osceola | | 5.8 (12) | 4.2 (2) | | * | | * | | 77.5 (12) |
| Oskaloosa | | * | * | | * | | * | | * |
| Oskaloosa | | 9.4 (50) | 12.3 (56) | | 11.0 (34) | | * | | 64.5 (48) |
| Oskaloosa | | * | * | | * | | * | | * |
| Ottumwa | | 6.6 (21) | 8.1 (28) | | 10.5 (27) | | 2.1 (10) | | 71.9 (27) |
| Ottumwa | | * | * | | * | | * | | * |
| Pella | | * | * | | * | | * | | * |
| Pella | | * | * | | * | | * | | * |
| Perry | | * | * | | * | | * | | * |
| Perry | | 12.9 (58) | 11.9 (53) | | 11.9 (39) | | * | | 69.3 (34) |
| Perry | | * | * | | * | | * | | * |
| Pocahontas | | * | * | | * | | * | | * |
| Pocahontas | | * | * | | * | | * | | * |
| Pocahontas | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Pomeroy | | * | * | | * | | * | | * |
| Postville | | * | * | | * | | * | | * |
| Postville | | 3.8 (2) | 5.7 (4) | | * | | * | | 81.0 (4) |
| Red Oak | | 5.9 (13) | 13.3 (57) | | 15.3 (48) | | 0.0 (1) | | 64.7 (47) |
| Red Oak | | * | * | | * | | * | | * |
| Redfield | | * | * | | * | | * | | * |
| Refugee Cln | | * | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | | * | | * |
| Rock Rapid | | * | * | | * | | * | | * |
| Rock Valle | | * | * | | * | | * | | * |
| Rock Valle | | * | * | | * | | * | | * |
| Rockwell C | | * | * | | * | | * | | * |
| Rockwell C | | * | * | | * | | * | | * |
| SDM | | 7.8 (32) | 9.3 (39) | | 10.0 (20) | | 2.7 (20) | | 76.1 (13) |
| Sac City | | * | * | | * | | * | | * |
| Sac City | | * | * | | * | | * | | * |
| Sheldon | | 5.7 (11) | 7.3 (15) | | 14.4 (46) | | * | | 75.8 (15) |
| Sheldon | | * | * | | * | | * | | * |
| Shenandoah | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Shenandoah | | * | * | | * | | * | | * |
| Sibley | | * | * | | * | | * | | * |
| Sibley | | * | * | | * | | * | | * |
| Sidney | | * | * | | * | | * | | * |
| Sidney | | * | * | | * | | * | | * |
| Sigourney | | * | * | | * | | * | | * |
| Sigourney | | * | * | | * | | * | | * |
| Sioux Cent | | 6.2 (16) | 8.5 (30) | | 8.1 (9) | | 2.7 (20) | | 81.2 (3) |
| Sioux Cent | | * | * | | * | | * | | * |
| Sioux City | | 7.8 (32) | 6.8 (7) | | 12.7 (43) | | 1.6 (5) | | 68.1 (39) |
| Sioux City | | 6.7 (22) | 8.0 (26) | | 9.4 (16) | | 2.0 (8) | | 72.2 (26) |
| Southridge | | 8.4 (43) | 9.7 (42) | | 10.2 (23) | | 3.7 (34) | | 80.4 (7) |
| Spencer | | 3.9 (3) | 6.2 (5) | | 8.1 (9) | | 0.0 (1) | | 75.4 (16) |
| Spencer | | * | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | | * | | * |
| Spirit Lak | | * | * | | * | | * | | * |
| Storm Lake | | * | * | | * | | * | | * |
| Storm Lake | | 5.2 (9) | 8.1 (28) | | 7.5 (7) | | 2.5 (15) | | 65.4 (46) |
| Stuart | | * | * | | * | | * | | * |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm (6) | | Full Term LBW (7) | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | | | | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Stuart | | * | * | | * | | * | | * |
| Tama | | 8.8 (48) | 11.3 (50) | | 12.5 (41) | | 3.2 (27) | | 66.5 (43) |
| Tama | | * | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | | * | | * |
| Tama/Settl | | * | * | | * | | * | | * |
| Tipton | | * | * | | * | | * | | * |
| Tipton | | * | * | | * | | * | | * |
| Traer | | * | * | | * | | * | | * |
| Traer | | * | * | | * | | * | | * |
| UIHC | | * | * | | * | | * | | * |
| Vinton | | * | * | | * | | * | | * |
| Vinton | | * | * | | * | | * | | * |
| WDM | | 7.8 (32) | 11.7 (52) | | 10.1 (21) | | 2.1 (10) | | 80.8 (6) |
| WDM | | * | * | | * | | * | | * |
| Wapello | | * | * | | * | | * | | * |
| Wapello | | * | * | | * | | * | | * |
| Washington | | * | * | | * | | * | | * |
| Washington | | * | * | | * | | * | | * |
| Waterloo | | 8.2 (38) | 7.6 (23) | | 9.6 (18) | | 3.4 (33) | | 69.0 (35) |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.

# WIC PNSS & PedNSS

## Table 8A

## 2016 - 2018 Pregnancy Nutrition Surveillance (1)

## Iowa

### Comparison of Infant Health Indicators by Clinic

### Combined 3 Years

**Comparison of Infant Health Indicators** (2,3)

| Clinic | Birthweight | | | Preterm | | Full Term LBW | | Ever Breastfed | |
|---|---|---|---|---|---|---|---|---|---|
| | | Low (4) | High (5) | (6) | | (7) | | | |
| | Number | % (rank) | % (rank) | Number | % (rank) | Number | % (rank) | Number | % (rank) |
| Waterloo | | 9.8 (51) | 3.8 (1) | | 10.6 (30) | | 5.2 (43) | | 61.5 (52) |
| Waukee | | * | * | | * | | * | | * |
| Waukee | | * | * | | * | | * | | * |
| Waukee | | * | * | | * | | * | | * |
| Waukon | | * | * | | * | | * | | * |
| Waukon | | * | * | | * | | * | | * |
| Waverly | | * | * | | * | | * | | * |
| Waverly | | * | * | | * | | * | | * |
| Wbstr Ct | | 6.8 (23) | 6.8 (7) | | 10.2 (23) | | 4.3 (41) | | 81.0 (4) |
| Wbstr Ct | | * | * | | * | | * | | * |
| West Union | | * | * | | * | | * | | * |
| Williamsburg | | * | * | | * | | * | | * |
| Winterset | | * | * | | * | | * | | * |
| Winterset | | * | * | | * | | * | | * |
| **Iowa** | | **7.6** | **8.3** | | **10.0** | | **2.9** | | **67.6** |

(1) Reporting period is January 1 through December 31.
(2) Excludes records with unknown data or errors.
(3) Rank compares one Clinic's rate to other Clinics. Rank 1 = best rate.
(4) <2500 g.
(5) >4000 g.
(6) <37 weeks gestation.
(7) 37 or more weeks gestation and <2500 g.
* Percentages are not calculated if <100 records are available for analysis after exclusions.